Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

| | | 1 | |
|---|--|---|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| • | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

aZ5076

A

United States - AIUST Department of Agriculture

Forest Service

Bibliographies and Literature of Agriculture No. 10

A

Rooting Habits of Selected Commercial Tree Species of the Eastern United States— A Bibliography

MATTL SEP 23 '93'
SEP 23 '93'
SATTL SEP 23 '93'
SEP 23

EA last

USDA, National Agricultural Library
NAL Bldg
10301 Baltimore Blvd
Belteville, MD 20705-2351

Rooting Habits of Selected Commercial Tree Species of the Eastern United States— A Bibliography

Compiled by Penninah Smith and Leanne Every

Formerly, on Intergovernmental Personnel Act assignment to the Eastern Region, from the University of Illinois; and research technician, Northeastern Forest Experiment Station, respectively.

U.S. Department of Agriculture

Forest Service

Bibliographies and Literature of Agriculture No. 10

November 1980

Contents

| | Pa | ge |
|---------------|----|-----|
| Introduction | | iii |
| Bibliography | | 1 |
| Appendix | | 38 |
| Key | | 38 |
| Subject index | | 39 |

Acknowledgments

We would like to acknowledge Robert Blomquist, regional silviculturist, Eastern Region, Forest Service, U.S. Department of Agriculture, for conceiving and initiating this project; Dr. Irving Holland, department head of the Forestry School at the University of Illinois, for cooperation in the project that furnished office space and supplies and full library privileges; and the Forestry School faculty.

Introduction

A great variety of research has been conducted to study the rooting habits of trees. However, little of this literature has been synthesized to describe the rooting habits of particular tree species or groups of species in ways that would be useful to field foresters. The objective of this bibliography is to facilitate review of literature pertaining to rooting habits of commercially important tree species occurring in the Eastern United States, especially those that occur in the Northeast (Region 9, Forest Service, U.S. Department of Agriculture).

We have attempted to be thorough, but the result is not complete. This list focuses on the years after 1930. The literature on the genus *Pinus* has been searched more thoroughly than the other genera. Aspects of disease and mycorrhizae that do not pertain directly to root character are not emphasized. Horticultural species, for the most part, have been excluded. References to commonly planted exotic species, such as *Pinus sylvestris* L., have been included, especially when the research was conducted in the Eastern United States or the species was judged to have commercial importance there.

The entries are primarily from Biological Abstracts, Forestry Abstracts, and references cited in articles. If a paper has been abstracted in Biological or Forestry Abstracts, the abstract volume and article number are given with "FA" or "BA" at the end of the entry.

The titles of journals and publishing agencies are abbreviated according to the International List of Periodical Title Word Abbreviations (ISO 833–1974 E) of the International Organization for Standardization, with certain exceptions (e.g. the use of USDA for U.S. Dep. Agric.). Journal titles thus appear approximately as they are shown in the annual BIOSIS List of Serials (BioSciences Information Services, Philadelphia), where the full titles, CODEN symbols, countries of origin, and information about mergers, changes of title, and the like may be found.

The bibliography is arranged alphabetically by the author, and the entries are numbered consecutively. A subject index is given in the appendix. To find articles on root grafting of white pine, for example, first find the symbol for root grafting (IIIA-2) in the key to the subject index. Then turn to *Pinus strobus* in the species column of the subject index and find IIIA-2 in the subject columns. The 12 numbers listed under IIIA-2 refer to entries in the bibliography that concern white pine root grafting.

The subjects covered in the bibliography are outlined in the following list, which is the basis for the key to the subject index.

- I. Root growth and development
 - A. Form and anatomy
 - B. Relation to top growth and development
 - C. Genetic variation, within species
 - D. Periodic, seasonal
 - E. Influenced by:
 - 1. Root environment
 - a. Moisture
 - i. Drought
 - ii. Flood
 - b. Temperature
 - c. Soil
 - i. Physical aspects
 - ii. Chemical aspects
 - iii. Microrelief
 - 2. Shoot
 - a. Light
 - b. Hormones
- II. Root functions
 - A. Uptake and upward translocation
 - 1. Water
 - 2. Nutrients
 - 3. Other substances
 - B. Translocation downward
 - 1. Carbohydrate reservation
 - 2. Exudation
 - C. Respiration
 - D. Anchorage

- III. Root interactions, biological
 - A. With herbaceous and woody plants
 - 1. Spatial relations (competition for nutrients and moisture)
 - 2. (Natural) root grafting
 - 3. Soil amelioration
 - a. Exudation
 - b. Root death
 - 4. Allelopathy
 - B. With other than herbaceous and woody plants
 - 1. Symbiotic relations
 - a. Mycorrhizae
 - i. Ecto-
 - ii. Endo-
 - iii. Eclendo-
 - b. Nitrogen fixation
 - i. Leguminous
 - ii. Non-leguminous
 - 2. Pathological relationships
 - a. With undetermined or mechanical agent
 - b. With determined agent
 - i. Insect
 - ii. Nematode
 - iii. Fungus
 - iv. Bacterium
- IV. Root response to nursery and forestry practices

Bibliography

1. Abrazhko, V.I.

1969. The suction power of roots as an indicator of intensity of competiton between trees and undergrowth for soil moisture [O sosushchey sile korney kak pokazatele napryazhennosti sorevnovaniya mezhdy derev'yami i podrostom iz-za pochvennoy vlagi]. Transl., Can. Dep. Sec. State Transl. Bur., Foreign Lang. Div. *from* Bot. Zh. 53(2):254–259 (1968). FA32:339 (Transl. Dep. Fish. For. Can. No. 00FF–61).

2. Adams, W.R., Jr.

1928. Effect of spacing in a jack pine plantation. Vt. Agric. Exp. Stn. Bull. 282.

3. Adams, W.R., Jr.

1934. Studies in tolerance of New England forest trees. XI. The influence of soil temperature on the germination and development of white pine seedlings. Vt. Agric. Exp. Stn. Bull. 379.

4. Adams, W.R., and G.L. Chapman.

1941. Competition influence on the root systems of jack and Norway pines. Vt. Agric. Exp. Stn. Bull. 472. FA4:218.

5. Adams, W.T., J.H. Roberds, and B.J. Zobel.

1973. Intergenotypic interactions among families of loblolly pine. (*Pinus taeda* L.). Theor. and Appl. Genet. 43(7):319–322.

6. Addoms, R.M.

1946. Entrance of water into suberized roots of trees. Plant Physiol. 21(1):109–111. FA8:897.

7. Addoms, R.M.

1950. Notes on the structure of elongating pine roots. Am. J. Bot. 37(3):208–211. BA24:33816.

8. Agnihotri, V.P., and O. Vaartaja.

1967. Root exudates from red pine seedlings and their effects on *Pythium ultimum*. Can. J. Bot. 45(7):1031–1040. FA29:4086.

9. Ahlgren, C.E.

1958. Adventitious roots and shoots of wilding white pine at the Quetico-Superior Wilderness Research Center. Minn. For. Notes 70.

10. Ahlgren, C.E.

1959. Some effects of fire on forest reproduction in northeastern Minnesota. J. For. 57(3):194–200.

11. Ahlgren, C.E.

1972. Some effects of inter- and intraspecific grafting on growth and flowering of some five-needle pines. Silvae Genet. 21(3/4):122–126. FA34:1637.

12. Ahlgren, C.E., and H.L. Hansen.

1957. Some effects of temporary flooding on coniferous trees. J. For. 55(9):647–650.

13. Aikman, J.M., A.L. McComb, and G.B. Cooper.

1948. Investigation of planting procedures and cultural practices in growing trees on steep, eroded soils for post, woodlot, and wildlife purposes. Ia. Agric. Exp. Stn. Rep. 1948:194–195. FA11:1029.

14. Aird, P.L.

1958. The effects of herbaceous vegetation on growth of planted poplar. Diss. Abstr. 18(3):739–740. FA19:4166.

15. Albertson, F.W., and J.E. Weaver.

1945. Injury and death or recovery of trees in prairie climate. Ecol. Monogr. 15(4):393–433. BA20:9627.

16. Alexander, S.A.

1973. Root depth and percentage infection by *Fomes annosus* in a loblolly pine plantation on two soil hazard types. (Abstr.) Phytopathology 63(7):801. FA35:1673.

17. Alexander, S.A., and J.M. Skelly.

1974. A comparison of isolation methods for determining the incidence of *Fomes annosus* in living loblolly pine. Eur. J. For. Pathol. 4(1):33–38. FA35:6177.

18. Alexander, S.A., J.M. Skelly, and C.L. Morris.

1975. Edaphic factors associated with the incidence and severity of disease caused by *Fomes annosus* in loblolly pine plantations in Virginia. Phytopathology 65(5):585–591. FA37:391.

19. Alexandrés, S.

1977. The inhibiting effect of harmful fertilizing on root formation in poplars in different soil types. For Abstr. 38(5):2131. *From* Dasika Hronika 176/177:26–34, 45 (1973).

20. Allaire, D., B. Bernier, and Y. LaFlamme.

1973. Microelement status of balsam fir in the Southern Laurentians. Can. J. For. Res. 3(3):379–386. BA57:54627.

21. Allen, J.C.

1950. Pine planting tests in the Copper Basin. J. Tenn. Acad. Sci. 25(3):199–216. FA12:2923.

22. Allen, R.M.

1969a. Contributions of tops and roots to variation in height growth of geographic sources of shortleaf pine. Silvae Genet. 18(1/2):38–39. BA51:45991.

23. Allen, R.M.

1969b. Racial variation in physiological characteristics of shortleaf pine roots. Silvae Genet. 18(1/2):40–43.

24. Allen, S.E.

1964. Nitrogen and phosphorus nutrition of loblolly pine, as shown by split-root techniques. Agron. Abstr. 1964:52.

25. Althen, F.W. von.

1968. Incompatability of black walnut and red pine. Bi-mon. Res. Notes 24(2):19.

26. American Phytopathological Society.

1963. Symposium on root diseases of forest trees, presented at the 54th Annual Meeting of the Am. Phytopathol. Soc. [Corvallis, Oregon, Aug. 1962]. Phytopathology 53(10):1120–1136. BA45:44661.

27. Amos, R.E.

1962. Longevity of *Ceratocystis fagacearum* (Bretz) Hunt in the presence of other fungi in roots of deep-girdled oak-wilt trees. (Abstr.) Phytopathology 52(2):162. BA38:23845.

28. Amos, R.E., and R.P. True.

1967. Longevity of *Ceratocystis fagacearum* in roots of deep-girdled oak-wilt trees in West Virginia. Phytopathology 57(10):1012–1015. BA49:26333.

29. Anderson, C.H., and E.G. Cheyney.

1934. Root development in seedlings in relation to soil texture. J. For. 32(1):32–34.

30. Anderson, N.A., and G.W. Anderson.

1963. White pine root rot at the Chittenden nursery. USDA For. Serv. Res. Note LS-26. FA25:3734.

31. Anderson N.A., and W.M. Bugbee.

1962. *Cylindrocladium* root rot of conifer seedlings in Minnesota. (Abstr.) Phytopathology 52(8):721. BA41:3653.

32. Anderson, N.A., D.W. French, and D.P. Taylor.

1962. Cylindrocladium root rot of conifers in Minnesota. For. Sci. 8(4):378–382. BA42:12036.

33. Anderson, R.L., and M. Berbee.

1976. Development of ectomycorrhizae on yellow birch from the Eveleth Nursery, Minnesota. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–18–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

34. Anderson, R.L., and D. Mosher.

1976. Abundance of ectomycorrhizae on red pine in Michigan State Nurseries. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–19–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

35. Anderson, R.L., and L.J. Wlazik.

1976a. Comparison of development of ectomycorrhizae on 1–4 red pine seedlings from Tourney Minnesota National Forest Nursery and adjacent land. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–20–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

36. Anderson, R.L., and L.J. Wlazik.

1976b. Ectomycorrhizae on four year-old red pine seedlings from Eveleth Nursery, Minnesota. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–9–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

37. Anderson, R.L., and L.J. Wlazik.

1976c. Occurrence of ectomycorrhizal fungus on white spruce at Eveleth and Tourney USFS Nurseries. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–16–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

38. Anderson, R.L., J.E. Borkenhagen, and L.J. Wlazik.
1976a. Comparison of ectomycorrhizae occurrence on pruned and unpruned roots on 3–0 red pine seedlings at Hayward State Nursery, Hayward, Wisconsin, 1976. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–17–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

39. Anderson, R.L., S.V. Krupa, P.R. Laidly, and E. Stewart. 1976b. Evaluation of ectomycorrhizae on red pine seedlings at Tourney and Eveleth National Forest Nurseries in the Lake States. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S-14-76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

40. Anderson, R.L., D. Mosher, and G.R. Ellis.
1976c. Endomycorrhizae development on black walnut seedlings from the Southern Michigan State Nursery. USDA For. Serv. For. Insect and Dis. Manage. Eval. Rep. S–10–76, Northeast. Area, State and Priv. For., Upper Darby, Pa.

41. Andresen, J.W.

1959. A study of pseudo-nanism in *Pinus rigida* Mill. Ecol. Monogr. 29(4):309–332.

42. Andrews, L.K.

1941. Effects of certain soil treatments on the development of loblolly pine nursery stock. J. For. 39(11):918–921. FA3:285.

43. Arend, J.L.

1948. Influences on red-cedar distribution in the Ozarks. USDA For. Serv. Note 58, South. For. Exp. Stn., New Orleans, La.

44. Arend, J.L.

1950. Influence of fire and soil on distribution of eastern red-cedar in the Ozarks. J. For. 48(2):129–130.

45. Armson, K.A.

1970. Soils, roots, and foresters. *In* Tree growth and forest soils. C.T. Youngberg, and C.B. Davey, eds., 3rd North Am. For. Soils Conf. Proc., N.C. State Univ. Raleigh, [August 1968], p. 513–522.

46. Armson, K.A.

1972. Distribution of conifer seedling roots in a nursery soil. For. Chron. 48(3):141–143. FA34:152.

Armson, K.A., and W.S. Millward.
 1970. A study of aeration of roots of two-year black spruce and one-year jack pine. Rep. Res. For. Bot. Glendon Hall Fac. For. Univ.

Toronto 1969/70:3. FA33(2):2217.

48. Armson, K.A., and S.R. Shea.

1970. Effect of soil texture on physical impedance of root growth.

Rep. Res. For. Bot. Glendon Hall Fac. For. Univ. Toronto 1969/
70:4. FA33(2):2218.

49. Armson, K.A., and H. Struik.

1968. The effect of oxygen levels in the growth of black spruce and red pine seedlings. Rep. Res. For, Bot. Glendon Hall Fac. For. Univ. Toronto 1967/68:4. FA31(1):338.

 Armson, K.A., and R. Van Den Driessche.
 1959. Natural root grafts in red pine (*Pinus resinosa* Ait.). For. Chron. 35(3):232–241. BA35:11536.

51. Armson, K.A., and J.R.M. Williams.
1960. The root development of red pine (*Pinus resinosa* Ait.) seedlings in relation to various soil conditions. For. Chron. 36(1):14–17.
BA35:64597.

52. Armstrong, W.

1968. Oxygen diffusion from the roots of woody species. Physiol. Plant. 21(3):539–543. BA49:123943.

Armstrong, W., and D.J. Read.
 1972. Some observations on oxygen transport in conifer seedlings.
 New Phytol. 71:55–62.

54. Arnett, J.D., Jr., and W. Witcher.
1974. Infection sites of *Fusarium solani* on yellow poplar. Plant Dis. Rep. 58(8):754–757. FA36:2117.

55. Artman, J.D.

1974. Greenhouse screening of loblolly pine of known parentage for resistance to *Fomes annosus*. Plant Dis. Rep. 58(5):409-411. FA35:7695

56. Artman, J.D., and E.L. Sharp.

1971. An inoculation test using *Peniphora gigantea* on stumps of eastern white pine. Plant Dis. Rep. 55(9):834–836. FA33:4672.

57. Ashby, W.C.

1960. Seedling growth and water uptake by *Tilia americana* at several root temperatures. Bot. Gaz. 121(4):228–233. BA36:18456.

58. Ashby, W.C.

1962. Root growth in American basswood. Ecology 43(2):336–339. BA39:24626.

Ashby, W.C., and J.N. Cummins.
 1968. Root growth and rooting of basswood. Bot. Gaz. 129(4):327–333. BA50:100479.

60. Ashe, W.W.

1915. Loblolly or North Carolina pine. N.C. Geol. and Econ. Surv. Bull. 24.

61. Baker, F.S.

1918. Aspen reproduction in relation to management. J. For. 16:389–398.

62. Baker, F.S.
1925. Aspen in the Central Rocky Mountain Region. USDA Dep.
Bull. 1291.

63. Baker, J.B.

1977. Tolerance of planted hardwoods to spring flooding. South. J. Appl. For. 1(3):23–25. BA65:8852.

64. Baker, J.B., and B.G. Blackmon.

1974. Biomass and nitrogen accumulation in a cottonwood plantation. Agron. Abstr. 1974:174.

65. Balatinecz, J.J., D.F. Forward, and R.G.S. Bidwell.
1966. Distribution of photoassimilated C¹⁴O₂ in young jack pine seedlings. Can. J. Bot. 44(3):362–364. FA28:139.

66. Baldwin, H.I.

1954. Needle blight in eastern white pine. Plant Dis. Rep. 38(10):725-727. FA16:3240.

67. Bannan, M.W.

1940. The root systems of Northern Ontario conifers growing in sand. Am. J. Bot. 27(2):108–114. BA14:9933.

68. Bannan, M.W.

1941a. Variability in wood structure in roots of native Ontario conifers. Bull. Torrey Bot. Cl. 68(3):173–194. BA15:13216.

69. Bannan, M.W.

1941b. Vascular rays and adventitious root formation in *Thuja* occidentalis L. Am. J. Bot. 28(6):457–463. BA15:19299.

70. Bannan, M.W.

1941c. Wood structure of *Thuja occidentalis*. Bot. Gaz. 103:295–309.

71. Bannan, M.W.

1942. Notes on the origin of adventitious roots in the native Ontario conifers. Am. J. Bot. 29(8):593–598. BA17:2517.

72. Bard, G.E.

1946. The mineral nutrient content of the foliage of forest trees on three soil types of varying limestone content. Proc. Soil Sci. Soc. Am. 10:419–422.

73. Barham, R.O., D.H. Marx, and J.L. Ruehle.

1974. Infection of ectomycorrhizal and nonmycorrhizal roots of shortleaf pine by nematodes and *Phytophthora cinnamomi*. Phytopathology 64(9):1260–1264. BA59:22142.

74. Barker, K.R., and G.D. Griffin.

1965. The distribution of nematodes in jack and red pine plantations and the parasitism of *Xiphinema americanum* on jack pine. Nematologica 11(1):33–34. FA27:2570.

75. Barnard, E.L., and J.R. Jorgensen.

1977. Respiration of field-grown loblolly pine roots as influenced by temperature and root type. Can. J. Bot. 55(6):740–743.

76. Barnes, B.V.

1959. Natural variation and clonal development of *Populus tremuloides* and *P. grandidentata* in Northern Lower Michigan. Diss. Abstr. 20(5):1511-1512.

77. Barnes, B.V.

1966. The clonal growth habit of American aspens. Ecology 47(3):439-447.

78. Barnes, R.L.

1958. Studies on physiology of isolated pine roots and root callus cultures. Diss. Abstr. 19(3):417–418. BA33:23431.

79. Barnes, R.L., and A.W. Naylor.

1959a. Effect of various nitrogen sources on growth of isolated roots of *Pinus serotina*. Physiol. Plant. 12(1):82–89. BA33:38899.

80. Barnes, R.L., and A.W. Naylor.

1959b. *In vitro* culture of pine roots and the use of *Pinus serotina* roots in metabolic studies. For. Sci. 5(2):158–168. BA33:46884.

81. Barnes, R.L., and A.W. Naylor.

1959c. Studies on the ornithine cycle in roots and callus tissues of *Pinus serotina* and *Pinus clausa*. Bot. Gaz. 121(2):63-69. BA35:54068.

82. Barney, C.W.

1951. Effects of soil temperature and light intensity on root growth of loblolly pine seedlings. Plant Physiol. 26(1):146–163. BA25:25250.

83. Barykina, R.P.

1958. Osobennosti obrazovaniya kornevykh otpryskov u beloi akatsii (*Robinia pseudoacacia* L.). [Features of the formation of suckers in *R. pseudoacacia*.] Byul. Mosk. O-va Ispy. Prir. Otd. Biol. 63(4):57–71. Ref. Zhur., Biol. 1959, No. 66578 (Transl.). BA47:119058, FA22:4324.

84. Basham, J.T.

1966. Heart rot of jack pine in Ontario. I. The occurrence of basidiomycetes and microfungi in defective and normal heartwood of living jack pine. Can. J. Bot. 44(3):275–295. FA28:802.

85. Basham, J.T.

1973. Heart rot of black spruce in Ontario. I. Stem rot, hidden rot and management considerations. Can. J. For. Res. 3(1):95-104. FA35:876.

86. Baskerville, G.L.

1966. Dry-matter production in immature balsam fir stands: Roots, lesser vegetation, and total stand. For. Sci. 12(1): 49–53. FA27:5605.

87. Bassett, J.R

1962. Soil moisture and loblolly pine growth relations on a perched water table site in Southeast Arkansas. Diss. Abstr. 22(7):2136. FA23:4875.

88. Bates, C.G.

1926. Some relations of plant ecology to silvicultural practice. Ecology 7(4):469–480.

89. Batra, L.R.

1974. Armillaria mellea on flowering dogwood (Cornus florida). Plant Dis. Rep. 58(8):719–721. FA36:3415.

90. Baxter, D.V.

1937. Devleopment and succession of forest fungi and diseases in forest plantations. Mich. Univ. Sch. For. Conserv. Circ. 1.

91. Baxter, P., and D. West.

1977a. The flow of water into fruit trees. I. Resistances to water flow through roots and stems. Ann. Appl. Biol. 87 (1):95–101.

92. Baxter, P., and D. West.

1977b. The flow of water into fruit trees. II. Water intake through a cut limb. Ann. Appl. Biol. 87(1):103–112.

93. Baxter, S.S.

1958. Subsurface problems of trees and utility structures. Trees Mag., Ohio 19(1):7–8. FA20:3199.

94. Beattie, D.

1976. A layman's introduction to mycorrhizae. U.S. Gov. Print. Off. 1976–641–313/5151.

95. Beaufait, W.R.

1955. Soil profile observations relating to drouth damage in black willow stands. J. For. 53(7):517.

96. Becking, J.H.

1961. A requirement of molybdenum for the symbiotic nitrogen fixation in alder (*Alnus glutinosa* Gaertn.). Plant and Soil 15(3):217–227. FA23:3298.

97. Becking, J.H.

1970. Plant-endophyte symbiosis in non-leguminous plants. Plant and Soil 32(3):611–654. FA32:323.

98. Beckman, C.H., and J.E. Kuntz.

1951. Translocation of poisons, dyes, and radioiodine, and its relation to oak wilt. (Abstr.) Phytopathology 41(1):2–3. FA12:3388.

99. Beeftink, H.H.

1951. Some observations on tamarack or eastern larch (*Larix laricina* DuRoi (Koch)) in Alberta. For. Chron. 27(1):38–39.

100. Bega, R.V.

1962. Tree killing by *Fomes annosus* in a genetics arboretum. Plant Dis. Rep. 46(2):107–110.

101. Bega, R.V.

1963. Fomes annosus. In Symposium of root diseases of forest trees. Invitational papers presented August 27, 1962, at the 54th Annual Meeting of the American Phytopathological Society at Corvallis, Oregon. Phytopathology 53(10):1120–1123.

102 Beineke, W.F.

1967. Genetic variation in the ability to withstand transplanting shock in loblolly pine (*Pinus taeda* L.). Diss. Abstr. Int. 27B(12, part 1):4197. FA29:1897.

103. Bell, D.T.

1975. Germination of *Quercus alba* L. following flood conditions. Ill. Agric. Exp. Stn. For. Res. Rep. 75–2.

104. Bell, D.T., and F.L. Johnson.

1974. Flood-caused tree mortality around Illinois reservoirs. Trans. Ill. State Acad. Sci. 67(1):28–37.

105. Bellefeuille, R.

1935. La reproduction des peuplements d'épinette noire dans les forêts du Nord-Québec. [The reproduction of the populations of black spruce in the forests of Northern Quebec.]. For. Chron. 11(4):323–340. [Engl. summ.]

106. Bendana, F.E.

1964. Effects of nodules on geotropism of roots. Diss. Abstr. 24(7):2663–2664. FA25:4652.

107. Bergdahl, D.R., and D.W. French.

1976. Needle droop: an abiotic disease of plantation red pine. Plant Dis. Rep. 60(6):472–476. FA38:932.

108. Bergman, F., and B. Haggstrom.

1976. Some important facts considering planting with rooted forest plants. Transl. by R.A. Hellenius. For. Chron. 52(6):266–273, from Saertr. ur Sver. Skogsvardsforbunds Tidskr. 6 (1973).

109. Berndt, H.W., and R.D. Gibbons.

1958. Root distribution of some native trees and understory plants growing on three sites within ponderosa pine watersheds in Colorado. USDA For. Serv., Rocky Mtn. For. and Range Exp. Stn., Fort Collins, Colo., Stn. Pap. 37.

110. Berry, C.R., and D.H. Marx.

1976. Sewage sludge and *Pisolithus tinctorius* ectomycorrhizae: Their effect on growth of pine seedlings. For. Sci. 22(3):351–358.

111. Berry, C.R., and D.H. Marx.

1977. Growth of loblolly pine seedlings in strip-mined kaolin spoil as influenced by sewage sludge. J. Environ. Qual. 6(4):379–381.

112. Berry, F.H.

1968. Spread of *Fomes annosus* root rot in thinned shortleaf pine plantations. USDA For. Serv. Res. Note NE-87. FA31:6667.

113. Berry, F.H., and T.W. Bretz.

1963. Attempts to transmit the oak-wilt fungus by soil and root inoculations. Plant Dis. Rep. 47(3):164. FA25:3759.

Beslow, D.T., E. Hacskaylo, and J.H. Melhuish, Jr.
 1970. Effects of environment on beaded root development in red maple. Bull. Torrey Bot. Club 97(5):248–252. BA52:28116.

 Bey, C.F., and R.E. Phares.
 [n.d. Circa 1969]. Seasonal growth pattern for five sources of black walnut. Cent. States For. Tree Improv. Conf. Proc. 6:44–47.

116. Beyer, L.E., and R.J. Hutnik. 1969. Acid and aluminum toxicity as related to strip-mine spoil banks in western Pennsylvania. Pa. State Univ. Res. Briefs 3:69–72.

117. Bibelriether, H.
1967. Root development of some tree species in relation to soil properties [Die Bewurzelung einiger Baumarten in Abhangigkeit von Bodeneigenschaften]. Transl., Can. Dep. Sec. State Transl. Bur., Foreign Lang. Div., from Allg. Forstz. 21/47:808–815 (1966). FA29:292 (Transl. Dep. For. Can. 95).

118. Bilan, M.V. 1960. Root development of loblolly pine seedlings in modified environments. Stephen F. Austin State Coll., Nacogdoches, Tex., Dep. For. Bull. 4. FA22:255.

Bilan, M.V.
 1962. Effect of planting date on regeneration and development of roots of loblolly pine seedlings. 13th Int. Union For. Res. Organ. Congr. Proc., [Vienna 1961], Part 2(1), Section 22–15. FA24:273.

 Bilan, M.V.
 1965. Initial root growth in loblolly pine. (Abstr.) Bull. Ecol. Soc. Am. 46(3):94. FA27:3558.

121. Bilan, M.V.
1966. Low temperature as a limiting factor of root growth in loblolly pine seedlings. (Abstr.) Bull. Ecol. Soc. Am. 47(3):103. FA29:1945.

122. Bilan, M.V. 1967a. Effect of low temperature on root elongation in loblolly pine seedlings. 14th Int. Union For. Res. Organ. Congr. Proc., [Munich 1967], Part IV, Section 23, p. 74–82. FA29:2004.

123. Bilan, M.V.
1967b. Growth and development of root system in loblolly pine during the first season of growth. J. For. 65(3):224.

124. Bilan, M.V. 1968. Effects of physical soil environment on growth and development of root systems in southern pines. *In* Forest fertilization, theory and practice. [Papers presented at the Symp. on For. Fert., April 1967, Gainesville, Fla.], Tenn. Val. Auth. Natl. Fert. Dev. Cent., Muscle Shoals, Ala., p. 15–19.

Bilan, M.V.
 1971. Some aspects of tree root distribution. *In* Mycorrhizae, E.
 Hacskaylo, ed., 1st North Am. Conf. on Mycorrhizae, Univ. Ill.,
 Urbana, April 1969. USDA For. Serv. Misc. Publ. 1189, p. 69–80.

126. Bilan, M.V., and S.W. Jan. 1968. Needle moisture content as indicator of cessation of root elongation in loblolly pine seedlings. (Abstr.) Bull. Ecol. Soc. Am. 49(3):109. FA30:2007.

Bilan, M.V., and J.J. Stransky.
1966. Pine seedling survival and growth response to soils of the Texas post-oak belt. Stephen F. Austin State Coll., Nacogdoches, Tex., Dep. For. Bull. 12.

Biswell, H.H.
 1935. Effects of environment upon the root habits of certain deciduous forest trees. Bot. Gaz. 96(4):676–708. BA10:2694.

129. Bjallovich, Ju.P.
1968. [Taxodium distichum in Samarkand, and the mystery of its pneumatophores.] Lesoved. Mosk. 1:87–91. FA30:2009. [Engl. summ.]

 Blakeslee, G.M.
 1975. Basidiospore germination and function in ectomycorrhizal synthesis by certain members of the Agaricales. Diss. Abstr. Int. 36B(6):2550–2551. FA37:6015.

Blaser, R.E.
 1976. Plants and de-icing salts. Am. Nurseryman 144(12):8–9, 48, 50, 52–53.

132. Bloomberg, W.J. 1963. The significance of initial adventitious roots in poplar cuttings and the effect of certain factors on their development. For. Chron. 39(3):279–289.

Boerker, R.H.1916. Ecological investigations upon the germination and early growth of forest trees. Nebr. Univ., Univ. Stud. 16(1-2):1-89.

Boisen, A.T., and J.A. Newlin.1910. The commercial hickories. USDA For. Serv. Bull. 80.

135. Bond, G.
1954. The formation and function of some non-legume root nodules.
In 8th Int. Bot. Congr., [Paris 1954], Part 10, Section 21, p. 14.
FA16:230.

Bond, G.
 1956. Evidence for fixtation of nitrogen by root nodules of alder (Alnus) under field conditions. New Phytol. 55(2):147-153.
 FA18:3780.

Bond, G.
1958. Symbiotic nitrogen fixation by non-legumes. *In* Nutrition of legumes. E.G. Hallsworth, ed., Univ. Nottingham 5th Easter Schl. Agric. Sci., Butterworths Sci. Publ., London 1958, p. 216.

Bond, G.1967. Fixation of nitrogen by higher plants other than legumes.Annu. Rev. Plant Physiol. 18:107–126.

139. Bond, G. 1976. The results of the IBP survey of root-nodule formation in non-leguminous angiosperms. *In* Symbiotic nitrogen fixation in plants. P.S. Nutman, ed., Int. Biol. Programme 7, Cambridge Univ. Press, p. 443–474.

Bond, G., and E.J. Hewitt.1962. Cobalt and the fixation of nitrogen by root nodules of *Alnus* and *Casuarina*. Nature (Lond.) 195(4836):94–95. FA24:257.

141. Bond, G., and E.J. Hewitt. 1967. The significance of copper for nitrogen fixation in nodulated Alnus and Casuarina plants. Plant and Soil 27(3):447-449. FA29:3555.

142. Bond, G. and J.T. MacConnell. 1955. Nitrogen fixation in detached non-legume root nodules. Nature (Lond.) 176(4482):606. FA17:223.

Bond, G., and G.D. Scott.
 1955. An examination of some symbiotic systems for fixation of nitrogen. Ann. Bot. (N.S.) 19(73):67–77.

144. Bond, G., W.W. Fletcher, and T.P. Ferguson.1954. The development and function of the root nodules of *Alnus*, *Myrica*, and *Hippophae*. Plant and Soil 5(4):309–323. FA16:2762.

Bond, J.J. 1972. Mycorrhizal fungi of oak in North Carolina and their influence on uptake of potassium by oak seedlings. Diss. Abstr. Int. 32B(12):6770. FA35:6726.

146. Bormann, F.H. 1954. Ecological implications of size and weight changes in *Pinus taeda* L. seedlings during the first growing season. (Abstr.) Bull. Ecol. Soc. Am. 35(3):59–60. FA16:1523. 147. Bormann, F.H.

1961a. Intraspecific root grafts and transport. *In* Recent advances in botany, 9th Int. Bot. Congr., [Montreal 1959], 2:40–41. BA37:3466.

148. Bormann, F.H.

196lb. Intraspecific root grafting and the survival of eastern white pine stumps. For. Sci. 7(3):247–256. BA37:7344.

149. Bormann, F.H.

1966. The structure, function, and ecological significance of root grafts in *Pinus strobus* L. Ecol. Monogr. 36(1):1–26. BA47:50490.

Bormann, F.H., and B.F. Graham, Jr.
 1959. The occurrence of natural root graftin

1959. The occurrence of natural root grafting in eastern white pine, *Pinus strobus* L., and its ecological implications. Ecology 40(4):677–691. BA35:9450.

151. Bormann, F.H., and B.F. Graham, Jr.

1960. Translocation of silvicides through root grafts. J. For. 58(5):402–403. BA35:48995.

152. Bormann, F.H., and B.F. Graham, Jr.

1961. A study of intraspecific root grafts in white pine, *Pinus strobus* L., by means of radioisotopes and dyes. *In* Recent advances in botany, 9th Int. Bot. Congr., [Montreal 1959], 2:1375–1377.

153. Boullard, B.

1959. A propos des mycorrhizes du *Pinus strobus* L. [Concerning the mycorrhizae of *Pinus strobus* L.] Bull. Trimest. Soc. Mycol. Fr. 75(2):194–200. FA21:2853.

154. Boullard, B., and T. Dominik.

1958. Badania porownawcze nad mikoryzami *Pinus strobus* L. z roznych stanowick we Francji i w Polsce. [Investigations on mycorrhizae of *P. strobus* L. at different sites in France and Poland.] Pr. Inst. badarv. Lesn. 178, p. 45–84. FA20:4267. [Fr., Russ., and Ger. summ.]

155. Boullard, B., and H.A. Ferchau.

1962. Endotrophic mycorrhizae of plants collected in some Eastern American and Canadian white pine communities. Phyton 19(1):65–71. FA24:4747.

156. Bourdeau, P.

1954. Oak seedling ecology determining segregation of species in Piedmont oak-hickory forests. Ecol. Monogr. 24(3):297–320.

157. Box, B.H.

1968. A study of root extension and biomass in a six year old pine plantation in Southeast Louisiana. Diss. Abstr. 28B(9):3545–3546. FA30:342.

158. Boyce, J.S., Jr.

1959. Root rot in pine plantations. For. Farmer 19(3):8,17–18.

159. Boyce, J.S., Jr. 1960. Distrib

1960. Distribution of *Ceratocystis fagacearum* in roots of wilt-infected oaks in North Carolina. Phytopathology 50 (10):775–776. BA36:15509.

160. Boyce, J.S., Jr.

1962a. *Fomes annosus* in white pine in North Carolina. J. For. 60(8):553–557. FA24:2287.

161. Boyce, J.S., Jr.

1962b. Greenhouse inoculations of coniferous seedlings with *Fomes annosus*. (Abstr.) Phytopathology 52(1):4. FA23:5420.

162. Bovce, J.S., Jr.

1966. Sporulation by *Peniophora gigantea* with reference to control of Annosus root rot. [Fomes annosus] For. Sci. 12(1):2-7. FA27:6204.

163. Boyer, W.D., R.M. Romancier, and C.W. Ralston.

1971. Root respiration rates of four tree species grown in the field. For. Sci. 17(4):492–493. BA54:16554.

164. Braun, E.L.

1936. Notes on root behavior of certain trees and shrubs of the Illinoian till plain of Southwestern Ohio. Ohio J. Sci. 36(3):141–146. BA11:5412.

165. Braun, H.J.

1969. Root germs in young and older stems of *Populus*. Transl. Dep. Fish. For. Can. OOFF-69. Transl. *from* Z. Bot., Stuttgart, 1963, 51(5):441-451. FA26:1744.

166. Braun, H.J., and J. Lulev.

1969. [Infection of intact, finger-thick spruce roots by *Fomes annosus*. I. The structure of the surface periderm, and first infection results.] Forstwiss. Centralbl. 88(6):327–338. FA32:2718. [In Ger. with Engl. summ.]

167. Braun, H.J., and J. Lulev.

1970. [Infection of intact, finger-thick spruce roots by *Fomes annosus*. II. Infection paths and the spread of the fungus immediately after infection.] Forstwiss. Centralbl. 89(5):269–275. FA32:2719. [In. Ger. with Engl. summ.]

168. Brender, E.V.

1955. Drought damage to pines. For. Farmer 14(10):7,15. FA17:513.

169. Bretz, T.W.

1951. Oak wilt. J. For. 49(3):169-171.

170. Brewer, C.W., and N.E. Linnartz.

1976. Loblolly pine root distribution influenced by physical properties of soil. Agron. Abstr. 1976:182.

171. Briggs, A.H.

1939. Report of planting experiment to determine the effect of root exposure on deciduous planting stock. J. For. 37(12):939–943. BA15:13410.

172. Brix, H.

1960. Determination of viability of loblolly pine seedlings after wilting. Bot Gaz. 121(4):220–223. FA23:4874.

173. Broadfoot, W.M.

1973. Water table depth and growth of young cottonwood. USDA For. Serv. Res. Note SO–167. FA35:7447.

174. Broadfoot, W.M., and H.L. Williston.

1973. Flooding effects on southern forests. J. For. 71(9):584–587.

175. Brooks, M.G.

1951. Effect of black walnut trees and their products on other vegetation. W. Va. Agric. Exp. Stn. Bull. 347.

176. Brown, A.B.

1935. Cambial activity, root habit and sucker shoot development in two species of poplar. New Phytol. 34(3):163–179. BA10:16893.

177. Brown, A.H.F., A. Carlisle, and E.J. White.

1966. Some aspects of the nutrition of Scots pine on peat. *In* Physiology in forestry. Rep. Soc. Forest. G. B. 6th Discuss. Meet., [Edinburgh, Scotland 7–9 January 1966], p. 78–87.

178. Brown, C.L., and P.P Kormanik.

1967. Suppressed buds on lateral roots of *Liquidambar styraciflua*. Bot. Gaz. 128(3/4):208–211. BA49:52908.

179. Brown, James Harold.

1968. Patterns of variation in root systems of Scotch pine provenances. Diss. Abstr. 28B(9):3546–3547.

180. Brown, J.H.

1969. Variations in roots of greenhouse grown seedlings of different Scotch pine provenances. Silvae Genet. 18(4):111-117. BA51:57739.

181. Brown, J.H., and F.C. Cech.

1972. Top and root characteristics of greenhouse grown seedlings of different black cherry provenances. Silvae Genet. 21(3/4):130–133. BA56:8145.

182. Brown, James Henry, Jr.

1965. A study of root distribution in two hardwood stands in the Duke Forest. Ph.D. thesis, Duke Univ., Durham, N.C. (Diss. Abstr. 26(2):593.)

183. Brown, J.H., Jr., and T.G. Bourn.

1973. Patterns of soil moisture depletion in a mixed oak stand. For. Sci. 19(1):23–30. FA35:80.

184. Brown, J.H., Jr., and F.W. Woods.

1968. Root extension of trees in surface soils of the North Carolina Piedmont. Bot. Gaz. 129(2):126–132. BA50:50059.

185. Brown, W.G.E., and D.S. Lacate.

1961. Rooting habits of white and red pine. Can. For. Res. Branch Tech. Note 108. FA23:1708.

186. Bryan, W.C., and P.P. Kormanik.

1977. Mycorrhizae benefit survival and growth of sweetgum seedlings in the nursery. South. J. Appl. For. 1(1):21–23.

Bryan, W.C., and J.L. Ruehle.
 1976. Growth stimulation of sweetgum seedlings induced by the endomycorrhizal fungus *Glomus mosseae*. Tree Plant. Notes 27 (2):9,14.

188. Bryan, W.C., and B. Zak. 1961. Synthetic culture of mycorrhizae of southern pines. For. Sci. 7(2):123–129. FA23:1703.

Bryan, W.C., and B. Zak.
 1962. Additional synthesis of mycorrhizae on shortleaf and loblolly pines. For. Sci. 8(4):384. FA24:3365.

Buckman, R.E., and L.H., Blankenship.
 1965. Repeated spring prescribed burning reduces abundance and vigor of aspen root suckering. J. For. 63(1):23–25. BA46:40688.

Buell, M.F., and H.F. Buell.
 1959. Aspen invasion of prairie. Bull. Torrey Bot. Club 86(4):264–265

Bugbee, W.M.
 1962. Host range and bioassay of field soil for *Cylindrocladium scoparium*. (Abstr.) Phytopathology 52(8):726. BA41:3666.

193. Buijtenen, J.P. van. 1966. Testing loblolly pines for drought resistance. Tex. For. Serv. Tech. Rep. 13.

194. Buijtenen, J.P. van, M.V. Bilan, and R.H. Zimmerman. 1976. Morpho-physiological characteristics related to drought resistance in *Pinus taeda*. *In* Tree physiology and yield improvement. M.G.R. Cannell, and F.T. Last, eds., Academic Press, N.Y., p. 349–359.

Bunger, M.T., and H.J. Thomson.
 1938. Root development as a factor in the success or failure of windbreak trees in the southern high plains. J. For. 36(8):790–803. BA14:9291.

Burger, H.
 1930. Bodenveränderung und wurzelbildung. Schweiz. Z. Forstwes. 2:67–71.

Burns, G.P.
 1914. Studies in tolerance of New England forest trees. I. Development of white pine seedlings in nursery beds. Vt. Agric. Exp. Stn. Bull. 178, p. 125–144.

198. Burns, G.P.
1937. Studies in tolerance of New England forest trees. XIII. The effect of root development on height and diameter growth. Vt. Agric. Exp. Stn. Bull. 422.

199. Burton, J.D. 1971. Prolonged flooding inhibits growth of loblolly pine seedlings. USDA For. Serv. Res. Note SO-124.

200. Busgen, M., and E. Munch.1929. The structure and life of forest trees. 3rd ed., Transl. from Ger. by T. Thomson. John Wiley and Sons, Inc., N.Y.

Bushey, D.J.
 1937. Root extension of shade trees. Natl. Shade Tree Conf. Proc.
 13:22-30.

02. Bushey, D.J. 1946. Regeneration of the root systems of pin oak (*Quercus palustris*) and American elm (*Ulmus americana*) following transplanting in the fall and spring. Natl. Shade Tree Conf. Proc. 22:46–56. BA21:20594.

203. Cabrera, H., and F.W. Woods. 1975. Effects of root deformation upon early growth of loblolly pine (*Pinus taeda* L.) Tenn. Farm and Home Sci. Progr. Rep. 93, p. 28–30. FA37:276. 204. Calderone, R.A., and R.P. True.
1967. Pathogenicity and physiology of *Pythium vexans*. (Abstr.)
Phytopathology 57(8):806. FA30:2491.

205. Campana, R.J.

1953. Corticum galactinum and white pine needle blight. (Abstr.) Phytopathology 43(9): 468. FA15:1620.

206. Campana, R.J.
1954. Corticum galactinum does not cause white pine needle blight.
Plant Dis. Rep. 38(4):297–303. FA15:1620.

Campbell, W.A.
 1967. Phytophthora spp.: Phytophthora spp., particularly P. cactorum (Lebert and Cohn) Schroet., P. cinnamomi Rands, and P. lateralis Tucker and Milbrath. In Important forest insects and diseases of mutual concern to Canada, the United States, and Mexico. A.G. Davidson, and R.M. Prentice, eds., Can. For. Serv. Publ. 1180, p. 57-60.

208. Campbell, W.A., and J.H. Miller. 1952. Windthrow of root-rotted oak shade trees. Plant Dis. Rep. 36(12):490. BA28:6934.

209. Canada Department of Agriculture.
 1953. Deterioration of birch. Can. Dep. Agric. Entomol. Br., Annu.
 Rep. For. Insect and Dis. Surv. 1953, p. 81. FA16:739.

Cannon, W.A.
 1949. A tentative classification of root systems. Ecology 30(4):542–548.

211. Cannon, W.A.
1954. A note on the grouping of lateral roots. Ecology 35(2):
293-295. FA15:3350.

Capps, S.R.
 1931. Glaciation in Alaska. U.S. Geol. Surv. Prof. Pap. 170–A.

Carleton, L.C.
 1963. Salt tolerances of *Pinus thunbergii*, *P. radiata*, *P. resinosa* and *P. taeda*. Diss. Abstr. 23(9):3090–3091. FA25:272.

214. Carlson, L.W.
1976. Root initiation of lodgepole pine and white spruce seedlings grown under varying light conditions. Bi-mon. Res. Notes 32(4):21-22.

215. Carlson, L.W.1977. The effect of defoliation on conifer seedling root initiation.Bi-mon. Res. Notes 33(1):1.

216. Carlson, L.W., and F. Endean. 1976. The effect of rooting volume and container configuration on the early growth of white spruce (*Picea glauca*) seedlings. Can. J. For. Res. 6(2): 221–224.

Carlson, L.W., and L.D. Nairn.
 1977. Root deformities in some container-grown jack pine in south-eastern Manitoba. For. Chron. 53(3):147–149.

Carlson, M.C.1950. Nodal adventitious roots in willow stems of different ages.Am. J. Bot. 37(7):555-561. BA25:8855.

219. Carlson, W.C.
1974. Root initation induced by root pruning in northern red oak. *In*Forestry research review—1974, Ohio Agric. Res. and Devel.
Cent., Wooster, p. 14–16.

220. Carpenter, I.W., and A.T. Guard.
1954. Anatomy and morphology of the seedling roots of four species of the genus *Quercus*. J. For. 52(4):269–274. FA15:3252.

221. Carrodus, B.B.
1967. Absorption of nitrogen by mycorrhizal roots of beech. II.
Ammonium and nitrate as sources of nitrogen. New Phytol. 66(1):1–
4. BA48:51595.

Carter, J.C.
 1941. Preliminary investigation of oak diseases in Illinois. Ill. Nat.
 Hist. Surv. Bull. 21:195–230.

 Chadwick, L.C.
 1936. Ohio shade tree fertilization experiments. Natl. Shade Tree Conf. Proc. 12:55-66. 224. Chadwick, L.C.

1939. The distribution of roots of Moline elms in relationship to fertilizer application. Natl. Shade Tree Conf. Proc. 15:38–51.

Chadwick, L.C., D. Bushey, and G. Pletcher.
 1937. Root distribution studies. Proc. Am. Soc. Hort. Sci. 35:734
 738. BA12:8741.

226. Chandler, R.F., Jr.

1941. The amount and mineral nutrient content of freshly fallen leaf litter in the hardwood forests of central New York. J. Am. Soc. Agron. 33(10): 859–871. FA3:284.

227. Chapman, A.G.

1935. The effects of black locust on associated species with special reference to forest trees. Ecol. Monogr. 5:37–60.

228. Cheyney, E.G.

1928. The root system of the hazel. J. For. 26(8):1046-1047. BA4:22320.

229. Cheyney, E.G.

1929. A study of the roots in a square yard of jack pine forest. J. For. 27(5):546–549. BA5:2041.

230. Cheyney, E.G.

1932. The roots of a jack pine tree. J. For. 30(8):929-932.

231. Childs, T.W.

1937. Variability of *Polyporus schweinitzii* in culture. Phytopathology 27(1):29–50.

232. Chouinard, L.

1959. Sur l'existence d'un centre quiescent au niveau de l'apex radiculaire juvénile de *Pinus banksiana* Lamb. [On the existence of a quiescent center in the root apical meristem of *P. banksiana*.] Laval Univ. For. Res. Found. Contr. 4, p. 27–31.

233. Christensen, C.M.

1938. Root rot of pines caused by *Armillaria mellea*. (Abstr.) Phytopathology 28(1):5.

234. Chung, H.H., and P.J. Kramer.

1975. Absorption of water and ³²P through suberized and unsuberized roots of loblolly pine. Can. J. For. Res. 5(2): 229–235. BA60:48892.

235. Clark, F.B.

1969. Factors affecting the production of fibrous roots on black walnut seedlings. Diss. Abstr. 29B(7):2252. FA30:5595.

236. Clark, F.B., and C.K. Losche.

1969. Importance of the A horizon in hardwood seedling establishment. J. For. 67(7):504–505. BA51:11139.

237. Clark, J.

1961. Birch dieback. *In* recent advances in botany, 9th Int. Bot. Congr., [Montreal 1959], 2:1551–1555.

Clements, F.E., J.E. Weaver, and H.C. Hanson.
 1929. Plant competition: An analysis of community functions. Carnegie Inst. Wash. Publ. 398.

239. Cleveland, T.E., H.E. Garrett, and H.G. Hedrick.
1976a. Characterization of associative soil fungi with mycorrhizae
on loblolly pine and white oak seedlings. (Abstr.) Proc. La. Acad.
Sci. 39:109.

240. Cleveland, T.E., H.E. Garrett, and H.G. Hedrick. 1976b. Effects of SO₂ and O₃ on colony growth of *Pisolithus tinctorius*—a mycorrhizal isolate on loblolly pine and white oak

seedlings. (Abstr.) Proc. La. Acad. Sci. 39:108.

241. Cleveland, T.E., H.E. Garrett, and H.G. Hedrick.

1976c. Effects of SO₂ and O₃ on colony growth of selected associative soil fungi with root surface mycorrhizae on loblolly pine and white oak seedlings. (Abstr.) Proc. La. Acad. Sci. 39:108–109.

242. Cleveland, T.E., H.E. Garrett, and H.G. Hedrick.
1976d. Effects of SO₂ and O₃ on sporulation of selected associative soil fungi with mycorrhizae on loblolly pine and white oak seedlings.
(Abstr.) Proc. La. Acad. Sci. 39:109.

243. Cobb, F.W., Jr., and W.W. Wilcox.
1967. Comparison of susceptibility of *Abies concolor* and *Pinus ponderosa* wood to decay by *Fomes annosus*. Phytopathology

57(12):1312–1314. FA29:4106.

244. Coile, T.S.

1937. Distribution of forest tree roots in North Carolina Piedmont Soils. J. For. 35(3):247–257.

245. Coile, T.S.

1940. Soil changes associated with loblolly pine succession on abandoned agricultural land of the Piedmont Plateau. Duke Univ. Sch. For. Bull. 5.

246. Coleman, E.

1945. Strange root formation in weeping willow (*Salix babylonica*). Victorian Nat. 62(7):121. BA20:11082.

Commonwealth Bureau of Horticultural and Plantation Crops.
 1965. Bibliography on the ion-exchange capacity of roots. Commomr. Bur. Hortic. Plant Crops, East Malling, Eng. FA27:137.

248. Connola, D.P., and E.C. Wixson. 1963a. Effects of soil and other environmental conditions on white pine weevil attack in New York. J. For. 61(6): 447–448. FA25:1012.

249. Connola, D.P., and E.C. Wixson.

1963b. White pine weevil attack in relation to soils and other environmental factors in New York. Bull. N.Y. State Museum and Sci. Serv. 389, Albany.

250. Conway, V.M.

1940. Aeration and plant growth in wet soils. Bot. Rev. 6(4):149–163.

251. Cook, D.B.

1961. Shoestring fungus and planted larch on cutover land. J. For. 59(11):824–826. FA23:3828.

252. Cook, D.B., and D.S. Welch.

1957. Backflash damage to residual stands incident to chemipeeling. J. For. 55(4):265–267. FA18:4109.

253. Cooper, W.S.

1911. Reproduction by layering among conifers. Bot. Gaz. 52(5):369–379.

254. Copeland, O.L, Jr.

1952. Root mortality of shortleaf and loblolly pine in relation to soils and littleleaf disease. J. For. 50(1): 21–25. FA13:3191.

Copeland, O.L., Jr., and R.G. McAlpine.
 1962. Soil characteristics associated with spot die-out in loblolly

pine plantations. For. Sci. 8(1):12–15. FA23:5631.

256. Cordell, C.E.

1973. Cylindrocladium root rot of black walnut and yellow-poplar nursery seedlings. In Southeast. U.S. Abstr. Pap., 2nd Int. Congr. Plant Pathol. 0689, Minneapolis. FA36:3407.

257. Cordell, C.E., and D.H. Marx.

1977. Mycorrhizae + tree seedlings = increased southern timber production. South. Lumberman, June 1, 1977.

258. Cordell, C.E., and S.J. Rowan.

1975. Cylindrocladium scoparium infection in a natural sweetgum stand. Plant Dis. Rep. 59(10):775–776. FA37:2988.

259. Cordell, C.E., W.J. Stambaugh, C.E. Affeltranger, and J.L. Knighten.

1970. Rhododendron and mountain laurel, new hosts of *Fomes annosus* in western North Carolina. Plant Dis. Rep. 54(7):560. FA32:2726.

260. Corovic, M., and L. Stjepanovic.

1958. Rastenje korena i korenskih dlakakod biljaka *Pinus nigra* L. i *Robinia pseudoacacia* L. na razlicitim temperaturama pod loge. [The growth of roots and root-hairs of *P. nigra* and *R. pseudoacacia* at different substratum temperatures.] Sumarstve 11(5/6):305–314. FA20:1551.

261. Corte, A.

1969. Richerche sull 'influenza dell' infezione micorrizica sullo sviluppo vegetativo, sulla vigoria e sullo stato sanitario di tre specie di pino. I. Risultati preliminari sul comportamento delle piantine in vivaio e nel primo anno di coltura in campo. [The influence of infection with mycorrhizae on the growth, vigour and health of three pine species. I. Preliminary results, on the behavior of plants in the nursery and during the first year in the field.] Arch. Bot. e Biogeogr. Ital. Modena 45(1/2):1–28. [Engl. summ.] FA31:4376.

262. Coultas, C.L.

1973. Response of pine seedlings to manipulations of the spodic horizon. Proc. Soil and Crop Sci. Soc. Fla. 32:121–125. FA38:3251.

263. Coutts, M.P., and W. Armstrong.

1976. Role of oxygen transport in the tolerance of trees to waterlogging. *In* Tree physiology and yield improvement. M.G.R. Cannell, and F.T. Last, eds., Academic Press, N.Y. p. 361–385.

264. Cox, R.S.

1951. A preliminary report on etiological and control studies of damping-off and root-rot in the conifer seedbed in Delaware. Plant Dis. Rep. 35(8):374–378. FA13:2070.

265. Cox, R.S.

1953. Etiology and control of a serious complex of diseases of conifer seedlings. (Abstr.) Phytopathology 43(9):469. FA15:1292.

266. Cox, R.S.

1954. Cylindrocladium scoparium on conifer seedlings. Univ. Del. Agric. Exp. Stn. (Tech.) Bull. 301.

267. Cox, T.L.

1972. Relationships of metabolic activity and mineral content to surface phenomena of four diameter classes of *Liriodendron* roots. Agron. Abstr. 1972:138.

268. Cox, T.L.

1973. Production, mortality and nutrient cycling in root systems of *Liriodendron* seedlings. Diss. Abstr. Int. 33B(11):5245–5246. FA36:2483.

269. Cox, T.L.

1975. Seasonal respiration rates of yellow-poplar roots by diameter classes. For. Sci. 21(2):185–188. BA60:48914.

270. Coyne, J.F.

1968. Simulated Nantucket pine tip moth attacks reduce root development of shortleaf pine seedlings. J. Econ. Entomol. 61(1):319–320. BA49:69827.

271. Crandall, B.S.

1938. A root and collar disease of pine seedlings caused by *Sphaeropsis ellisii*. Phytopathology 28(3):227–229.

272. Crandall, B.S., G.F. Gravatt, and M.M. Ryan.

1945. Root disease of *Castanea* species and some coniferous and broadleaf nursery stocks, caused by *Phytophthora cinnamomi*. Phytopathology 35(3):162–180. FA7:99.

273. Crawford, R.M.M., and M.A. Baines.

1977. Tolerance of anoxia and the metabolism of ethanol in tree roots. New Phytol. 79:519–526.

274. Crossley, D.I.

1940. The effect of a compact subsoil horizon on root penetration. J. For. 38(10):794–796. FA2:293.

275. Cummings, W.H.

1942. Exposure of roots of shortleaf pine stock. J. For. 40(6):490–492

276. Curtis, J.D.

1944. Northern white-cedar on upland soils in Maine. J. For. 42(10):756-759.

277. Cuthbert, R.A., W.N. Cannon, Jr., and J.W. Peacock. 1975. Relative importance of root grafts and bark beetles to the spread of Dutch elm disease. USDA For. Serv. Res. Note NE–206. FA37:3003.

278. Daft, M.J., and E, Hacskaylo.

1977. Growth of endomycorrhizal and nonmycorrhizal red maple seedlings in sand and anthracite spoil. For. Sci. 23(2):207–216.

 Daft, M.J., E. Hacskaylo, and T.H. Nicolson.
 1975. Arbuscular mycorrhizae in plants colonizing coal spoils in Scotland and Pennsylvania. *In* Endomycorrhizas. F.E. Sander, B. Mosse, and P.B. Tinker, eds., Academic Press, London, p. 561–580. FA37:6013.

280. Dale, J., A.L. McComb, and W.E. Loomis.

1955. Chlorosis, mycorrhizae and the growth of pines on a high-lime soil. For. Sci. 1(2):148–157. FA17:218.

281. Daly, G.T.

1966. Nitrogen fixation by nodulated *Alnus rugosa* NS. Can. J. Bot. 44(12):1607–1621.

282. Dandeno, J.B.

1909. Mutual interaction of plant roots. Mich. Acad. Sci. Annu. Rep. 11:24–25.

283. Danielson, R.M., and C.B. Davey. 1969. Microbial recolonization of a fumigated nursery soil. For. Sci. 15(4):368–380. FA31:4367.

284. Davey, C.B., and W.A. Bland.

1970. Nutrient accumulation patterns of loblolly pine (*Pinus taeda* L.) seedlings in nursery seedbeds. Agron. Abstr. 1970:158.

285. Davidson, D.W.

1963. Living stumps of *Tsuga canadensis* (L.) Carr. (hemlock) in northern New Jersey. Bull. Torrey Bot. Club 90(3):204–207. FA25:291.

286. Davidson, D.W., and R.M. Davis

1964. Further observations on living stumps of *Tsuga canadensis* (L.) Carr. (hemlock) in northern New Jersey. Bull. Torrey Bot. Club 91(3):233–234. FA26:1903.

287. Davidson, H.

1970. The effect of root system and depth of planting on the survival and growth of Scots pine when planted in a highway right-of-way. Tree Plant. Notes 21(2):26–27.

288. Davidson, R.W., and R.F. Patton.

1961. *Paxillus atrotomentosus* causes brown root rot in dead jack pine in plantations in Wisconsin. Plant Dis. Rep. 45(11): 836–838. FA24:2294.

289. Davies, G.

1973. Response of loblolly pine (*Pinus taeda* L.) seedlings from two seed sources to favorable and unfavorable moisture regimes. M.S. Thesis, Stephen F. Austin State Univ., Nacogdoches, Tex.

290. Davis, D.E.

1959. Some effects of calcium deficiency on the anatomy of *Pinus taeda*. Am. J. Bot. 36(3):276–282.

291. Davis, E.F.

1928. The toxic principle of *Juglans nigra* as identified with synthetic juglone, and its toxic effects on tomato and alfalfa plants. (Abstr.) Am. J. Bot. 15(10, suppl.):620.

292. Davis, J.B., and R.L. Barnes.

1973. Effects of soil-applied fluoride and lead on growth of loblolly pine and red maple. Environ. Pollut. 5(1):35–44. FA36:4805.

293. Davis, S.H.

1942. Sclerotium bataticola a cause of damping-off in seedling conifers. Science 95:70. FA4:55.

294. Dawson, J.O.

1978. Nitrogen fixation, photosynthesis and early growth of *Alnus glutinosa*. Ph.D. Thesis, Ia. State Univ., Ames.

295. Day, M.W.

1941. The root system of red pine saplings. J. For. 39(5):468–472. BA15:24271.

296. Day, M.W.

1944. The root system of aspen. Am. Midl. Nat. 32(2):502–509. BA19:2203.

297. Day, M.W.

1945. A comparison of the root systems of jack pine and tamarack. J. For. 43(1):41–42. BA19:17332.

298. Day, M.W., and F.H. Vogel.

1944. Silviculture and utilization of balsam poplar. J. For. 42(7):512–514.

299. Day, R.J., and G.R. MacGillivray.

1975. Root regeneration of fall-lifted white spruce nursery stock in relation to soil moisture content. For. Chron. 51(5):196–199. FA37:4395.

300. Day, R.J., J.P. Stupendick, and J.M. Butler.

1976. Root periodicity and root regeneration potential are keys to successful plantation establishment. *In* Ont. Minist. of Nat. Resour.—Great Lakes For. Res. Cent., Plant. Establ. Symp. FA39:396.

301. Day, W.R.

1950. The drought-crack of conifers. *In* 25th Imperial For. Inst. Rep., Oxford, 1948–9, p. 12–13. FA11:3245.

302. Dean, G.W.

1927. Rate of autumnal root growth in white pine. Unpubl. ms., Yale Univ. Sch. For.

303. DeByle, N.V.

1962. Tracing and evaluating functional root connections among stems within aspen clones. Diss. Abstr. 23(1):9–10. FA24:1871.

304. DeByle, N.V.

1964. Detection of functional intraclonal aspen root connections by tracers and excavation. For Sci. 10(4):386–396. BA46:26946.

305. DeByle, N.V., and I.C.M. Place.

1959. Rooting habits of oak and pine on Plainfield sand in Central Wisconsin. Univ. Wis. For. Res. Notes 44.

306. DeGruchy, J.H.B.

1956. Water fluctuations as a factor in the life of six of the higher plants of Central Oklahoma. Proc. Okla. Acad. Sci. 37:45–46.

307. DeMent, J.A., and E.L. Stone.

1968. Influence of soil and site on red pine plantations in New York. II. Soil type and physical properties. Cornell Univ. Agric. Exp. Stn. Bull. 1020, Ithaca, N.Y.

308. Denne, M.P.

1972. A comparison of root- and shoot-wood development in conifer seedlings. Ann. Bot. (N.S.) 36(146):579–587. BA54:68795.

309. DenUyl, D.

1961. Some observations on bald cypress in Indiana. Ecology 42(4):841-843. FA23:3306.

310. Denyer, W.B.G.

1960. Cultural studies of *Flammula alnicola* (Fr.) Kummer and *Flammula conissans* (Fr.) Gillet. Can. J, Bot. 38(6):909–920.

311. Denyer, W.B.G., and C.G. Riley.

1964. Dieback and mortality of tamarack caused by high water. For. Chron. 40(3):334–338. FA26:2316.

312. Detwiler, S.B.

1916. The bald cypress (*Taxodium distichum*): Identification and characteristics. Am. For. 22(274):577–581.

313. Dhillon, P.S., W.R. Byrnes, and C. Merritt.

1967. Simazine and phosphorus interactions in red pine seedlings. Weeds 15(4):339–343. FA30:760.

314. Dickson, R.E., T.C. Broyer, and C.M. Johnson.

1972. Nutrient uptake by tupelo gum and bald cypress from saturated or unsaturated soil. Plant and Soil 37(2):297–308. BA55:30917.

315. Dickson, R.E., J.F. Hosner, and N.W. Hosley.

1965. The effects of four water regimes upon the growth of four bottomland tree species. For. Sci. 11(3):299–305.

316. Diebold, C.H.

1933. Root distribution and penetration of soil layers. J. For. 31(4):481-482.

317. Dimbleby, G.W.

1953. Natural regeneration of pine and birch on the heather moors of North-east Yorkshire. Forestry 26:41–52.

318. Dirr, M.A.

1974. Tolerance of honeylocust seedlings to soil-applied salt. Hort-Science 9(1):53–54. FA36:4023.

319. Dirr, M.A.

1976. Selection of trees for tolerance to salt injury. J. Arboric. 2(11):209-216.

320. DiSanzo, C.P., and R.A. Rohde.

1969. Xiphinema americanum associated with maple decline in Massachusetts. Phytopathology 59(3):279–284.

321. Dittmer, H.J.

1949. Root hair variations in plant species. Am. J. Bot. 36(2):152–154.

322. Doak, K.D.

1934a. Cortical parasitism of conifer-seedling roots in pure culture by mycorrhizal and non-mycorrhizal fungi. (Abstr.) Phytopathology 24:6–7.

323. Doak, K.D.

1934b. Fungi that produce ectotrophic mycorrhizae of conifers. (Abstr.) Phytopathology 24:7.

324. Doak, K.D.

1934c. Mycorrhizae and their relation to shade trees. Natl. Shade Tree Conf. Proc. 10:99–105. BA11:203.

325. Doak, K.D.

1955a. Pine root reaction in sterile culture to mycorrhizal and other fungi. Am. Midl. Nat. 54(2):443–451. FA18:2596.

326. Doak, K.D.

1955b. Mineral nutrition and mycorrhizal association of bur oak. Lloydia 18(3):101–108.

327. Dochinger, L.S.

1967. *Leptographium* root decline of eastern white pine. (Abstr.) Phytopathology 57(8):809. FA30:2544.

328. Dosser, R.C., and R.R. Hicks, Jr.

1975. Ortet and season of collection significantly affect rooting of river birch stem cuttings. Tree Plant. Notes 26(2):11–12.

329. Drake, C.R., J.E. Kuntz, and A.J. Riker.

1957. Chemical control of the oak wilt. Univ. Wis. For. Res. Note 35. FA19:3170.

330. Dreisinger, B.R., D.V. Baxter, and S.H. Spurr.

1956. Decline of red pine on heavy loam soils. Mich. For. 14. FA18:702.

331. Drobnov, P.P., and L.S. Savel'eva.

1973. [Morphological changes in the stem and roots of *Robinia pseudoacacia* in shelterbelts choked with an accumulation of dust.] Byul. VNII agrolesomelior 12(66):37–39. [In Russ. with Engl. summ.] FA36:282.

332. Dullaart, J.

1970. The auxin content of root nodules and roots of *Alnus glutinosa* (L.) Vill. J. Exp. Bot. 21(69):975–984. FA32:3942.

333. Dumbroff, E.B., and D.C.W. Brown.

1976. Cytokinin and inhibitor activity in roots and stems of sugar maple seedlings through the dormant season. Can. J. Bot. 54(3/4):191–197. BA62:4781.

334. Dumbroff, E.B., and D.P. Webb.

1978. Physiological characteristics of sugar maple and implications for successful planting. For. Chron. 54(2): 92–95.

335. Duncan, W.H.

1935. Root systems of woody plants of old fields of Indiana. Ecology 16(4):554–567. BA10:20371.

336. Duncan, W.H.

1941. A study of root development in three soil types in the Duke Forest. Ecol. Monogr. 11(2):141–164. BA15:21213.

337. Dunning, D.

1946. Roots of forest trees. A selected list of references. USDA For. Serv. Res. Note 52, Pac. Southwest For. and Range Exp. Stn. FA9:775.

338. Dykeman, W.R., and A.S. deSousa. 1966. Natural mechanisms of copper tolerance in a copper swamp forest. Can. J. Bot. 44:871–878.

339. Edmonds, A.S., J.M. Wilson, and J.L. Harley. 1976. Factors affecting potassium uptake and loss by beech mycorrhiza. New Phytol. 76(2):307–315. BA62:13866.

340. Egger, A.
1975. [Otiorrhynchus ovatus, a serious pest of nurseries and young plantations.] Allg. Forstztg. 85(1):7–11. [In Ger.] FA35:5375.

341. Eghbaltalab, M., G. Gay, and G. Bruchet.
1975. [Antagonism between 15 species of basidiomycetes and three pathogenic fungi of tree roots.] Bull. Mens. Soc. Linn. de Lyon.
44(7):203-208. [In Fr.] FA37:5915.

342 Einspahr, D.W.
1971. Growth and nutrient uptake of aspen hybrids using sand culture techniques. Silvae Genet. 20(4):132–137. BA54:25658.

343. Eis, S. 1970. Root-growth relationships of juvenile white spurce, alpine fir, and lodgepole pine on three soils in the interior of British Columbia. Can. For. Serv. Publ. 1276. FA32:5633.

344. Eis, S.
1977. Root forms in habitats with heavy shrub competition. Bi-mon.
Res. Notes 33(4):27-29.

345. Elder, W.C., and J.E Webster.
1959. Food reserves in post oak stumps and roots. Okla. Agric. Exp.
Stn. Tech. Bull. T-80. BA35:30202.

Eliason, E.J.
 1928. Comparative virulence of certain strains of *Pythium* in direct inoculation of conifers. Phytopathology 18:361–367.

1938. Buckwheat as a factor in the root rot of conifers. (Abstr.)
Phytopathology 28:7.

348. Eliasson, L.
1961. The influence of growth substances on the formation of shoots from aspen roots. Physiol. Plant. 14(1):150–156. BA36:45925.

349. Emerson, F.W.
1921. Subterranean organs of bog plants. Bot. Gaz. 72–359–374.
350. Enescu, V.

1960. Influenta provenientei (saminta, lastari, drajoni) si a virstei asupra sistemului de inradacinare al salcimului. [The influence of origin (seed, coppice shoots, root suckers) and age, on the root system of *Robinia pseudoacacia*.] Rev. Padurilor 75(7):396–399. FA22:4323.

Engler, A.
 1903. Untersuchungen über das Wurzelwachstum der Holzarten.
 Mitt. Schwiez. Centralanst. Forstl. Versuchswes. 7:247–272.

352. Enright, L.J.
1959. Growth response of rooted cuttings of pine and spruce. J. For.
57(7):509-510. BA34:8585.

353. Ewan, H.E., and R.J. Finnegan.
1967. Pine root-collar weevil *Hylobius radicis* Buch. *In* Important forest insects and diseases of mutual concern to Canada, the United States, and Mexico. A.G. Davidson, and R.M. Prentice, eds., Can. For. Serv. Publ. 1180, p. 111-114.

354. Fabricius, L. 1905. Untersuchungen über den Stärke-und Fettgehalt der Fichte auf der oberbayerischen Hochebene. Naturw. Zeit. Land-u. Forstw. 3:137-176.

355. Farmer, R.E., Jr.
1962a. Aspen root sucker formation and apical dominance. For. Sci. 8(4):403–410. BA42:11643.

Farmer, R.E., Jr.1962b. Depth and diameter of the parent roots of aspen root suckers.Mich. For. 23. FA24:1869.

357. Farmer, R.E., Jr.
1962c. Some physiological aspects of root sucker initiation and early growth in *Populus tremuloides* and *P. grandidentata*. Diss. Abstr. 23(1):10. FA24:1870.

358. Farmer, R.E., Jr.
1970. Variation and inheritance of eastern cottonwood growth and wood properties under two soil moisture regimes. Silvae Genet.
19(1):5-8. FA31:5950.

359. Farmer, R.E., Jr.
1975. Dormancy and root regeneration of northern red oak. Can. J.
For. Ras. 5(2):176–185. BA60:48873.

Farrar, J.L.
 1953. Distribution of radiophosphorus in red pine seedlings. Can.
 For. Branch Silv. Leafl. 78. FA15:3242.

Farrar, J.L., and G.D. Huntly.
 1969. The effect of root pruning on root extension growth of *Pinus banksiana* and *Picea glauca* seedlings. Rep. Res. For. Bot. Glendon Hall Fac. For. Univ. Toronto 1968/69:5-6. FA31:6287.

362. Farrell, E.P., and A.L. Leaf.
1974. Effects of fertilization and irrigation on root numbers in a red pine plantation. Can. J. For. Res. 4(3):366-371. BA59:1960.

363. Fassi, B., and A. Fontana.

1966. Richerche sulle micorrize ectotrofiche del pino strobo in vivaio. II. Micorrize di *Thelephora terrestris* Ehrh. ex Fries, di *Laccaria laccata* (Scop.) Berk. et Br. e di *Hebeloma mesophaeum* Pers. ex Fries. [Researches on ectotrophic mycorrhizae of *Pinus strobus* in nurseries. II. Mycorrhizae of *T. terrestris*, of *L. laccata*, and of *H. mesophaeum*.] Allionia 12:47–53. [Engl. summ.] FA29:1988.

364. Fassi, B., and A. Fontana.

1969. Sintesi micorrizica tra *Pinus strobus* e *Tuber maculatum*. II.

Sviluppo dei semenzali trapiantati e produzione di ascocarpi.

[Mycorrhizal synthesis between *Pinus strobus* and *Tuber maculatum*. II. Development of transplanted seedlings and production of ascocarps.] Allionia 15:115–120. [Engl. summ.] FA32:2205.

365. Fassi, B., and M. Palenzona.
1969. Sintesi micorrizica tra Pinus strobus, Pseudotsuga douglasii ed Endogene lactiflua. [Mycorrhizal associations of Endogene lactiflua with Pinus strobus and Pseudotsuga menziesii.] Allionia 15:105-114. [Engl. summ.] FA32:2204.

366. Fassi, B., and E. de Vecchi.
1962. Richerche sulle micorrize ectotrofiche de pino strobo in vivaio. I. Descrizione di alcune forme piu diffuse in Piemonte.
[Researches in ectotrophic mycorrhizae of *Pinus strobus* in nurseries. I. Description of some of the most common forms in Piedmont.]
Allionia 8:133-152. [Engl. summ.] FA24:4748.

367. Fassi, B., G. Bressy, and T.M.C. Diallo. 1969a. The effect of substrate and fungicides on damping-off of *Pinus strobus* and on root development. Phytopathol. Mediterr., Bologna 8(1):28-40. [In Fr. with Fr., Ital., and Engl. summ.] FA31:2754.

368. Fassi, B., C. Diallo, and M. Palenzona.
1969b. Root rot of *Pinus strobus* caused by *Phytophthora cactorum*.
Ann. Phytopathol. Paris 1(Spec No.):187–192. [In Ital. with Fr. summ.] FA31:4773.

369. Fassi, B., A. Fontana, and J.M. Trappe.
1969c. Ectomycorrhizae formed by *Endogene lactiflua* with species of *Pinus* and *Pseudotsuga*. Mycologia 61:412-413. FA31:2179.

370. Faulkner, M.E., and D.C. Malcolm.
1972. Soil physical factors affecting root morphology and stability of Scots pine on upland heaths. Forestry 45(1):23-36. BA54:60261.
371. Fayle, D.C.F.

1964. Layering habit of sugar maple. For. Chron. 40(1): 116-121.
372. Fayle, D.C.F.

1965. Rooting habit of sugar maple and yellow birch. Can. Dep. For. Publ. 1120. BA47:94198.

373. Fayle, D.C.F.

1966. Root sucker origin on bitternut hickory. Bi-mon. Res. Notes 22(4):2.

374. Fayle, D.C.F.

1968a. Patterns of radial growth in tree roots. Diss. Abstr. 28B(10):4017. FA30:1342.

375. Fayle, D.C.F.

1968b. Radial growth in tree roots: Distribution, timing, anatomy. Univ. Toronto Fac. For. Tech. Rep. 9. BA50:116891.

376. Fayle, D.C.F.

1975a. Archy and diameter of primary xylem in horizontal and vertical roots of red pine. Can. J. For. Res. 5(1):122-129. BA60:9592.

377. Fayle, D.C.F.

1975b. Distribution of radial growth during the development of red pine root systems. Can. J. For. Res. 5(4):608–625.

378. Fayle, D.C.F.

1975c. Extension and longitudinal growth during the development of red pine root systems. Can. J. For. Res. 5(1):109-121. FA36:6137.

379. Fayle, D.C.F.

1976. Stem sway affects ring width and compression wood formation in exposed root bases. For. Sci. 22(2):193–194.

380. Fayle, D.C.F.

1978. Poor vertical root development may contribute to suppression in a red pine plantation. For. Chron. 54(1):99–103.

381. Fayle, D.C.F., and G. Pierpoint.

1975. Evolvement of root and shoot systems of red pine on soils of different moisture availability. *In* Forest soils and forest land management. B. Bernier and C.H. Winget, eds., 4th North Am. For. Soils Conf. Proc., Laval Univ., 1973, Press l'Univ. Laval, Quebec, p. 11-25.

382. Fegel, A.C.

1941. Comparative anatomy and varying physical properties of trunk, branch and root wood in certain northeastern trees. Bull. N.Y. State Coll For. Syracuse Univ., Tech. Publ. 55.

383. Fenton, R.H.

1955. Windthrow a hazard in Virginia pine strip cuttings. USDA For. Serv. Res. Note 53, Northeast. For. Exp. Stn.

384. Fenton, R.H.

1965. Root grafts and translocation of 2, 4, 5–T in young sweetgum stands. J. For. 63(1):16–18. BA46:40689.

385. Fessenden, R.J., R.O. Waito, and R.C. Gregory.

1973. Copper toxicity of jack pine seedlings. Agron. Abstr. 1973:138.

386. Feucht, J.R.

1962. Root initiation and development in air-layered pine and spruce. Diss. Abstr. 23(1):15–16. BA41:3626.

387. Fisher, P.L.

1941. Germination reduction and radicle decay of conifers caused by certain fungi. J. Agric. Res. 62:87–95.

388. Fisher, R.F., Jr.

1969. Increased availability of nitrogen and phosphorus in the root zone of conifers. Diss. Abstr. 29B(11):3993. FA31:2167.

389. Fisher, R.F., Jr., and E.L. Stone.

1968. Soil and plant moisture relations of red pine growing on a shallow soil. Proc. Soil Sci. Soc. Am. 32(5):725-728.

390. Fisher, R.F., Jr., and E.L. Stone.

1969. Increased availability of nitrogen and phosphorus in the root zone of conifers. Proc. Soil Sci. Soc. Am. 33(6):955-961. BA51:63443.

391. Foil, R.R.

1965. The effects of compaction on soil characteristics and seedling growth. Diss. Abstr. 26(6):2955. FA27:5223.

392. Foil, R.R, and C.W. Ralston.

1967. The establishment and growth of loblolly pine seedlings on compacted soils. Proc. Soil Sci. Soc. Am. 31(4):565–568.

393. Fontana, A., and M., Palenzona.

1969. Sintesi micorrizica di *Tuber albidum* in coltura pura, con *Pinus strobus* e pioppo Euroamericano. [Mycorrhizal synthesis of *T. albidum* in pure culture, on *P. strobus* and Euroamerican poplar.] Allionia 15:99–104. [Engl. summ.] FA32:2206.

394. Forbes, R.D.

1925. The roots of trees. Am. For. 31:201-204,234,236.

395. Fortin, J.A.

1970. Interaction entre Basidiomycetes mycorrhizateurs et racines de pine en presence d'acide indol-3y1-acetique. [The interaction between mycorrhiza-forming Basidiomycetes and pine roots in the presence of IAA.] Physiol. Plant. 23(2):365–371. FA31:6072.

396. Foster, N.W., and I.K. Morrison.

1976. Distribution and cycling of nutrients in a natural *Pinus banksiana* ecosystem. Ecology 57(1):110–120.

397. Fowler, R.F.

1970. The white grub in Upper Michigan and northern Wisconsin 1–2 year old red pine plantations: Population-damage correlation, high-grub-hazard area indentification and chemical application techniques. Diss. Abstr. Int. 31B(3):989. FA33:1160.

398. Fowler, R.F., and L.F. Wilson.

1971. White grub populations, *Phyllophaga* spp., in relation to damaged red pine seedlings in Michigan and Wisconsin plantations (Coleoptera: Scarabaeidae). Mich. Entomol. 4(1):23-28. FA34:3000.

399. Fowler, R. F., and L.F. Wilson.

1974. Injury to aldrin-treated and untreated red pine by white grubs (Coleoptera: Scarabaeidae) and other agents during first five years after planting. Great Lakes Entomol. 7(3):81–88. FA37:6903.

400. Fraser, A.I.

1962. The soil and roots as factors in tree stability. Forestry 35(2):117-127. FA24:3804.

401. Fraser, A.I.

1966. Current Forestry Commission root investigations. *In Physiology in forestry*. Soc. For. G.B. 6th Discuss. Meet. Rep., [Edinburgh, 7–9 January 1966], p. 89–93. BA47:104249.

402. Fraser, D.A., and D. McGuire.

1969. Total growth of a black spruce (*Picea mariana*) tree at Chalk River, Ontario, Canada. Can. J. Bot. 47(1):73–84. BA50:77996.

403. Freeman, F.W.

1963. Factors affecting the differential tolerance of tree species to herbicides, primarily simazine. Diss. Abstr. 24(2):456-457. FA25:2322.

404. Froelich, R.C., E.G. Kuhlman, C.S. Hodges, M.J. Weiss, and J.D. Nichols.

1977. Fomes annosus root rot in the South: Guidelines for prevention. USDA For. Serv., August 1977, South. For. Exp. Stn., Southeast. For. Exp. Stn., and Southeast. Area, State and Priv. For.

405. Föelich, H.J., and W. Dietze.

1970. Untersuchungen über Wurzelentwicklung an Pflanzen der Gattung *Populus*, Sektionen Aigeiros, Leuce and Tacamahaca. [Studies on the root development on plants of the genus *Populus*, sections Aigeiros, Leuce and Tacamahaca.] Silvae Genet. 19(4):131–142. [Engl. summ.] BA52:118666.

406. Frothingham, E.H.

1914. White pine under forest management. USDA Bull. 13.

407. Frothingham, E.H.

1915. The eastern hemlock. (*Tsuga canadensis* (Linn.) Carr.). USDA Bull. 152.

408. Fuller, G.D.

1913. Reproduction by layering in the black spruce. Bot. Gaz. 55:452-457.

409. Funk, D.T.

1971. Pot size, pot shape and soil mix all influence black walnut seedling growth. Plant Propagator 17(1):10–14.

410. Gabriel, W.J.

1975. Allelopathic effects of black walnut on white birches. J. For. 73(4):234–237. FA37:168.

411. Gager, C.S.

1929. Aeration of tree roots—thoery. Natl. Shade Tree Conf. Proc. 5:26–27. BA6:6585.

412. Gaiser, R.N.

1947. The growth of loblolly and pond pine seedlings under differing conditions of soil flooding. Unpubl. ms., Dep. Bot., Duke Univ., Durham, N.C.

413. Gaiser, R.N.

1952. Root channels and roots in forest soils. Proc. Soil Sci. Soc. Am. 16(1):62–65. BA26:35837.

414. Gaiser, R.N., and J.R. Campbell.

1951. The concentration of roots in the white oak forests of south-eastern Ohio. USDA For. Serv. Tech. Pap. 120, Cent. States For. Exp. Stn. FA13:1038.

415. Gant, R.E., and E.E.C. Clebsch.

1975. The allelopathic influences of *Sassafras albidum* in old-field succession in Tennessee. Ecology 56(3):604–615. FA37:170.

416. Gardner. F.E.

1929. The relationship between tree age and the rooting of cuttings. Proc. Am. Soc. Hort. Sci. 26:101–104.

417. Garin, G.I.

1942. Distribution of roots of certain tree species in two Connecticut soils. Conn. Agric. Exp. Stn. Bull. 454.

418. Garrett, H.E.

1975. Root initiation and development in sycamore seedlings and cuttings. Tree Plant. Notes 26(3):19–20. FA37:5551.

419. Garrett, H.E.

1977. First-year, root-shoot growth observations of eastern cotton-wood seedlings and cuttings. Tree Plant. Notes 28(1):27–28.

420. Gary, H.L.

1963. Root distribution of five-stamen tamarisk, seepwillow and arrowweed. For. Sci. 9(3):311–314. FA25:1905.

421. Gates, F.C.

1938. Layering in black spruce (*Picea mariana* (Mill.) BSP.). Am. Midl. Nat. 19:589–594.

422. Gevorkiantz, S.R., P.O. Rudolf, and P.J. Zehngraff.

1943. A tree classification for aspen, jack pine, and second-growth red pine. J. For. 41(4):268–274.

423. Gier, L.J.

1945. Development of adventitious roots inside the trunks of trees. Trans. Kans. Acad. Sci. 48:198. BA20:5693.

424. Glifford, G.F.

1966. Aspen root studies on three sites in northern Utah. Am. Midl. Nat. 75(1):132–141. BA47:55568.

425. Gifford, G.F.

1967. The influence of growth media, temperatures, and light intensities on aspen root and top growth. USDA For. Serv. Res. Note INT-67.

426. Gill, C.J.

1970. The flooding tolerance of woody species—a review. For. Abstr. Leading Art. Ser. For. Bur., Oxford 44.

427. Gill, C.J.

1975. The ecological significance of adventitious rooting as a response to flooding in woody species, with special reference to *Alnus glutinosa* (L.) Gaertn. Flora 164(1):85–97. FA37:3836.

428. Gillespie, W.H., and R.E. Adams.

1962. An awl nematode, *Dolichodorus silvestris* n. sp., from West Virginia. Nematologica 8(2):93–98. FA25:1626.

429. Gillgren, I.

1974. [Root tangling, deformation and strangulation.] Skogen 61(1):18–21. [In. Swed.] FA35:5015.

430. Gilmore, A.R.

1957. Physical and chemical characteristics of loblolly pine seedlings associated with drought resistance. 4th South. Conf. For. Tree Improv. Proc., Athens, Ga., [8–9 January 1957], p. 34–39.

431. Gilmore, A.R.

1961. The carbohydrate reserves of loblolly pine (*Pinus taeda* L.) roots in relation to early survival of outplanted seedlings. Diss. Abstr. 21(12):3576. BA37:15545.

432. Gilmore, A.R.

1964. Food reserves of transplanted loblolly pine seedlings and root growth. Tree Plant. Notes 66, p. 15–16. FA26:2102.

433. Gilmore, A.R.

1965. The apparent source of a root growth stimulus in loblolly pine seedlings. Ill. Agric. Exp. Stn. For. Note 112. FA27:161.

434. Gilmore, A.R.

1966. Survival of transplanted yellow-poplar seedlings and root food reserves. Trans. Ill. State Acad. Sci. 59(4):405–406. BA48:46144.

435. Ginns. J.H., and W.H. Gillespie.
1962. *Fomes* root rot found in five thinned native white pine stands

in West Virginia. Plant Dis. Rep. 46(10):734. BA42:15527.
436. Girouard, R.M.
1967. Anatomy of adventitious root formation in stem cuttings.

Proc. Plant Propag. Soc. 17:289–302. FA30:3892.

1955. Tree growth. II. Growth rings and climate. Bot. Rev. 21(1/3):73–188.

438. Goff, E.S.

1897. A study of the roots of certain perennial plants. Wis. Agric. Exp. Stn. Annu. Rep. 14:286–298.

439. Goff, E.S.

1898. The resumption of root growth in spring. Wis. Agric. Exp. Stn. Annu. Rep. 14:220–228.

440. Good, R.E., and N.F. Good.

1975. Growth characteristics of two populations of *Pinus rigida* Mill. from the pine barrens of New Jersey. Ecology 56(5):1215–1220.

441. Good, R.E., and N.F. Good.

1976. Growth analysis of pitch pine seedlings under three temperature regimes. For. Sci. 22(4):445–448.

442. Gosselin, R.

1941. Notes sur le *Polyporus circinatus* Fr. Ann. Ass. Canad.-franc. Sci. 7:57. FA3:339.

443. Goyer, R.A., and D.M. Benjamin.

1972. Influence of soil fertility on infestation of jack pine plantations by the pine root weevil. For. Sci. 18(2):139–147. BA55:30630.

444. Graham, B.F., Jr.

1959. Root-grafts in eastern white pine, *Pinus strobus* L.: Their occurrence and ecological implications. Diss. Abstr. 20(2):466–467. BA35:5369.

445. Graham, B.F., Jr.

1960. Transfer of dye through natural root grafts of *Pinus strobus* L. Ecology 41(1):56–64. BA36:8595.

446. Graham, B.F., Jr., and F.H. Bormann.

1966. Natural root grafts. Bot. Rev. 32(3):255-292. BA48:66913.

447. Graham, B.F., Jr., and A.L. Rebuck.

1958. The effect of drainage on the establishment and growth of pond pine (*Pinus serotina*). Ecology 39(1):33–36. FA20:1798.

148. Graham, S.A.

1958. Problems caused by insects in Lake States forest plantations. Proc. 10th Int. Congr. Entomol., [Montreal 1956], 4:261–272. FA20:3508.

449. Graham, S.A., R.P. Harrison, Jr., and C.E. Westell, Jr.

1963. Aspens: Phoenix trees of the Great Lakes Region. Univ. Mich. Press, Ann Arbor.

450. Grand, L.F., and W.W. Ward.

1969. An antibiotic detected in conifer foliage and its relation to *Cenococcum graniforme* mycorrhizae. For. Sci. 15(3):286–288. FA31:2180.

451. Grano, C.X.

1953. Wind-firmness of shortleaf and loblolly pines. South. Lumberman 187(2345):116.

452. Graves, A.H.

1915. Root rot of coniferous seedlings. Phytopathology 5:213–217.

453. Graves, A.H.

1929. The comparative resistance of root and shoot of the American chestnut to the chestnut bark disease. Nat. Shade Tree Conf. Proc. 5:56–57.

454. Green, R.J., Jr., and R.G. Pratt.

1970. Root rot of black walnut seedlings caused by *Phytophthora citricola*. Plant Dis. Rep. 54(7):583–585. FA32:2708.

455. Green, W.E.

1947. Effect of water impoundment on tree mortality and growth. J. For. 45(2):118–120.

456. Greenidge, K.N.H.

1953. Further studies of birch dieback in Nova Scotia. Can. J. Bot. 31(5):548–559.

457. Creig, B.J.W.

1962. Fomes annosus (Fr.) Cke. and other root-rotting fungi in conifers on ex-hardwood sites. Forestry 35(2):164–182. BA42:19557.

458. Gries, G.A.

1943. Juglone—the active agent in walnut toxicity. North. Nut. Grow. Assoc. Rep. 34:52–55. BA19:17443.

459. Griffin, G.D.

1963. Pathological and ecological relationships between *Xiphinema americanum* Cobb and commercial spruce. Diss. Abstr. 23(12, part 1):4492–4493. FA25:1040.

460. Griffith, B.G., E.W. Hartwell, and T.E. Shaw.

1930. The evolution of soils as affected by the old field white pine-mixed hardwood succession in central New England. Harv. For. Bull. 15.

461. Grisyuk, N.M.

1973. Nektorye osobennosti stroeniya kornevykh sistem derev'ev i kustarnikov semeistva bobovykh. [Some characteristics of the structure of the root system of trees and shrubs of the pea family.] Byull. Gl. Bot. Sada. 87:12–20. [In Russ.] BA58:24061.

462. Grossenbacher, J.G.

1915. The periodicity and distribution of radial growth in trees and their relation to the development of 'annual rings'. Trans. Wis. Acad. Sci., Arts, and Lett. 18:1–77.

463. Gruschow, G.F.

1959. Observations on root systems of planted loblolly pine (*Pinus taeda*). J. For. 57(12):894–896. BA35:19773.

464. Gurskii, A.V.

1928/1929. Kornyevihye sistyemih *Fraxinus excelsior* L., *Fraxinus pennsylvanica* Marsch. i *Acer negundo* L. [The root systems of *F. excelsior*, *F. pennsylvanica* and *A. negundo* on the black soil of Kuban.] Tr. Prikl. Bot., Ghyenetikye i Syelyektsii 21(3):145–183. [Engl. summ.] BA5:11696.

465. Gurskii, G.F.

1939. [Root systems of trees in steppe and desert soils.] Dokl. vsesoyuzn. Akad. s.-kh. Nauk 1939(5/6):45–49. [In Russ.] FA3:112–113.

466. Guttay, A.J.R.

1976. Impact of deicing salts upon the endomycorrhizae of roadside sugar maples. J. Soil Sci. Soc. Am. 40(6):952–954. FA39:146.

467. Gysi, C., C.H. Winget, and B. Bernier.

1975. Interactions of P, Mn, and Cu in nutrient uptake by sugar maple. Can. J. For. Res. 5(1):105–108. BA60:11002.

468. Haas, A.R.C., and H.S. Reed.

1926. The absorption of ions by citrus and walnut seedlings. Hilgardia 2(4):67–106.

469. Habeck, J.R.

1958. White cedar ecotypes in Wisconsin. Ecology 39(3):457–463. FA20:150.

470. Hacskaylo, E.

1951. A study of the roots of *Pinus virginiana* in relation to certain Hymenomycetes suspected of being mycorrhizal. J. Wash. Acad. Sci. 41(12):399–400.

471. Hacskaylo, E.

1959. The role of mycorrhizae in the mineral nutrition of trees. Duke Univ. Sch. For. Bull. 15, Durham, N.C. p. 111–115.

472. Hacskaylo, E.

1962. Research on mycorrhizae in the United States. 13th Int. Union For. Res. Organ. Congr. Proc., [Vienna 1961], Part 2(1), Section 24–6.

473. Hacskaylo, E.

1965. *Thelephora terrestris* and mycorrhizae of Virginia pine. For. Sci. 11(4):401–404. FA27:5553.

474. Hacskaylo, E., and G. Bruchet.

1972. Hebelomas as mycorrhizal fungi. Bull. Torrey. Bot. Club $99(1){:}17{-}20.\ FA34{:}147.$

475. Hacskaylo, E., and J.G. Palmer.

1955. Hymenomycetous species forming mycorrhizae with *Pinus virginiana*. Mycologia 47(1):145–147.

476. Hacskaylo, E., and A.G. Snow, Jr.

1959. Relation of soil nutrients and light to prevalence of mycorrhizae on pine seedlings. USDA For. Serv. Stn. Pap. 125, Northeast. For. Exp. Sta.

477. Hacskaylo, E., J.G. Palmer, and J.A. Vozzo.

1965. Effect of temperature on growth and respiration of ectotrophic mycorrhizal fungi. Mycologia 57(5):748–756. FA27:5548.

478. Hadi, S.

1974. Epidemology and control of *Cylindrocladium* stem canker and root rot of conifers in nurseries. Diss. Abstr. Int. 35B(4):1473–1474. FA37:3867.

479. Haines, B., and G.R. Best.

1976. The influence of an endomycorrhizal symbiosis on nitrogen movement through soil columns under regimes of artificial throughfall and artificial acid rain. *In* First Int. Symp. on Acid Precipitation and the For. Ecosystem Proc., USDA For. Serv. Gen. Tech. Rep. NE–23, p. 951–961.

480. Hall, J.P., P. Singh, and H.O. Schooley.

1971. Survival and growth of some exotic firs in Newfoundland. For. Chron. 47(5):279–281.

481. Hall, T.F., W.T. Penfound, and A.D. Hess.

1946. Water level relationships of plants in the Tennessee Valley with particular reference to malaria control. J. Tenn. Acad. Sci. 21(1):18–59

482. Hall, T.F., and G.E. Smith.

1955. Effects of flooding on woody plants, West Sandy Dewatering Project, Kentucky Reservoir. J. For. 53(4):281–285.

483. Hallmen, U.

1975. Translocation and complex formation of root-applied 2, 4–D and picloram in susceptible and tolerant species. Physiol. Plant. 34(3):266–272. FA38:2301.

484. Hamilton, L.S.

1955. Silvicultural characteristics of American beech. USDA For. Serv. Northeast. Tech. Comm. on the Util. of Beech, Beech Util. Series 13.

485. Hanes, R.E., L.W. Zelazny, and R.E. Blaser.

1970. Effects of deicing salts on water quality and biota; literature review and recommended research. Natl. Coop. Highw. Res. Program Rep. 91, Highw. Res. Bd., Natl. Res. Counc.

486. Hannah, P.R.

1972. Yellow birch root occupancy related to stump and breast height diameters. Vt. Agric. Exp. Stn. Bull. 669. FA34:4461.

487. Hansbrough, T., and J.P. Hollis.

1957. The effect of soil furnigation for the control of parasitic nematodes on the growth and yield of loblolly pine seedlings. Plant Dis. Rep. 41(12):1021–1025. FA19:2929.

488. Hansbrough, T.

1959. The effect of soil fumigation on the growth and yield of loblolly pine seedlings in the nursery. Tree Plant. Notes 37, p. 13–16. FA21:1715.

Hansen, T.S., W.H. Kenety, G.H. Wiggin, and E.C. Stakman.
 1923. A study of the damping-off disease of coniferous seedlings.
 Minn. Agric. Exp. Stn. Tech. Bull. 15.

490. Harley, J.L.

1940. A study of the root system of the beech in woodland soils with especial reference to mycorrhizal infection. J. Ecol. 28(1):107–118. BA14:8074.

491. Harley, J.L., and J.K. Brierley.

1954. The uptake of phosphate by excised mycorrhizal roots of the beech. VI. Active transport of phosphorus from the fungal sheath into the host tissue. New Phytol. 53(2):240–252. BA29:17235.

492. Harley, J.L., C.C. McCready, and J.K. Brierley. 1958. The uptake of phosphate by excised mycorrhizal roots of the beech. VIII. Translocation of phosphorus in mycorrhizal roots. New Phytol. 57(3):353–362. BA33:27482.

493. Hart, J.H.

1965. Root rot of oak associated with *Cylindrocarpon radicicola*. Phytopathology 55(10):1154–1155. BA47:24316.

494. Hartig, T.

1863. Ueber die Zeit des Zuwachses der Bäume. Bot. Zeit. 21:288–289.

495. Hartmann, F.

1962. Root types as site indicator. Can. Dep. For., Ottawa, Transl. by D.C.F. Fayle from Der Waldboden: Humus-, Boden-, und Wurzeltypen als Standortsanzeiger. Osterreichisches Prod. -Zentrum, Vienna 1951, p. 123–152. FA24:3371.

496. Hatch, A.B.

1936. The role of mycorrhizae in afforestation. J. For. 34(1):22-29.

497. Hatch, A.B

1937. The physical basis of mycotrophy in *Pinus*. Black Rock For. Bull. 6.

498. Hatch, A.B., and K.D. Doak.

1933. Mycorrhizal and other features of the root systems of *Pinus*. J. Arnold Arbor. 14(1):85–99. BA8:1837.

499. Hatch, A.B., and C.T. Hatch.

1933. Some hymenomycetes forming mycorrhizae with *Pinus strobus* L. J. Arnold Arbor. 14:324–334.

500. Hatchell, G.E.

1970. Soil compaction and loosening treatments affect loblolly pine growth in pots. USDA For. Serv. Res. Pap. SE-72. BA52:77851.

501. Hathaway, R.L., and D. Penny.

1975. Root strength in some *Populus* and *Salix* clones. N.Z. J. Bot. 13(3):333–344. FA37:4989. BA61:36990.

502. Havis, J.R.

1976. Root hardiness of woody ornamentals. Hort. Sci. 11(4):385–386. BA63:8271.

503. Hawboldt, L.S., and A.J. Skolko.

1948. Investigation of yellow birch dieback in Nova Scotia in 1947. J. For. 46(9):659–671.

504. Hay, R.L.

1969. Effects of planting deformation on growth of loblolly pine seedlings: A study of root deformation. Diss. Abstr. Int. 30B(6):2485. FA32:5942.

505. Hay, R.L., and F.W. Woods.

1968. Distribution of available carbohydrates in planted loblolly pine root systems. For. Sci. 14(3):301–303. BA50:4874.

506. Hay, R.L.

1974. Root deformation correlated with sapling size for loblolly pine. J. For. 72(3):143–145. FA35:7553.

507. Hay, R.L.

1974. Shape of root systems influences survival and growth of loblolly seedlings. Tree Plant. Notes 25(3):1–2.

508. Hay, R.L.

1975. Distribution of carbohydrates in deformed seedling root systems. For. Sci. 21(3):263–267. BA61:25435.

509. Hayes, F.A., and J.H. Stoeckeler.

1935. Possibilities of shelterbelt planting in the Plains region. Section 12—Soil and forest relationships of the shelterbelt zone. USDA For. Serv. Spec. Rep. Lake States For. Exp. Stn., p. 111–153.

510. Heinsdorf, D., and D. Schulzke.

1969. [Distribution of fine roots of young Scots pine and red oaks in a humus-poor sandy soil after deep ploughing and mineral fertilization.] Arch. Forstw. 18(7):731–745. [In Ger. with Russ. and Eng. summ.] FA31:2385.

511. Heinicke, A.J.

1932. The effect of submerging the roots of apple trees at different seasons of the year. Proc. Am. Soc. Hort. Sci. 29:205–207. BA8:11811.

512. Henderson, G.S.

1969. The influence of mycorrhizae on nutrient uptake and growth of conifer seedlings. Diss. Abstr. 29B(11):3993.

513. Hendrix, F.F., Jr., and W.A. Campbell.

1968. Pythiaceous fungi isolated from southern forest nursery soils and their pathogenicity to pine seedlings. For. Sci. 14(3):292–297.

514. Hendrix, F.F., Jr., and W.M. Powell.

1968. Nematode and *Pythium* species associated with feeder root necrosis of pecan trees in Georgia. Plant Dis. Rep. 52(4):334–335. BA49:86278.

515. Hendrix, F.F., Jr., E.G. Kuhlman, C.S. Hodges, Jr., and E.W. Ross. 1964. *Fomes annosus*, a serious threat to regeneration of pine. USDA For. Serv. Res. Note SE-24. FA26:827.

516. Heninger, R.L., and D.P. White.

1974. Tree seedling growth at different soil temperatures. For. Sci. 20(4):363–367. BA60:2056. FA36:5489.

517. Henry, B.W.

1953. A root rot of southern pine nursery seedlings and its control by soil fumigation. Phytopathology 43(2):81–88. FA14:3314.

518. Hepting, G.H.

1971. Diseases of forest and shade trees of the United States. USDA Agric. Handb. 386.

519. Hepting, G.H., and A.A. Downs.

1944. Root and butt rot in planted white pine at Biltmore, North Carolina. J. For. 42(2):119–123.

520. Herrera, S.

1968. Pudricion radicular en plantulas de pino insigne (*Pinus radiata* D. Don), pino oregon (*Pseudotsuga menziesii*) y nogal negro (*Juglans nigra*). [Root rot in seedlings of *P. radiata*, *P. menziesii* and *J. nigra*.] Agric. Tec. (Santiago) 28(1):43-45. [Engl. summ.] BA50:50100.

521. Hewitt, E.J.

1966. A physiological approach to the study of forest tree nutrition. *In* Physiology in forestry. Soc. For. G.B. 6th Discuss. Meet. Rep., [Edinburgh, 7–9 January 1966], p. 49–59.

522. Hicks, R.R., Jr.

1972. The aspen rooting test: A new bioassay. For. Sci. 18(1):21–22.

523. Hicks, R.R., Jr., and W.T. Gladstone.

1971. Some anatomical aspects of rooting quaking aspen. 11th South. Conf. For. Tree Improv. Proc. p. 265–274. FA34:4400.

524. Hilf, H.H.

1927. Wurzelstudien an Waldbäumen. Die Wirzelausbreitung und ihre waldbauliche Bedeutung. [Root systems of forest trees and their silvicultural significance.] M. and H. Schaper, Hanover. BA3:9052.

Hilton, R.J., and H. Khatamian.
 1973. Diurnal variation in elongation rates of roots of woody plants.
 Can. J. Plant Sci. 53(3):699-700. FA35:3483.

526. Himelick, E.B.

1959. Experimental control studies on the oak wilt disease. Ph.D. thesis, Univ. Ill., Urbana.

527. Himelick, E.B.

1959. Experimental control studies on the oak wilt disease. Diss. Abstr. 20(5):1554-1555. FA21:3365.

528. Himelick, E.B., and D. Neely. 1962. Root grafting of city-planted American elms. Plant Dis. Rep. 46(2):86-87. BA40:8155.

529. Hemelick, E.B., and D. Neely. 1962. Frequency of root grafting between city-planted American elms. Arborist's News 27(4):29–31. BA41:20333.

530. Himes, W.E, and J.M. Skelly.
1972. An association of the black turpentine beetle, *Dendroctonus terebrans*, and *Fomes annosus* in loblolly pine. (Abstr.) Phytopathology 62(6):670. FA34:1132.

531. Hock, W.K., and W.L. Klarman.
1966. Increased disease resistance of Virginia pine seedlings caused by morphological changes in maturing root tissue. (Abstr.) Phytopathology 56(6):584. FA28:779.

532. Hocking, D., and D.L. Mitchell.
1975. The influences of rooting volume-seedling escapement and substratum density on greenhouse growth of lodgepole pine (*Pinus contorta*), white spruce (*Picea glauca*), and Douglas fir (*Pseudotsuga menziesii*) grown in extruded peat cylinders. Can. J. For. Res. 5(3):440-451.

533. Hodges, C.S., Jr.
1962. Black root rot of pine seedlings. Phytopathology 52(3):210–219. FA23:5401.

534. Hodges, C.S., Jr.
1963. Black root rot of pine. *In* Symposium on root diseases of forest trees, Corvallis, Oregon 1962. Phytopathology 53(10):1132–1134.

BA45:57445

535. Hodges, C.S., Jr.
1964. The effect of competition by *Peniophora gigantea* on the growth of *Fomes annosus* in stumps and roots. (Abstr.) Phytopathology 54(6):623. FA26:844.

536. Hodges, C.S., Jr.
1969. Relative susceptibility of loblolly, longleaf, and slash pine roots to infection by *Fomes annosus*. (Abstr.) Phytopathology 59(8):1031. FA31:2786.

537. Hodges, C.S., and E. Jorgensen.
1967. Fomes annosus root rot Fomes annosus (Fr.) Cke. In Important forest insects and diseases of mutual concern to Canada, the United States, and Mexico. A.G. Davidson and R.M. Prentice, eds., Can. For. Serv. Publ. 1180, p. 49–52.

538. Hodges, C.S., and E.G. Kuhlman.
1974. Spread of *Fomes annosus* in roots of redcedar and loblolly pine. Plant Dis. Rep. 58(3):282-284. FA35:7001.

539. Hodges, C.S., and L.C. May.
1972. A root disease of pine, araucaria and eucalyptus in Brazil
caused by a new species of *Cylindrocladium*. Phytopathology
62(8):898-901. FA34:1726.

540. Hoffmann, G.
1964. [The effectiveness and host specificity of the nodule bacteria of *Robinia pseudoacacia*.] Arch. Forstw. 13(6):563–576. [In Ger. with Russ. and Engl. summ.] FA26:294.

541. Hoffmann, G.
1966. The influence of soil heating on root growth of black locust (Robinia pseudoacacia L.). Pap. 4th World For. Congr., [Madrid].

542. Hoffmann, G. 1968. [The effect of soil heating on the growth of roots and shoots of Robinia pseudoacacia.] Arch. Forstw. 17(4):431-435. [In. Ger. with Russ. and Engl. summ.] FA30:297. 543. Hoffmann, G.
1972. [The growth rhythms of roots and shoot axis in forest trees.]
Flora 161(3):303-319. [In Ger. with Engl. summ.]

544. Holch, A.E.
1931. Development of roots and shoots of certain deciduous tree seedlings in different forest sites. Ecology 12(2):259-298.
BA8:16267.

545. Hollett, B.P., and M.T. Jackson.
1976. Quantitative aspects of the association of *Cenococcum graniforme* with *Fagus grandifolia* in Indiana. For. Sci. 22(2):127–130. FA38:1208.

546. Holm, T. 1909. Nyssa sylvatica Marsh. Am. Midl. Nat. 1:128-137.

 Holm, T.
 1921. Morphological study of Carya alba and Juglans nigra. Bot. Gaz. 72:375–388.

548. Holm, T.
1931. The seedling of *Hamamelis virginiana* L. Rhodora 33(388):81-92.

549. Hong, S.G. 1975. Test of hypothesis that cytokinin will stimulate potassium ion transport in roots of honeylocust (*Gleditsia triacanthos* L.). Diss. Abstr. Int. 35B(7):3536. FA37:3478.

550. Hong, S.G., and S.K. Hyun. 1970. Anatomical investigation on root formation in hypocotyl cuttings of pines. Res. Rep. Inst. For. Genet. Korea 8, p. 15-22. FA33:2472.

551. Hook, D.D. 1969. Growth and development of swamp tupelo (Nyssa sylvatica var. biflora (Walt.) Sarg.) under different root environments. Diss. Abstr. 29B(12, part 1):4549. FA31:2139.

Hook, D.D.1974. Root (botany). McGraw-Hill Yearb. of Sci. and Technol.McGraw-Hill, New York.

553. Hook, D.D., and C.L. Brown.
1973. Root adaptations and relative flood tolerance of five hardwood species. For. Sci. 19(3):225–229. BA57:43014.

Hook, D.D., C.L. Brown, and P.P. Kormanik.
 1970a. Lenticel and water root development of swamp tupelo under various flooding conditions. Bot. Gaz. 131(3):217–224.

 Hook, D.D., P.P. Kormanik, and C.L. Brown.
 1970b. Early development of sweetgum root sprouts in coastal South Carolina. USDA For. Serv. Res. Pap. SE-62.

Hook, D.D., O.G. Langdon, J. Stubbs, and C.L. Brown.
 1970c. Effect of water regimes on the survival, growth, and morphology of tupelo seedlings. For. Sci. 16(3):304–311.

557. Hook, D.D., C.L. Brown, and P.P. Kormanik.
1971. Inductive flood tolerance in swamp tupelo (*Nyssa sylvatica* var. *biflora* (Walt.) Sarg.). J. Exp. Bot. 22(70):78–89.

Hook, D.D., C.L. Brown, and R.H. Wetmore.
 1972. Aeration in trees. Bot. Gaz. 133(4):443–454.

559. Hopkins, H.T., Jr., and R.L. Donahue.
1939. Forest tree root development as related to soil morphology.
Proc. Soil Sci. Soc. Am. 4:353. BA15:6161.

560. Hord, H.H.V., and M.J. Hildebrand.
1956. Armillaria mellea in relation to regeneration of red pine, white pine, and white spruce. Bi-mon. Progr. Rep. Div. For. Biol., Dep. Agric. Can. 12(1):2. FA17:4088.

561. Horsley, S.B.
1971. Root tip injury and development of the paper birch root system. For. Sci. 17(3):341–348. BA53:16386.

Horsley, S.B., and B.F. Wilson.
 1971. Development of the woody portion of the root system of Betula papyrifera. AM. J. Bot. 58(2):141-147. BA52:51455.

 Horton, K.W., and E.J. Hopkins.
 1965. Influence of fire on aspen suckering. Can. For. Serv. Publ. 1095. FA27:330. 564. Horton, K.W., and J.C. Lees.

1961. Black spruce in the foothills of Alberta. Can. For. Res. Branch Tech. Note. 110.

565. Hosner, J.F.

1959. Survival, root and shoot growth of six bottomland tree species following flooding. J. For. 57(12):927-928. BA35:19774.

Hosner, J.F., and S.G. Boyce.

1962. Tolerance to water saturated soil of various bottomland hardwoods. For. Sci. 8(2):180-186.

567. Hosner, J.F., and D.L. Graney.

1970. The relative growth of three forest tree species on soils associated with different successional stages in Virginia. Am. Midl. Nat. 84(2):418-427. FA32:3994.

568. Hosner, J.F., and A.L. Leaf.

1962. The effect of soil saturation upon the dry weight, ash content, and nutrient absorption of various bottomland tree seedlings. Proc. Soil Sci. Soc. Am. 26(4):401-404. FA24:1768.

569. Hosner, J.F., A.L. Leaf, R. Dickson, and J.B. Hart, Jr. 1964. Effects of varying soil moisture upon the nutrient uptake of four bottomland tree species. Agron. Abstr. 1964:53.

Hough, A.F.

1951. Tree roots. Pa. For. 36(346):22-26.

Hough, A.F., and R.D. Forbes.

1943. The ecology and silvics of forests in the high plateaus of Pennsylvania. Ecol. Monogr. 13:299-320.

Houston, D.R., and H.G. Eno.

1969. The use of soil fumigants to control spread of Fomes annosus. USDA For. Serv. Res. Pap. NE-123.

573. Howe, V.K.

1973. Site changes and root damage. Proc. 49th Int. Shade Tree Conf., p. 25-28.

574. Howell, F.C., and W.J. Stambaugh.

1972. Rates of pathogenic and saprophytic development of Fomes annosus in roots of dominant and suppressed eastern redcedar. Plant Dis. Rep. 56(11):987-990. BA55:51255.

575. Hoyle, M.C.

1965. Addition of phosphorus to subsoil promotes root development of yellow birch. USDA For. Serv. Res. Note NE-42. FA28:289.

1970. Growth and nutrition of yellow birch as affected by the nutrient status of a podzol soil. In Tree growth and forest soils. C.T. Youngberg, and C.B. Davey, eds., 3rd North Am. For. Soils Conf. Proc., N.C. State Univ., Raleigh, [August 1968], Oreg. State Univ. Press, Corvallis. p. 221-233.

577. Hoyle, M.C.

1971. Effects of the chemical environment on yellow birch root development and top growth. Plant and Soil 35(3):623-633. BA54:39320.

578. Huberman, M.A.

1940. Normal growth and development of southern pine seedlings in the nursery. Ecology 21(3):323-334.

Hudak, J., and P. Singh.

1970. Incidence of Armillaria root rot in balsam fir infested by balsam woolly aphid. Can. Plant Dis. Surv. 50(3):99-101.

580. Hunt, F.M.

1951. Effects of flooded soil in growth of pine seedlings. Plant Physiol. 26(2):363-368.

581. Hunt, T.N., and M.H. Farrier.

1974. Oviposition and feeding preference of pales weevil (Coleoptera: Curculionidae) for five types of loblolly pine bark. Ann. Entomol. Soc. Am. 67(3):407-408. FA36:1025.

582. Huntly, J.H., J.D. Cafley, and E. Jorgensen.

1961. Armillaria root rot in Ontario. For. Chron. 37(3):228-232. BA37:24623.

583. Husch, B.

1959. Height growth of white pine in relation to selected environmental factors on four sites in southeastern New Hampshire. N.H. Agric. Exp. Stn. Tech. Bull. 100. FA21:2842.

584. Hutchinson, M.T., and J.P. Reed.

1959. The pine cystoid nematode in New Jersey. Plant Dis. Rep. 43(7):801-802. FA23:956.

585. Hutnik, R.J.

1954. Effect of seedbed condition on paper birch reproduction. J. For. 52(7):493-495.

586. Hutnik, R.J.

1964. Functionality of root grafts between scarlet oak trees. (Abstr.) Bull. Ecol. Soc. Am. 45(4):150-151. FA26:3449.

587. Ibberson, J.E.

1949. A preliminary report on the diseased condition of red pine in southeastern Pennsylvania. Pa. For. and Waters 1(6):124-125, 141. FA11:2268.

588. Ike, A.F., and E.L. Stone.

1956. Soil nitrogen accumulation and availability under black locust stands in New York. Agron. Abstr. 1956:50.

589. Ike, A.F., and E.L. Stone.

1958. Soil nitrogen accumulation under black locust. Proc. Soil Sci. Soc. Am. 22(4):346-349.

590. Illick, J.S., and E.F. Brouse.

1926. The Ailanthus tree in Pennsylvania. Pa. Dep. For. and Waters

591. International Symposium: Ecology and Physiology of Root Growth. 1974. [Potsdam, 1971] Akademie-Verlag, Berlin. FA35:7455.

592. Iverson, T.H., and K. Siegel.

1976. The geotropic curvature in roots of Norway spruce (Picea abies) containing anthocyanins. Physiol. Plant. 37(4):283-287. BA63:29165.

593. Iyer, J.G., and S.A. Wilde.

1974. Micronutrients in tree nursery soils: Their behavior, and importance, and an appraisal of their deficiencies. Soil Sci. 118(4):267-269. BA59:14296.

594. Jackson, L.W.R.

1938. Cylindrocladium associated with diseases of tree seedlings. Plant Dis. Rep. 22:84-85.

595. Jackson, L.W.R.

1945. Root defects and fungi associated with the little-leaf disease of southern pines. Phytopathology 35(2):91-105.

596. Jackson, L.W.R.

1970. Nematode parasitic on shortleaf pine roots. Plant Dis. Rep. 54(6):465-466. FA32:1204.

597. Jackson, L.W.R., and G.H. Hepting.

1964. Rough bark formation and food reserves in pine roots. For. Sci. 10(2):174-179. BA45:84175.

598. Jagels, R.

1963. Gelatinous fibers in the roots of quaking aspen. For. Sci. 9(4):440-443. BA45:52872.

599. Janouch, K.L.

1927. Effect of spacing and root pruning on the development of transplants. J. For. 25(1):62-67.

600. Jeffrey, W.W.
1959. White spruce rooting modifications on the fluvial deposits of the lower Peace River. For. Chron. 35(4):304-311. FA21:2857.

601. Jensen, C.A.

1907. Some mutual effects of tree-roots and grasses on soils. Science (N.S.) 25:871-874.

602. Johansson, M., and O. Theander.

1974. Changes in sapwood of roots of Norway spruce, attacked by Fomes annosus. Part I. Physiol. Plant. 30(3):218-225. BA58:27755.

603. Johnson, A.W., T.J. Ratcliffe, and G.C. Freeman. 1970. Control of *Meloidogyne incognita* on dogwood seedlings by chemical dips. Plant Dis. Rep. 54(11):952-955. FA32:6548.

604. Johnson, J.R., and J.R. Havis.

1977. Photoperiod and temperature effects on root cold acclimation. J. Am. Soc. Hort. Sci. 102(3):306-308. FA39:169.

605. Johnson, R.L., and R.M. Krinard.

1976. Hardwood regeneration after seed tree cutting. USDA For. Serv. Res. Pap. SO-123.

606. Johnston, H.R., and C.B. Eaton.

1939. White grubs in forest nurseries of the Carolinas. USDA Bur. Entomol. and Plant Quar. E-486. FA2:41.

607. Johnston, W.F.

1970. Planting black spruce on brushy lowland successful if done in unshaded sphagnum. Tree Plant. Notes 21(3):20-22. FA32:733.

1968. A note on the dimensions of shoots and roots of planting stock. Forestry 41(2):199-206. BA50:66842.

609. Jones, J.R.

1974. Aspen sucker growing from an Engelmann spruce stump. USDA For. Serv. Res. Note RM-264. FA36:6145.

610. Jones, K.L.

1967. Observations on root tips and associated microorganisms in soil. Pap. Mich. Acad. Sci. 52:43-54. FA30:340.

611. Jones, L.R., and W.J. Morse.

1903. The shrubby cinquefoil as a weed. Vt. Agric. Exp. Stn. Annu. Rep. 16:173-190.

612. Jones, T.W.

1961. First report of pine mortality caused by Fomes annosus root rot in Ohio. Plant Dis. Rep. 45(12):980. BA38:11588.

613. Jones, T.W., and A.D. Partridge.

1961. The importance of root grafts in oak wilt spread in Missouri. Plant Dis. Rep. 45(7):506-507. BA37:3626.

614. Jordan, M.J.

1975. Effects of zinc smelter emissions and fire on a chestnut-oak woodland. Ecology 56(1):78-91. FA36:7027.

615. Jorgensen, E.

1959. Fomes annosus root rot. In Recent advances in botany. Proc. 9th Int. Bot. Congr. [Montreal 1959] 2:40-41. BA36:82414.

616. Jorgensen, E.

1961. On the spread of Fomes annosus (Fr.) Cke. Can. J. Bot. 39(6):1437-1445. FA23:3834.

617. Jorgensen, J.R.

1968. Root growth of direct-seeded southern pine seedlings. USDA For. Serv. Res. Note SO-79.

618. Kahn, M.S.

1969. Fungal associates of Juglans nigra L. Diss. Abstr. 29B(12, part 1):4471.

619. Kalela, E.K.

1950. Männiköiden ja kuusikoiden juurisuhteista I. [On the horizontal roots in pine and spruce stand I.] Acta For. Fenn. 57(2):1-79. [Engl. summ.] BA26:9774.

620. Kalisz, S.P., nd J.H. Brown, Jr.

1976. Starch content of oak roots on campsites. Sci. Biol. J. 2(4):160-165. BA63:11170.

621. Katzman, G.B.

1971. White spruce in northern New York root by layering. Tree Plant. Notes 22(4):15-16.

622. Katznelson, H., J.W. Rouatt, and E.A Peterson.

1962. The rhizosphere effect of mycorrhizal and non-mycorrhizal roots of yellow birch seedlings. Can. J. Bot. 40(3):377-382. BA39:7722.

623. Kaufman, C.M.

1945. Root growth of jack pine on several sites in the Cloquet Forest, Minnesota. Ecology 26(1):10-23. BA19:7950.

624. Kaufmann, M.R.

1968. Water relations of pine seedlings in relation to root and shoot growth. Plant Physiol. 43(2):281-288. FA30:305.

625. Kearby, W.H.

1966. The bionomics of the pine root weevil Hylobius rhizophagus Millers and the weevil complex associated with plantation deterioration. Diss. Abstr. 26(9):4978-4979. FA28:989.

626. Kearby, W.H., and D.M. Benjamin.

1969. Life history and damage of the pine root tip weevil, Hylobius rhizophagus, in Wisconsin. Ann. Entomol. Soc. Am. 62(4):838-843. BA51:17021.

627. Keller, T.

1967. Beitrag zur Kenntnis der Wurzelatmung von Koniferenjungpflanzen. [Contribution to the knowledge of root respiration of young conifers.] 14th Int. Union For. Res. Organ. Congr. Proc., [Munich 1967], Part IV, Section 23, p. 329-340. [Engl. and Fr. summ.] FA29:1846.

628. Kelley, A.P.
1937. The form and occurrence of mycorrhizae in the genus *Popu*lus. Landenberg Rev. 1937:54-56. FA1:76.

629. Kelley, A.P.

1941. The variations in form of mycorrhizal short roots of Pinus virginiana Mill. associated with certain soil series. Landenberg Lab., Landenberg, Pa. FA3:20.

630. Kelley, A.P.

1943. Note on the mycorrhizae of Pinus virginiana Mill. Landenberg Lab., Landenberg, Pa. FA5:11.

631. Kelley, A.P.

1950. Mycotrophy in plants. Chron. Bot. Co., Waltham, Mass.

632. Kelley, A.P.

1960. The root endings of beech, maple, and dogwood as found in the eastern U.S.A. Folia for. polon. (Lesn.) 4, p. 45-88 FA23:237.

633. Kelman, A., and G.V. Gooding, Jr.

1965. A root and stem rot of yellow-poplar caused by Cylindrocladium scoparium. Plant Dis. Rep. 49(9):797-801. BA47:24326.

634. Kelman, A., G.V. Gooding, Jr., and G.K. Slocum.

1959. Cylindrocladium root rot of yellow-poplar (Liriodendron tulipifera L.). J. Elisha Mitchell Sci. Soc. 75(2):66-67. BA36:66068.

635. Kennedy, G.E.

1937. Root growth on white oak acorns. USDA For. Serv. R-9 Bull. 10(3):7.

636. Keresztesi, B.

1955. Néhány adat az akác gyökérrendszeréről. [Some data on the root system of Robinia pseudoacacia.] Erdö 4(3):113-121. FA17:2558.

637. Kessler, K.J.

1966. Growth and development of mycorrhizae of sugar maple (Acer saccharum Marsh.). Can. J. Bot. 44(10):1413-1425.

Khashes, Ts. M.

1975. Determining the viability of root systems of seedlings. Inst. Pap. Chem., Appleton, Wis. [Engl. transl.] FA37:3740.

Khatamian, H.

1971. Seasonal and diurnal root growth patterns of certain woody plants. M.S. thesis, Univ. Guelph, Ontario, Canada.

640. Khatamian, H., and R.J. Hilton.

1971. Development of ectomycorrhiza on red pine roots. Can. J. For. Res. 1(4):269-272. BA53:68898.

641. Kienholz, R.

1934. Leader, needle, cambial, and root growth of certain conifers and their inter-relations. Bot. Gaz. 96(1):73-92. BA9:10142.

642. Kinden, D.A.

1975. An ultrastructural study of vesicular-arbuscular mycorrhizae on yellow-poplar. Diss. Abstr. Int. 35B(9):4310. FA36:7534.

643. Kirby, C.S., and H.G. McPhee.

1963. Some field observations on the cicada, *Okanagana rimosa* (Say). Bi-mon Progr. Rep., For. Entomol. Pathol. Branch, Dep. For. Can. 19(5):1. FA25:2431.

644. Kirby, H.W., and L.F. Grand.

1975. Susceptibility of *Pinus strobus* and *Lupinus* spp. to *Phytophthora cinnamomi*. Phytopathology 65(6):693–695. FA37:1698.

645. Kitteredge, J., and S.R. Gevorkiantz.

1929. Forest possibilities of aspen lands in the Lake States. Minn. Agric. Exp. Stn. Tech. Bull. 60.

646. Kochenderfer, J.N.

1973. Root distribution under some forest types native to West Virginia. Ecology 54(2):445–448. BA56:48300.

647. Koenigs, J.W.

1960. Fomes annosus: A bibliography with subject index. USDA For. Serv. Occ. Pap. 181. South. For. Exp. Stn. FA23:761.

648. Koenigs, J.W.

1971. Borax: Its toxicity to *Fomes annosus* in wood and its diffusion, persistence, and concentration in treated stumps of southern pines. Phytopathology 61(3):269–274. FA32:6269.

649. Kolesnikov, V.A.

1930. The root system of fruit tree seedlings. J. Pomol. and Hort. Sci. 8(3):197–203. BA6:25384.

650. Korać, M.

1970. Competition and root systems coalescence in nest seeding of some forest tree species with a special account of beech treatment. Silvae Genet. 19(2/3):99–101. FA32:345.

651. Kormanik, P.P., and C.L. Brown.

1967. Root buds and the development of root suckers in sweetgum. For. Sci. 13(4):338–345. BA49:27685.

652. Korstian, C.F.

1927. Factors controlling germination and early survival in oaks. Yale Univ. Sch. For. Bull. 19, p. 7–115. BA2:18493.

653. Korstian, C.F., and T.S. Coile.

1938. Plant competition in forest stands. Duke Univ. Sch. For. Bull. 3, Durham, N.C.

654. Köstler, J.N.

1962. Untersuchungen zur Wurzelbildung. [Investigations on root formation.] Allg. Forst Z. 28:413–417.

655. Kovaleva, A.N.

1972. Kornevaya sistema drevesnykh i kustarnikovykh rastenii. Ukazatel'otechestvennoi i inostrannoi literatury. Chast' I 1900–1965. [The root system of trees and shrubs. A bibliography of Russian and foreign literature. Part I 1900–1965.] Moscow, USSR, MGU. FA34:5644.

656. Kozlowski, T.T.

1949. Light and water in relation to growth and competition of Piedmont forest tree species. Ecol. Monogr. 19(3):207–231.

657. Kozlowski, T.T.

1968. Growth and development of *Pinus resinosa* seedlings under controlled temperatures. Adv. Front. Plant Sci. 19:17–27. FA 30:5345

658. Kozlowski, T.T., and J.H. Cooley.

1960. Natural root grafting in forest trees. Univ. Wis. For. Res. Notes 56. FA22:254.

659. Kozlowski, T.T., and J.H. Cooley.

1961. Natural root grafting in northern Wisconsin. J. For. 59(2):105-107. BA36:38773.

660. Kozlowski, T.T., and W.L. Scholtes.

1948. Growth of roots and root hairs of pine and hardwood seedlings in the Piedmont. J. For. 46(10):750–754. BA23:19598.

661. Kozlowski, T.T., and C.H. Winget.

1961. Patterns of water uptake in forest trees. Univ. Wis. For. Res. Notes 72.

662. Kozlowski, T.T., and C.H. Winget.

1964a. Contributions of various plant parts to growth of pine shoots. Univ. Wis. For. Res. Notes 113.

663. Kozlowski, T.T., and C.H. Winget.
1964b. The role of reserves in leaves, branches, stems, and roots on shoot growth of red pine. Am. J. Bot. 51(5):522–529. BA45:88632.

664. Kramer, P.J.

1933. The intake of water through dead root systems and its relation to the problem of absorption by transpiring plants. Am. J. Bot. 20:481–492.

665. Kramer, P.J.

1942. Species differences with respect to water absorption at low soil temperatures. Am. J. Bot. 29:828–832. FA5:90.

666. Kramer, P.J.

1946. Absorption of water through suberized roots of trees. Plant Physiol. 21(1):37–41. FA8:896.

667. Kramer, P.J.

1951a. Causes of injury to plants resulting from flooding of the soil. Plant Physiol. 26(4):722–736.

668. Kramer, P.J.

1951b. Effects of respiration inhibitors on accumulation of radioactive phorsphorus by roots of loblolly pine. Plant Physiol. 26(1):30–36. FA13:1853.

669. Kramer, P.J., and H.C. Bullock.

1966. Seasonal variations in the proportions of suberized and unsuberized roots of trees in relation to the absorption of water. Am. J. Bot. 53(2):200–204. FA28:316.

670. Kramer, P.J., and R.H. Hodgson.

1954. Differences between mycorrhizal and non-mycorrhizal roots of loblolly pine. Proc. 8th Int. Bot. Congr., [Paris 1954], Part 5, Section 13, p. 133–134. FA16:222.

671. Kramer, P.J., and H.H. Wiebe.

1952. Longitudinal gradients of P³² absorption in roots. Plant Physiol. 27(4):661–674. BA27:7686.

672. Kramer, P.J., and K.M. Wilbur.

1949. Absorption of radioactive phosphorus by mycorrhizal roots of pine. Science 110(2844):8–9.

673. Kramer, P.J., W.S. Riley, and T.T. Bannister.

1952. Gas exchange of cypress knees. Ecology 33(1):117–121.

674. Krebill, R.G.

1962. Etiology of a jack and red pine plantation decline characterized by root weevil injury and fungal deterioration. Diss. Abstr. 23(6):1861. FA24:4010.

675. Krebill, R.G., and R.F. Patton.

1962. Wounds in jack pine (*P. banksiana*) roots as entry points for a succession of fungi. (Abstr.) Phytopathology 52(8):739. BA41:24683.

676. Krebill, R.G., K.R. Barker, and R.F. Patton.

1967. Plant-parasitic nematodes of jack and red pine stands in Wisconsin. Nematologica 13(1):33–42. FA28:6070.

677. Kriebel, H.B.

1963. Selection for drought resistance in sugar maple. World Consult. on For. Genet. and Tree Improv., Section 3, FAO, Stockholm, Sweden.

678. Krupa, S., and J.E. Nylund.

1972. Studies on ectomycorrhizae of pine. III. Growth inhibition of two root pathogenic fungi by volatile organic constituents of ectomycorrhizal root systems of *Pinus sylvestris* L. Eur. J. For. Pathol. 2(2):88–94. BA54:68650.

679. Krupa, S., J. Andersson, and D.H. Marx.

1973. Studies on ectomycorrhizae of pine. IV. Volatile organic compounds in mycorrhizal and nonmycorrhizal root systems of *Pinus echinata* Mill. Eur. J. For. Pathol. 3(4);194–200. FA36:694.

680. Krywolap, G.N., L.F. Grand, and L.E. Casida, Jr.

1964. The natural occurrence of an antibiotic in the mycorrhizal fungus *Cenococcum graniforme*. Can. J. Microbiol. 10(3):323–328. FA26:299.

681. Kuhlman, E.G. 1969a. Variation in susceptibility of some forest tree seedlings to infection by *Fomes annosus*. (Abstr.) Phytopathology 59(8):1036. FA31:2785.

682. Kuhlman, E.G.

1969b. Inoculation of loblolly pine seedlings with *Fomes annosus* in the greenhouse. Can. J. Bot. 47(12):2079–2082. FA31:6653.

683. Kuhlman, E.G.

1969c. Number of conidia necessary for stump root infection by *Fomes annosus*. Phytopathology 59(8):1168–1169. FA31:2783.

684. Kuhlman, E.G.

1970a. Decomposition of loblolly pine bark and soil- and root-inhabiting fungi. Can. J. Bot. 48(10):1787–1793. BA52;16343.

685. Kuhlman, E.G. 1970b. Seedling i

1970b. Seedling inoculations with *Fomes annosus* show variation in virulence and in host susceptibility. Phytopathology 60(12):1743–1746. FA32:6249.

686. Kuhlman, E.G.

1972. Susceptibility of loblolly and slash pine progeny to *Fomes annosus*. USDA For. Serv. Res. Note SE-176. FA34:5240.

687. Kuhlman, E.G.

1973. Rate of infection of loblolly pine roots on high- and low-hazard sites by *Fomes annosus*. (Abstr.) Phytopathology 63(4):444. FA35:223.

688. Kuhlman, E.G., C.S. Hodges, Jr., and R.C. Froelich. 1976. Minimizing losses to *Fomes annosus* in the southern United States. USDA For. Serv. Res. Pap. SE–151.

689. Kuntz, J.E.

1961. Oak wilt: Its development, spread, and control. *In* Recent advances in botany, 9th Int. Bot. Congr. [Montreal 1959] 2:1544–1548.

690. Kuntz, J.E., and A.J. Riker.

1950a. Oak wilt in Wisconsin. Wis. Conserv. Bull. 15(6): 20–23. FA12:3389.

691. Kuntz, J.E., and A.J. Riker.

1950b. Root grafts as a possible means for local transmission of oak wilt. (Abstr.) Phytopathology 40(1):16–17. FA11:3293.

692. Kuntz, J.E., and A.J. Riker.

1950c. The translocation of poisons between oak trees through natural root grafts. 7th Annu. North Cent. Weed Control Conf. Res. Rep., Milwaukee, Wis., p. 242. FA13:1208.

693. Kuntz, J.E., and A.J. Riker.

1956. The use of radioactive isotopes to ascertain the role of root grafting in the translocation of water, nutrients, and disease-inducing organisms among forest trees. Int. Conf. Peaceful Uses Atomic Energy Proc. [Geneva 1955] 12:144–148. FA18:1283.

694. Kurz, H., and K. Wagner.

1952. The role of adventitious roots in survival of cypress. (Abstr.) J. Tenn. Acad. Sci. 27(3):201. FA14:3168.

695. Lachance, D.

1974. Développement de *Odontia bicolor* chez le sapin baumier infecté artificiellement. [Development of *O. bicolor* in artificially infected balsam fir.] Can. J. For. Res. 4(3):327–334. [Engl. summ.]

696. Lachance, D.

1975. Inoculations of balsam fir with *Odontia bicolor:* Early establishment and associated microorganisms. Can. J. For. Res. 5(1):130–138. FA36:6381.

697. Lachance, D.

1976. Étude de carie dans une plantation d'épinettes blanches fertilisées. [Study of decay in a fertilized white spruce plantation.] Rapp. d'Inf. Cent. Rech. For. des Laurentides. LAU-X-18.

698. Ladefoged, K.

1948. Analysis of the root sap of the birch. Plant and Soil 1(2):127–134. BA23:19647.

699. LaFond, A.

1950. The rate of respiration of jack pine root tips as influenced by extracts from different types of humus. Proc. Soil Sci. Soc. Am. 15:357–359. BA26:2105.

700. (Laing, E.V.)

1932. Studies on tree roots. (G. B.) For. Comm. Bull. 13. BA8:19453.

701. Lampky, J.R., and J.E. Peterson.

1963. *Pisolithus tinctorius* associated with pines in Missouri. Mycologia 55(5):675–678. FA25:4572.

702. Lane, C.L., and A.T. Shearin.

1972. Aldrin affects nematodes and mycorrhizal growth on loblolly pine seedlings. For. Sci. 18(4):319–320. FA34:5305.

703. Lane, R.D., and A.L. McComb.

1953. Effects of grass competition upon the establishment of hardwood plantations in Iowa. Ia. Agric. Exp. Stn. Res. Bull. 399, p. 433–459. FA15:1175.

704. Larsen, J.A.

1935. Natural spreading of planted black locust in southeastern Ohio. J. For. 33(4):616–619.

705. Larson, M.M.

1970. Root regeneration and early growth of red oak seedlings: Influence of soil temperature. For. Sci. 16(4): 442-446. BA52:36841.

706. Larson, M.M.

1974. Effects of temperature on early growth of oak seedlings. *In* Forestry research review—1974. Agric. Res. and Dev. Cent., Wooster, Ohio. p. 6–9. FA36:3212.

707. Larson, M.M.

1974. Effects of soil moisture on early growth of oak seedlings. *In* Forestry research review—1974. Agric. Res. and Dev. Cent., Wooster, Ohio, p. 10–13. FA36:4524.

708. Larson, M.M., and I. Palashev.

1973. Effects of osmotic water stress and gibberellic acid on initial growth of oak seedlings. Can. J. For. Res. 3(1):75–82. BA56:48890.

709. Larson, M.M., and F.W. Whitmore.

1970. Moisture stress affects root regeneration and early growth of red oak seedlings. For. Sci. 16(4):495–498. BA52:36845.

710. La Rue, C.D.

1934. Root grafting in trees. Am. J. Bot. 21(3):121-126. BA8:20838.

711. Laycock, W.A.

1967. Distribution of roots and rhizomes in different soil types in the pine barrens of New Jersey. U.S. Geol. Surv. Prof. Pap. 563–C. FA29:2099.

712. Leaf, A.L.

1957. Diagnosis of deficiencies of available potassium calcium, and magnesium in forested soils. Diss. Abstr. 17(9):1859. FA19:1472.

713. Leaf, A.L., and D.H. Bickelhaupt.

1975. Possible mutual prediction between black cherry and sugar maple foliar analysis data. Proc. Soil Sci. Soc. Am. 39(5):983–985. BA61:31319.

714. Leaf, A.L., and S.A. Wilde.

1957. Decomposition products of sloughed root tissues promote release of available nutrients. Univ. Wis. For. Res. Note 40. FA19:2634. (see next entry)

715. Leaf, A.L., and S.A. Wilde.

1959. Decomposition products of sloughed root tissues promote release of available nutrients. Univ. Wis. For. Res. Note 49. FA21:1330.

716. Leaf, A.L., R.E. Leonard, and J.V. Berglund.

1971. Root distribution of a plantation-grown red pine in an outwash soil. Ecology 52(1):153–158. BA52:77849.

717. Leaphart, C.D. 1960. A root stain disease of eas

1960. A root stain disease of eastern white pine. Plant Dis. Rep. 44(9):704–706. BA36:42591.

718. Leaphart, C.D.

1963. Armillaria root rot. USDA For. Serv. For. Pest Leafl. 78.

LeBarron, R.K.
 1944. Influence of controllable environmental conditions on regeneration of jack pine and black spruce. J. Agric. Res. 68(3):97–119.

720. LeBarron, R.K.
1945. Adjustment of black spruce root systems to increasing depth of peat. Ecology 26(3):309–311. BA19:18067.

721. Leclercq, W.L.
1960. Dennenmoorder en *Prunus serotina*. [Fomes annosus and P. serotina.] Ned. Bosb. Tijdschr. 32(2):74–75.

722. Ledig, F.T., and T.O. Perry. 1966. Physiological genetics of the shoot-root ratio. Proc. Soc. Am. For. 1965, p. 39–43. FA28:342.

Ledig, F.T., F.H. Bormann, and K.F. Wenger.
 1970. The distribution of dry matter growth between shoot and roots in loblolly pine. Bot. Gaz. 131(4):349–359. BA52:89184.

724. Ledig, F.T., A.P. Drew, and J.G. Clark.
1976. Maintenance and constructive respiration, photosynthesis, and net assimilation rate in seedlings of pitch pine (*Pinus rigida* Mill.). Ann. Bot. 40(166):289–300. BA62:63012.

725. Lee, C.I., B.C. Moser, and C.E. Hess. 1974. Root regeneration of transplanted pin and scarlet oak. New Horiz. Hort. Res. Inst., Washington, D.C., p. 10–13.

726. Leiser, A.T. 1968. A mucilaginous root sheath in Ericaceae. Am. J. Bot. 55(3):391–398.

727. Lemon, P.C. 1945. Wood as a substratum for perennial plants in the Southeast. Am. Midl. Nat. 34(3):744-749. BA20:19423.

728. Levisohn, I. 1952. Forking in pine roots. Nature 169(4304):715. FA13:3681.

729. Levisohn, I.
1954. Aberrant root infections of pine and spruce seedlings. New Phytol. 53:284–290.

730. Lewis, F.J., E.S. Dowding, and E.H. Moss. 1928. The vegetation of Alberta. II. The swamp, moor, and bog forest vegetation of central Alberta. J. Ecol. 16(1):19–70.

731. Leyton, L., and L.Z. Rousseau.

1958. Root growth of tree seedlings in relation to aeration. *In* The physiology of forest trees. A symposium. K.V. Thimann, ed. Maria Moors Cabot Found. for Bot. Res. Int. Symp. For. Tree Physiol., Harvard Univ. Cambridge, Mass. Ronald Press Co., New York. p. 467–475. BA33:19470.

732. Lightle, P.C. 1960. Fomes annosus root rot of loblolly pine. Plant Dis. Rep. 44(6):423. BA36:8859.

733. Liming, F.G., and J.P. Johnston. 1944. Reproduction in oak-hickory forest stands of the Missouri Ozarks. J. For. 42(3)175–180.

734. Limstrom, G.A., R.F. Finn, and G.H. Deitschman. 1955. Planting stock grades for yellow poplar. J. For. 53(1):28–32. BA30:11328.

735. Ling-lee, M., G.A. Chilvers, and A.E. Ashford.
1975. Polyphosphate granules in three different kinds of tree mycorrhiza. New Phytol. 75(3):551–554. FA37:3603.

736. Lister, G.R.
1968. Observations on the growth and physiology of *Pinus strobus*L. seedlings grown under various conditions of soil moisture and nitrogen and phosphorus nutrition. (Parts I and II). Diss. Abstr. 29B(4):1277. FA30:3677.

737. Lister, G.R., V. Slankis, G. Krotkov, and C.D. Nelson.
1968. The growth and physiology of *Pinus strobus* L. seedlings as affected by various nutritional levels of nitrogen and phosphorus.
Ann. Bot. (N.S.) 32(125):33–43. BA49:112933.

738. Little, S. 1973. Survival, growth of loblolly, pitch, shortleaf pines established by different methods in New Jersey. Tree Plant. Notes 24(4):1–5. FA35:6848.

739. Little, S., and H.A. Somes. 1964. Root systems of direct-seeded and variously planted loblolly, shortleaf, and pitch pines. USDA For. Serv. Res. Pap. NE-26. BA46:63682.

740. Loach, K., and C.H.A. Little.
1973. Production, storage and use of photosynthate during shoot elongation in balsam fir (*Abies balsamea*). Can. J. Bot. 51(6):1161–1168. FA35:2165.

741. Lodewich, J.E.1928. Seasonal activity of the cambium in some northeastern trees.Bull. N.Y. State Coll. For., Tech. Publ. 23.

742. Loftus, N.S. 1971. Yellow-poplar root development on Hartsells subsoils. USDA For. Serv. Res. Note SO–131. FA33:4038.

743. Logan, K.T. 1965. Growth of tree seedlings as affected by light intensity. I. White birch, yellow birch, sugar maple and silver maple. Can. Dep. For. Publ. 1121.

744. Logan, K.T. 1966. Growth of tree seedlings as affected by light intensity. II. Red pine, white pine, jack pine and eastern larch. Can. Dep. For. Publ. 1160.

745. Logan, K.T.
1966. Growth of tree seedlings as affected by light intensity. III.
Basswood and white elm. Can. Dep. For. Publ. 1176.

746. Logan, K.T.
1969. Growth of tree seedlings as affected by light intensity. IV.
Black spruce, white spruce, balsam fir, and eastern white cedar.
Can. Dep. Environ. For. Serv. Publ. 1256.

747. Logan, K.T.
 1973. Growth of tree seedlings as affected by light intensity. V.
 White ash, beech, eastern hemlock, and general conclusions. Can.
 Dep. Environ. For. Serv. Publ. 1323.

748. Lohman, M.L. 1926. Occurrence of mycorrhiza in Iowa forest plants. Ia. Univ. Stud. in Nat. Hist. 11(10):33–58.

749. Lorio, P.L., Jr., and J.D. Hodges.
1968. Microsite effects on oleoresin exudation pressure of large loblolly pines. Ecology 49(6):1207–1210. FA30:5201.

750. Lorio, P.L., Jr., and J.D. Hodges.
1971. Microrelief, soil water regime, and loblolly pine growth on a wet, mounded site. Proc. Soil Sci. Soc. Am. 35(5):795–800.
BA53:31362.

751. Lorio, P.L., Jr., V.K. Howe, and C.N. Martin.
1972. Loblolly pine rooting varies with microrelief on wet sites.
Ecology 53(6):1134-1140. BA55:59708.

752. Lortie, M.
1966. Observations sur l'enracinement du bouleau à papier à la suite d'études sur le dépérissement. [Observations on the rooting habit of white birch following studies on its decline.] Nat. Can. 93(3):199–206. [Engl. summ.] BA48:9655.

753. Love, H.S., Jr. 1972. The detection and nature of functional underground connections between members of certain woody species of the Oklahoma Cross Timbers. Diss. Abstr. Int. 33B(2):600. FA35:6728. 754. Luckoff, H.A.

1955. Two hitherto unrecorded fungal diseases attacking pines and eucalyptus in South Africa. J. S. Afr. For. Assoc. 26, p. 47–61. FA18:578.

755. Luncz, G.

1931. Recent research work on the root systems of forest trees. Int. Rev. Agric, 22:239T-243T.

756. Lunt, H.A.

1934. Distribution of soil moisture under isolated forest trees. J. Agric. Res. 49(8):695–703. BA10:1645.

757. Lutz, H.J.

1930. Effect of cattle grazing on vegetation of a virgin forest in northwestern Pennsylvania. J. Agric. Res. 41(7):561–570.

758. Lutz, H.J.

1939. Layering in eastern white pine. Bot. Gaz. 101:505-507. FA2:32.

759. Lutz, H.J.

1956. Ecological effects of forest fires in the interior of Alaska. USDA Tech. Bull. 1133.

760. Lutz, H.J., J.B. Ely, Jr., and S. Little, Jr.

1937. The influence of soil profile horizons on root distribution of white pine (*Pinus strobus* L.). Yale Univ. Sch. For. Bull. 44.

761. Lyarskaya, R.P.

1959. Nekotorye osobennosti estestvennogo vegetativnogo razmnozheniya uksusnogo dereva (*Rhus typhina* L.) kornevymi otpryskami. [Certain specific features of the natural vegetative reproduction of *R. typhina* by means of root suckers.] Nauch Dok. Vyssh. Shk. Biol. Nauk. 1:117–123. BA40:16411.

762. Lyford, W.H., and B.F. Wilson.

1964. Development of the root system of *Acer rubrum* L. Harvard For. Pap. 10. BA47:9504.

763. Lyford, W.H., and B.F. Wilson.

1966. Controlled growth of forest tree roots: Technique and application. Harvard For. Pap. 16. BA48:71969.

764. Lyle. E.S., Jr., and F. Adams.

1971. Effect of available soil calcium on taproot elongation of loblolly pine (*Pinus taeda* L.) seedlings. Proc. Soil Sci. Soc. Am. 35(5):800–805. BA53:31363.

765. Lyon, C.J.

1949. Secondary growth of white pine in bog and upland. Ecology 30(4):549–552. BA24:3286.

766. Lyr, H., and G. Hoffmann.

1967. Growth rates and growth periodicity of tree roots. Int. Rev. For. Res. 2:181–236. FA29:3567.

767. MacAloney, H.J.

1944. Relation of root condition, weather, and insects to the management of jack pine. J. For. 42(2):124–129.

768. McAlpine, R.G.

1959. Flooding kills yellow-poplar. For. Farmer 19(3):9, 13–14.

769. McAlpine, R.G.

1961. Yellow-poplar seedlings intolerant to flooding. J. For. 59(8):566–568.

770. McAlpine, R.G., and L.W.R. Jackson.

1959. Effect of age on rooting of loblolly pine air-layers. J. For. 57(8):565–566. BA34:14315.

771. McArdle, R.E.

1932. The relation of mycorrhizae to conifer seedlings. J. Agric. Res. 44(4):287–316.

772. McClurkin, I.T., and D.C. McClurkin.

1967. Cytochemical demonstration of a sodium-activated and a potassium-activated adenosine triphosphatase in loblolly pine seedling root tips. Plant Physiol. 42(8):1103–1110. FA29:1986.

773. McColl, J.G.

1973. Soil moisture influence on growth, transpiration, and nutrient uptake of pine seedlings. For. Sci. 19(4):281–288. FA35:5813.

774. McComb, A.L.

1938. The relation between mycorrhizae and the development and nutrient absorption of pine seedlings in a prairie nursery. J. For. 36(11):1148–1153. FA1:16.

775. McComb, A.L.

1943. Mycorrhizae and phosphorus nutrition of pine seedlings in a prairie soil nursery. Ia. Agric. Exp. Stn. Res. Bull. 314, p. 582–612. BA17:19568.

776. McComb, A.L., and J.E. Griffith.

1946. Growth stimulation and phosphorus absorption of mycorrhizal and nonmycorrhizal northern white pine and Douglas fir seedlings in relation to fertilizer treatment. Plant Physiol. 21:11–17.

777. MacConnell, J.T.

1959. The oxygen factor in the development and function of the root nodules of alder. Ann. Bot. (N.S.) 23(90):261–268. BA35:11174.

778. McDermott, R.E.

1954. Effects of saturated soil on seedling growth of some bottom-land hardwood species. Ecology 35(1):36–41.

779. McDougall, W.B.

1914. On the mycorrhizas of forest trees. Am. J. Bot. 1(2):51-74.

780. McDougall, W.B.

1916. The growth of forest tree roots. Am. J. Bot. 3(7):384–392.

781. McDougall, W.B.

1921a. A preliminary key to some forest tree roots. Trans. Ill. Acad. Sci. 14:87–91.

782. McDougall, W.B.

1921b. Thick-walled root hairs of *Gleditsia* and related genera. Am. J. Bot. 8(3):171–175.

783. McDougall, W.B.

1922. Mycorrhizas of coniferous trees. J. For. 20(3):255-260.

784. McEwen, J.K.

1966. An effect of sphagnum on the growth of black spruce. For. Chron. 42(2):175–183. FA28:275.

785. McGauley, B., and M. Hubbes.

1976. Relative susceptibility of selected pure and hybrid pines to *Fomes annosus* (Fr.) Cke. Eur. J. For. Pathol. 6(3):167–176. FA38:1437.

786. McKenzie, J.S., C.J. Weiser, and P.H. Li.

1974. Changes in water relations of *Cornus stolonifera* during cold acclimation. J. Am. Soc. Hort. Sci. 99(3):223–228. FA36:3214.

787. McLaughlin, S.B., N.T. Edwards, and J.J. Beauchamp.

1977. Spatial and temporal patterns in transport and respiratory allocation of [14C] sucrose by white oak (*Quercus alba*) roots. Can. J. Bot. 55(23):2971–2980.

788. McLintock, T.F.

1954. Factors affecting wind damage in selectively cut stands of spruce-fir in Maine and northern New Hampshire. USDA For. Serv. Stn. Pap. 70. Northeast. For. Exp. Stn.

789. McQuilkin, W.E.

1935. Root development of pitch pine, with some comparative observations on shortleaf pine. J. Agric. Res. 51(11):983–1016. BA10:6818.

790. McVean, D.N.

1956. Ecology of *Alnus glutinosa* (L.) Gaertn. III. Seedling establishment. J. Ecol. 44(1):195–218. FA17:3703.

791. McVean, D.N.

1956. Ecology of *Alnus glutinosa* (L.) Gaertn. IV. Root system. J. Ecol. 44(1):219–225. FA17:3704.

792. Mai, W.F., H.W. Crittenden, and W.R. Jenkins.

1960. Distribution of stylet-bearing nematodes in the northeastern United States. N.J. Agric. Exp. Stn. Bull. 795.

793. Maini, J.S.

1968. Silvics and ecology of *Populus* in Canada. *In* Growth and utilization of poplars in Canada. J.S. Maini and J.H. Cayford, eds., Can. Dep. For. Publ. 1205, p. 20–69.

794. Maini, J.S., and K.W. Horton.

1966. Vegetative propagation of *Populus* spp. I. Influence of temperature on formation and initial growth of aspen suckers. Can. J. Bot. 44(9):1183–1189. BA48:4151.

795. Maisenhelder, L.C.

1957. Propagation of some Delta hardwoods by rooting. 4th South. Cong. For. Tree Improv. Proc., Athens, Ga., [8–9 January 1957], p. 55–58.

796. Maki, T.E., and H. Marshall.

1945. Effects of soaking with indolebutyric acid on root development and survival of tree seedlings. Bot. Gaz. 107(2):268–276. BA20:9036.

797. Malek, R.B., and J.D. Smolik.

1975. Effect of *Xiphinema americanum* on growth of shelterbelt trees. Plant Dis. Rep. 59(2):144–148. FA36:5808.

798. Mann, W.F., Jr.

1950. Competition for light, water, and nutrients. USDA For. Serv. South. For. Notes 69. South. For. Exp. Stn., p. 3.

799. Manolova, Y., and I. Palashev.

1974. [The water-retaining capacity of the roots of several forest tree species.] Gorskostop. Nauka 11(2): 14–19. [In Bulg. with Russ. and Engl. summ.] FA36:129.

800. Marshall, R.P.

1948. A white root rot which merits study as possible cause of white pine blight. Scientific Tree Topics 1(9): 66–68. Bartlett Tree Res. Lab., Stamford, Conn. FA10:2242.

801. Martin, J.L.

1965. Living stumps aid insect control. Can. For. Ind. 85(7):40-43.

802. Marx, D.H.

1966. The role of ectotrophic mycorrhizal fungi in the resistance of pine roots to infection by *Phytophthora cinnamomi* Rands. Diss. Abstr. 27B(6):1692. FA28:5873.

803. Marx. D.H.

1969a. The influence of ectotrophic mycorrhizal fungi in the resistance of pine roots to pathogenic infections. I. Antagonism of mycorrhizal fungi to root pathogenic fungi and soil bacteria. Phytopathology 59(2):153–163. FA30:5389.

804. Marx., D.H

1969b. The influence of ectotrophic mycorrhizal fungi on the resistance of pine roots to pathogenic infections. II. Production, identification, and biological activity of antibiotics produced by *Leucopaxillus cerealis* var. *piceina*. Phytopathology 59(4):411–417.

805. Marx, D.H.

1970. The influence of ectotrophic mycorrhizal fungi on the resistance of pine roots to pathogenic infections. V. Resistance of mycorrhizae to infection by vegetative mycelium of *Phytophthora cinnamomi*. Phytopathology 60(10):1472–1473.

806. Marx, D.H.

1973. Growth of ectomycorrhizal and nonmycorrhizal shortleaf pine seedlings in soil with *Phytophthora cinnamomi*. Phytopathology 63(1):18–23.

807. Marx, D.H.

1975. Mycorrhizae and establishment of trees on strip-mined land. Ohio J. Sci. 75(6):288–297. FA37:5550.

808. Marx, D.H.

1976. Synthesis of ectomycorrhizae on loblolly pine seedlings with basidiospores of *Pisolithus tinctorius*. For. Sci. 22(1):13–20.

809. Marx, D.H.

1977a. Manipulation of selected mycorrhizal fungi to increase forest biomass. TAPPI, For. Biol./Wood Chem. Conf., Madison, Wis.

810. Marx, D.H.

1977b. Tree host range and world distribution of the ectomycorrhizal fungus *Pisolithus tinctorius*. Can. J. Microbiol. 23(3):217–223.

811. Marx, D.H., and W.C. Bryan.

1969. Effect of soil bacteria on the mode of infection of pine roots by *Phytophthora cinnamomi*. Phytopathology 59(5):614–619. FA31:895.

812. Marx, D.H., and W.C. Bryan.

1970. Pure culture synthesis of ectomycorrhizae by *Thelephora* terrestris and *Pisolithus tinctorius* on different conifer hosts. Can. J. Bot. 48:639–643. FA32:332.

813. Marx, D.H., and W.C. Bryan.

1971. Influence of ectomycorrhizae on survival and growth of aseptic seedlings of loblolly pine at high temperature. For. Sci. 17(1):37–41. BA52:83967.

814. Marx, D.H., and W.C. Bryan.

1975a. Growth and ectomycorrhizal development of loblolly pine seedlings in fumigated soil infested with the fungal symbiont *Pisolithus tinctorius*. For. Sci. 21(3):245–254. BA61:25436.

815. Marx, D.H., and W.C. Bryan.

1975b. The significance of mycorrhizae to forest trees. *In* Forest soils and forest land management. B. Bernier and C.H. Winget, eds., p. 107–117. 4th North Am. For. Soils Conf. Proc., Laval Univ., 1973. Press l'Univ. Laval, Quebéc. FA37:2174.

816. Marx, D.H., and C.B. Davey.

1967. Ectotrophic mycorrhizae as deterrents to pathogenic root infections. Nature 213(5081):1139. BA49:31620.

817. Marx, D.H., and C.B. Davey.

1969a. The influence of ectotrophic mycorrhizal fungi on the resistance of pine roots to pathogenic infection. III. Resistance of aseptically formed mycorrhizae to infection by *Phytophthora cinnamomi*. Phytopathology 59(5):549–558. BA51:98534.

818. Marx, D.H., and C.B. Davey.

1969b. The influence of ectotrophic mycorrhizal fungi on the resistance of pine roots to pathogenic infections. IV. Resistance of naturally occurring mycorrhizae to infections by *Phytophthora cinnamomi*. Phytopathology 59(5):559–565. BA51:98533.

819. Marx, D.H., W.C. Bryan, and C.B. Davey.

1970a. Influence of temperature on aseptic systhesis of ectomycorrhizae by *Thelephora terrestris* and *Pisolithus tinctorius* on loblolly pine. For. Sci. 16(4):424–431.

820. Marx, D.H., W.C. Bryan, and L.F. Grand.

1970b. Colonization, isolation, and cultural descriptions of *Thelephora terrestris* and other ectomycorrhizal fungi of shortleaf pine seedlings grown in fumigated soil. Can. J. Bot. 48(2):207–211.

821. Marx, D.H., W.C. Bryan, and C.E. Cordell.

1976. Growth and ectomycorrhizal development of pine seedlings in nursery soils infested with the fungal symbiont *Pisolithus tinctorius*. For. Sci. 22(1):91–100.

822. Marx, D.H., W.C. Bryan, and C.E. Cordell.

1977. Survival and growth of pine seedlings with *Pisolithus* ectomy-corrhizae after two years on reforestation sites in North Carolina and Florida. For. Sci. 23(3):363–373.

823. Marx, D.H., A.B. Hatch, and J.F. Mendicino.

1977. High soil fertility decreases sucrose content and susceptibility of loblolly pine roots to ectomycorrhizal infection by *Pisolithus tinctorius*. Can. J. Bot. 55(12):1569–1574. BA64:50025.

824. Massey, A.B.

1925. Antagonism of the walnuts (*Juglans nigra* L. and *J. cinerea* L.) in certain plant associations. Phytopathology 15:773–784.

825. Mattoon, W.R.

1915. The southern cypress. USDA Bull. 272.

826. Medve, R.J.

1969. Development and morphology of the beaded rootlets, mycorrhizae, and associated root fan structures of red maple (*Acer rubrum* L.). Diss. Abstr. Int. 30B(1):82–83. FA32:2208.

827. Medve, R.J.

1970. Beaded rootlets and associated root fan structures of *Acer rubrum* L. Pa. Acad. Sci. 44:122–126.

328. Medve, R.J

1970. Influence of genotype and soil on root fan structures of *Acer rubrum*. Can. J. Bot. 48(1):147–152. BA51:63045.

829. Medve, R.J.

1970. The relationship of beaded rootlets and mycorrhizae of red maple (*Acer rubrum* L.). Am. Midl. Nat. 83(2):631-634. FA32:2207.

830. Medve, R.J.

1971. Beaded rootlets in Acer. Pa. Acad. Sci. 45:98-100.

831. Medve, R.J., F.M. Hoffman, and T.W. Gaither.
1977. The effects of mycorrhizal-forming amendments on the revegetation of bituminous stripmine spoils. Bull. Torrey Bot. Club 104(3):218–225.

832. Melin, E., H. Nilsson, and E. Hacskaylo.
1958. Translocation of cations to seedlings of *Pinus virginiana* through mycorrhizal mycelium. Bot. Gaz. 119 (4):243–246.

833. Menge, J.A., L.F. Grand, and L.W. Haines.
1977. The effect of fertilization on growth and mycorrhizae numbers in 11-year-old loblolly pine plantations. For. Sci. 23(1):37–44.

834. Mergen, F.
1954. Mechanical aspects of wind-breakage and windfirmness. J.
For. 52(2):119–125.

835. Mergen, F., and J. Worrall.
1965. Effect of environment and seed source on mineral content of jack pine seedlings. For. Sci. 11(4):393–400. FA27:5391.

836. Merrit, C.
1959. Studies on the root growth of red pine (*Pinus resinosa* Ait.).
PhD. thesis, Univ. Mich., Ann Arbor.

837. Merrit, C.
1968. Effect of environment and heredity on the root-growth pattern of red pine. Ecology 49(1):34–40. BA49:86074.

838. Merz, R.W., and S.G. Boyce. 1956. Age of oak 'seedlings'. J. For. 54(11):774-775. BA31:25786.

839. Meserve, A.W.
1937. Effect of changing grades around trees and suggested treatment. Natl. Shade Tree Conf. Proc. 13:36–51.

840. Meyer-Brenken.
1954. Wurzelausbildung der roteiche. [Root development of *Quercus borealis*.] Forest-u. Holzw. 9(22):464–465. FA17:1339.

841. Migunova, E.S.
1976. [Root systems of woody species on saline soils in the southern
Ukrainian SSR.] Lesovedenie 6:27–36. [In Russ, with Russ. and
Engl. summ.] BA64:13694.

842. Mikola, P.
1952. Effect of forest humus on parasitic fungi causing damping-off disease of coniferous seedlings. Phytopathology 42(4):202–203. FA13:3785.

843. Mikola, P.
1953. An experiment on the invasion of mycorrhizal fungi into prairie soil. Karstenia 2, p. 33–34. FA17:227.

844. Miller, A.E., and M. Reines.
1974. Survival and water relations in loblolly pine seedlings after root immersion in alginate solution. For. Sci. 20(2):192–194. FA36:2034.

845. Miller, L., and F.W. Woods. 1965. Root-grafting in loblolly pine. Bot. Gaz. 126(4): 252–255. BA47:54141.

846. Miller, O.K.
1959. Red pine mortality from root rot (*Fomes annosus* (Fries)
Cooke). Fox For. Note 75, Caroline A. Fox Res. and Demonstration
For., N.H. FA21:3356.

847. Miller, T.
1964. Host factors influencing infection and growth of *Fomes annosus* in roots of *Pinus taeda*. (Abstr.) Phytopathology 54(8):901.
BA46:45190.

848. Miller, T., and A. Kelman.
1966. Growth of *Fomes annosus* in roots of suppressed and dominant loblolly pines. For. Sci. 12(2):225–233. BA47:109517.

849. Mirov, N.T. 1967. The genus *Pinus*. Univ. Calif. Press, Berkeley.

850. Mitchell, H.L.
1934. Pot culture tests of forest soil fertility, with observations on the effect of varied solar radiation and nutrient supply on the growth and nitrogen content of Scots and white pine seedlings. Black Rock For. Bull. 5. BA9:16639.

851. Mitchell, H.L.
1939. The growth and nutrition of white pine (*Pinus strobus* L.) seedlings in cultures with varying nitrogen, phosphorus, potassium, and calcium, with observations on the relation of seed weight to seedling yield. Black Rock For. Bull. 9.

852. Mitchell, H.L., and R.F. Finn.
1937. The relative feeding power of oaks and maples for soil phosphorus. Black Rock For. Pap. 1(2):5–9.

853. Mitchell, H.L., and R.O. Rosendahl.
1939. The relationships between cumulative solar radiation and the dry weight increase of nursery-grown white pine and red pine seedlings. Black Rock For. Pap. 1(13):88–93. FA1:75.

854. Mogensen, H.L.
1967. A contribution to the developmental anatomy of the root of *Quercus* L. Ia. State J. Sci. 41(4):413–423. BA48:87618.

855. Monk, C.D.
1966. Root-shoot dry weights in loblolly pine. Bot. Gaz.
127(4):246-248. BA48:56539.

856. Monk, C.D.
1967. The root-shoot weight relationship of four widely separated seedling populations of red maple. Bull. Torrey Bot. Club 94(3):197–199. BA48:109879.

Monk, R., and H.B. Peterson.1962. Tolerance of some trees and shrubs to saline conditions. Proc. Am. Soc. Hort. Sci. 81:556–561.

858. Moore, A.W.
1964. Note on non-leguminous nitrogen-fixing plants in Alberta.
Can. J. Bot. 42(7):952–955. FA26:1892.

859. Moore, B.
1922a. Humus and root systems in certain northeastern forests in relation to reproduction and competition. J. For. 20(3):233–254.

 Moore, B.
 1922b. Influence of certain soil factors on the growth of tree seedlings and wheat. Ecology 3(1):65–83.

Moreland, D.E.
 1950. A study of the translocation of radioactive phosphorus in loblolly pine (*Pinus taeda* L.). J. Elisha Mitchell Sci. Soc. 66(2):175–181. FA13:938.

862. Moriondo, F. 1959. Un singolare attaccodi *Armillaria* sp. su *Robinia pseudoaca-cia* L. [A peculiar attack by *A. mellea* on *R. pseudoacacia*.] Ital. For. Mont. 14(5):207–209. [Fr. summ.] FA21:1990.

Morris, R.C., T.H. Filer, J.D. Solomon, F.I. McCracken, N.A. Overgaard, and M.J. Weiss.
 1975. Insects and diseases of cottonwood. USDA For. Serv. Gen. Tech. Rep., SO–8. FA37:6241.

864. Morrison, I.K.
1970. The absorption of nutrients by roots of coniferous seedlings in relation to root characteristics and soil conditions. Diss. Abstr. Int. 31B(2):460. FA33:330.

865. Morrison, I.K.
1974. Dry-matter and element content of roots of several natural stands of *Pinus banksiana* Lamb. in northern Ontario. Can. J. For. Res. 4(1):61–64. BA58:13804.

866. Morrison, R.H., and D.W. French.
1969. Taxonomy of *Cylindrocladium floridanum* and *C. scoparium*.
Mycologia 61(5):957–966.

867. Morrow, R.R.
1950. Periodicity and growth of sugar maple surface layer roots. J.
For. 48(12):875–881. BA25:29061.

868. Mosher, D.G., and L.F. Wilson.1977. Scotch pine deterioration in Michigan caused by pine root

weevil complex. Great Lakes Entomol. 10(4):169-172. BA65:32913.

869. Moss, E.H.

1938. Longevity of seed and establishment of seedlings in species of *Populus*. Bot. Gaz. 99(3):529–542.

870. Motto, C.K.

1964. Cation exchange capacities of tree roots in relation to their mineral compositions. M.S. thesis, Univ. Ill., Urbana.

Moxley, L., and H. Davidson.
 1973. Salt tolerance of various woody and herbaceous plants. Mich.
 State Univ. Hort. Rep. 23.

872. Mueller, O.P., and M.G. Cline. 1959. Effects of mechanical soil barriers and soil wetness on rooting of trees and soil-mixing by blow-down in central New York. Soil Sci. 88(2):107–111. BA34:11332.

873. Mueller-Dombois, D.
1964. Effect of depth to water table on height growth of tree seed-lings in a greenhouse. For. Sci. 10(3):306–316. FA26:1873.

874. Mullin, R.E. 1963. Growth of white spruce in the nursery. For. Sci. 9(1):68–72. FA24:4881.

Mullin, R.E.
 1966. Root pruning of nursery stock. For. Chron. 42(3):256–264.
 BA48:15147.

876. Mullin, R.E.
 1967. Root exposure of white spruce nursery stock. For. Chron.
 43(2):155–160. BA48:113886.

877. Mullin, R.E.1978. Root exposure, root dipping and extended spring planting of white pine seedlings. For. Chron. 54(2):84–87.

878. Musselman, R.C., D.T. Lester, and M.S. Adams. 1975. Localized ecotypes of *Thuja occidentalis* L. in Wisconsin. Ecology 56(3):647–655. BA60:42576.

879. Myren, D.T., and R.F. Patton.
1971. Establishment and spread of *Polyporus tomentosus* in pine and spruce plantations in Wisconsin. Can. J. Bot. 49(6):1033–1040. FA33:945.

880. Myren, D.T., H.L. Gross, and E.B. Dorworth. 1975. *Cylindrocladium floridanum* in an Ontario forest nursery. Bi-mon. Res. Notes 31(5):34. FA37:4585.

881. Nair, V.M.G., and J.E. Kuntz.

1963. Ceratocystis fagacearum in roots of infected bur oaks and the recurrence and spread of oak wilt. (Abstr.) Phytopathology 53(8):884. BA45:44124.

882. Neary, D., M. Day, and G. Schneider. 1972. Density-growth relationships in a nine-year-old red pine plantation. Mich. Acad. 5(2):219–232. BA56:14103.

Neely, D., and E.B. Himelick.
 1963. Root graft transmission of Dutch elm disease in municipalities. Plant Dis. Rep. 47(2):83–85. BA45:79790.

Nelson, T.C.1951. A reproduction study of northern white-cedar. Mich. Dep. Conserv., Game Div.

885. Nemeth, J.C.
1973. Dry matter production in young loblolly (*Pinus taeda* L.) and slash pine (*Pinus elliottii* Engelm.) plantations. Ecol. Monogr. 43(1):21–41. FA34:6852.

886. Nesbitt, W.A.
1942. Aerial roots from old tree wound. Plant Physiol. 17(4):689–690. BA17:12276.

887. Nicholas, A.K., and R.J. Hutnik.
1971. Ectomycorrhizal establishment and seedling response on variously treated deep-mine coal refuse. Pa. State Univ., Coll. Earth and Mineral Sci. Spec. Res. Rep. SR-89. FA36:2691.

888. Norton, J.A., J.P. Walter, Jr., and J.B. Storey.
1970. The effect of herbicides on lateral roots and nut quality of pecans. Weed Sci. 18(4):520–522. BA51:133010.

Nova Scotia Department of Lands and Forests.
 1946. Root washings of yellow birch. Rep. Dep. Lands and For.
 1945, p. 45–47, 52. FA8:1459.

890. O'Leary, J.W.
1964. Exudation from root systems of woody plants. Diss. Abstr.
25(5):2727. BA46:63389.

891. O'Leary, J.W.
1965. Root-pressure exudation in woody plants. Bot. Gaz.
126(2):108-115. FA27:1708.

892. O'Leary, J.W., and P.J. Kramer.
 1964. Root pressure in conifers. Science 145(3629):284–285.
 BA46:12782.

893. O'Neil, W.J. 1928. More about living stumps. J. For. 26(2):244–245. BA3:9085.

 Oosting, H.J., and P.J. Kramer.
 1946. Water and light in relation to pine reproduction. Ecology 27(1):47-53.

895. Opalicki, K.
1969. [Occurrence of root nematodes in forest nurseries.] Sumar.
List 93(3/4):126–130. [In Serb. with Engl. summ.] FA31:4974.

896. Orlič, S., M. Harapin, M. Halambek, and B. Mayer.
1973. Sušeje američkog borovca (*Pinus strobus* L.) u kulturama.
[Die-back of eastern white pine (*P. strobus* L.) in plantations.]
Sumar. List 97(9/10):375–385. [Engl. summ.] FA35:5400.

897. Ortmann, C.
1961. Vorläufige Untersuchungsergebnisse zur Frage der Selektionstypen für die Frühdiagnose von *Salix alba-*Populationen. [Preliminary investigations on the question of the type selection for the early diagnoses of *S. alba* populations.] Silvae Genet. 10(2):43–48. [Engl. summ.] FA23:133.

898. Oskamp, J.
 1932. The rooting habit of deciduous fruits on different soils. Proc.
 Am. Soc. Hort. Sci. 29:213–219. BA9:5636.

899. Otrosina, W.J., and D.H. Marx.
1975. Populations of *Phytophthora cinnamomi* and *Pythium* spp. under shortleaf and loblolly pines in littleleaf disease sites. Phytopathology 65(11):1224–1229. FA37:3398.

900. Ouellette, G.B., and G. Bard.
 1961. Notes on the occurrence of *Armillaria* root rot. Bi-mon. Progr.
 Rep., For. Entomol. Pathol. Branch, Dep. For. Can. 17(6):1-2.

901. Ouellette, G.B., G. Bard, and R. Cauchon.
1971. Self-strangulation of roots: Points of entry of root-rot fungi in
the Grand'Mère white spruce plantations. Phytoprotection
52(3):119-124.

902. Ovington, J.D.
1957. Dry-matter production by *Pinus silvestris* L. Ann. Bot. (N.S.)
21:287.

903. Page, F.S. 1927. Living stumps. J. For. 25(6):687–690. BA3:2766.

904. Paine, L.A.
1956. Développement des racines de l'épinette blanche en sol sablonneux. [Development of the roots of white spruce in sandy soil.] (Abstr.) Ann. ACFAS (Assoc. Con. Fr. Av. Sci.) 22:54. FA19:2813.

905. Park, J.Y. 1970. A change in color of aging mycorrhizal roots of *Tilia america-na* formed by *Cenococcum graniforme*. Can. J. Bot. 48(7):1339–1341. BA52:1933.

Parker, J.
 1950. The effects of flooding on the transpiration and survival of some southeastern forest tree species. Plant Physiol. 25(3):453–460.

907. Parker, J.

1959. Seasonal variations in sugars of conifers with some observations on cold resistance. For. Sci. 5(1):56–63. BA33:35079.

908. Parker, J.

1968. Drought resistance of roots of white ash, sugar maple, and red oak. USDA For. Serv. Res. Pap. NE-95.

909. Parker, J.

1970. Effects of defoliation and drought on root food reserves in sugar maple seedlings. USDA For. Serv. Res. Pap. NE-169.

910. Parker, J.

1974. Effects of defoliation, girdling, and severing of sugar maple trees on root starch and sugar levels. USDA For. Serv. Res. Pap. NE-306.

911. Parker, J.

1975. The nature of starch storage fiber cells in *Acer saccharum*. Adv. Front. of Plant Sci. 30:137–150. FA37:74.

912. Parker, J., and D.R. Houston.

1971. Effects of repeated defoliation on root and root collar extractives of sugar maple trees. For. Sci. 17(1):91–95. BA52:83977.

913. Parker, J., and R.L. Patton.

1975. Effects of drought and defoliation on some metabolites in roots of black oak seedlings. Can. J. For. Res. 5(3):457–463. BA61:16540.

914. Patton, R.F., and R.V. Bravo.

1967. Armillaria root rot: Armillaria mellea (Vahl. ex Fr.) Kummer. In Important forest insects and diseases of mutual concern to Canada, the United States, and Mexico. A. G. Davidson, and R.M. Prentice, eds. Can. For. Serv. Publ. 1180, p. 37–38.

915. Patton, R.F., and R.G. Krebill.

1960. Deterioration of immature jack and red pine plantations in Wisconsin. Univ. Wis. For. Res. Notes 64. FA22:3425.

916. Patton, R.F., and R.G. Krebill.

1960. Deterioration of immature jack and red pine plantations in Wisconsin. (Abstr.) Phytopathology 50(9):642–643. FA22:2175.

917. Patton, R.F., and D.T. Myren.

1970. Root rot induced by *Polyporus tomemtosus* in pine and spruce plantations in Wisconsin. *In* Root diseases and soil-borne pathogens. T.A. Toussoun, R.V. Bega, and P.E. Nelson, eds. Univ. Calif. Press, Berkeley, p. 167–170.

918. Patton, R.F., and A.J. Riker.

1954. Top growth and root development of rooted white pine cuttings. J. For. 52(9):675–677.

919. Patton, R.F., and A.J. Riker.

1958. Rooting cuttings of white pine. For. Sci. 4(2):116-127.

920. Patton, R.F., and A.J. Riker.

1959. Artificial inoculations of pine and spruce trees with *Armillaria mellea*. Phytopathology 49(9):615–622. FA21:1989.

921. Pawsey, R.G., and M.A. Rahman.

1976. Chemical control of infection by honey fungus, *Armillaria mellea*: A review. J. Aboric. 2(9):161–169.

922. Pearsall, W.H.

1943. The mycorrhizal relations of *Pinus*. J. Ecol. 31(1):41–44. BA17:23140.

923. Pellett, N.E.

(n. d. Circa 1975) Salt tolerance of trees and shrubs. Univ. Vt. Ext. Serv. Brieflet 1212.

924. Perevertailo, B.I.

1971. Root system of *Juglans nigra* in young plantations. [Kornevaya sistema orekha chernogo v molodykh kul'turakh]. Transl., Can. Dep. Sec. State Transl. Bur., Foreign Lang. Div. *from* Lesn. Zh. 12(2):151–154 (1969). (Transl. Environ. Can. No. OOENV–10). FA31:4113.

925. Perry, G.S.

1932. Some tree antagonisms. Proc. Pa. Acad. Sci. 6:136-141.

926. Perry, T.O.

1971. Winter-season photosynthesis and respiration by twigs and seedlings of deciduous and evergreen trees. For. Sci. 17(1):41–43. FA33:122.

927. Peterson, E.B

1965. Inhibition of black spruce primary roots by a water-soluble substance in *Kalmia angustifolia*. For. Sci. 11(4):473–479. FA27:5571.

928. Peterson, L.O.T., and H.A. Worden.

1968. Root borers in poplar cuttings beds. Summ. Rep. Tree Nursery Sask. 1967, p. 7–9. FA29:5949.

929. Pharis, R.P., R.L. Barnes, and A.W. Naylor.

1964. Effects of nitrogen level, calcium level and nitrogen source upon the growth and composition of *Pinus taeda* L. Physiol. Plant. 17(3):560–572. FA26:4868.

930. Pillai, A.

1964. Root apical organization in gymnosperms—some conifers. Bull. Torrey Bot. Club 91(1):1–13. BA45:56822.

931. Pizelle, G.

1975. [Seasonal variations in the nitrogenase activity of root nodules in *Alnus glutinosa*, *A. incana*, and *A. cordata*.] C. R. Hebd. des Séances de l'Acad. Sci., France, D 281(23):1829–1832. [In Fr. with Engl. summ.] FA37:6016.

932. Platt, W.D., E.B. Cowling, and C.S. Hodges, Jr.

1964. Resistance of coniferous root wood and stem wood to decay by *Fomes annosus*. (Abstr.) Phytopathology 55(2):130–131. BA46:72910.

933. Platt, W.D.

1965. Comparative resistance of coniferous root wood and stem wood to decay by isolates of *Fomes annosus*. Phytopathology 55(12):1347–1353. BA47:49395.

934. Plice, M.J., and G.W. Hedden.

1931. Selective girdling of hardwoods to release young growth of conifers. J. For. 29(1):32–40.

935. Plowman, A.B.

1915. Is the box-elder a maple? A study of the comparative anatomy of *Negundo*. Bot. Gaz. 60:169–172.

936. Pomerleau, R., and M. Lortie.

1956. Relation entre le dépérissement du bouleau blanc et l'état des racines. [Relation between the dieback of white birch and the condition of the roots.] (Abstr.) Ann. ACFAS (Assoc. Con. Fr. Av. Sci.) 22:55. FA19:3257.

937. Pomerleau, R., and M. Lortie.

1962. Relationships of dieback to the rooting depth of white birch (Betula papyrifera). For. Sci. 8(3):219–224. BA41:16423.

938. Pomeroy, K.B.

1949. The germination and initial establishment of loblolly pine under various surface soil conditions. J. For. 47(7):541–543.

939. Potzger, J.E.

1937. Vegetative reproduction in conifers. Am. Midl. Nat. 18:1001-1004.

940. Powers, H.R., Jr., and C.S. Hodges, Jr.

1970. Annosus root rot in eastern pines. USDA For. Serv. For. Pest Leafl. 76.

941. Prasad, R.

1975. Protection of elm trees against the Dutch elm disease (*Ceratocystis ulmi* Buism/Moreau) in the urban environments: Some factors affecting uptake and translocation of the systemic fungicide, Benomyl, by roots. Inf. Ref. CC–X Chem. Control Res. Inst. 133.

942. Prasad, R., and R.P. Moody.

1974. A preliminary study on the chemical control of maple roots (*Acer saccharinum* L.) with herbicides. Chem. Control Res. Inst., Can. Rep. CC-X-77. FA36:6989.

943. Pritchett, W.L.

1972. The effect of nitrogen and phosphorus fertilizers on the growth and composition of loblolly and slash pine seedlings in pots. Proc. Soil and Crop Sci. Soc. Fla. 32:161–165. FA36:3944.

944. Pronos, J., and R.F. Patton.

1977. Armillaria root rot of red pine planted on oak sites in Wisconsin. Plant Dis. Rep. 61(11):955–958.

945. Pudova, R.A.

1974. [Effect of seed pre-treatment with trace elements on the growth of the root system in *Fraxinus pennsylvanica*.] Byull. Gl. Bot. Sada 92, p. 74–79. [In Russ.] FA36:2006.

946. Pulling, H.E.

1918. Root habit and plant distribution in the far north. Plant World 21:223–233.

947. Raabe, R.D.

1962. Host list of the root rot fungus, *Armillaria mellea*. Hilgardia 33(2):25–88. FA24:5165.

948. Raju, M.V.S., R.T Coupland, and T.A. Steeves. 1966. On the occurrence of root buds on perennial plants in Saskatchewan. Can. J. Bot. 44(1):33–37. FA27:5572.

949. Ralston, C.W.

1973. Annual primary productivity in a loblolly pine plantation. *In* Biomass studies. H.E. Young, ed. Int. Union For. Res. Organ. Congr. Univ. Maine Press, Orono. p. 105–117.

Ralston, C.W., R.C. Chapman, and R.S. Kinerson, Jr.
 1972. Biomass distribution in a loblolly pine plantation. US-IBP
 East. Deciduous For. Biome Memo. Rep. 72-79.

951. Rangnekar, P.V., and D.F. Forward. 1972. Foliar nutrition and growth in re-

1972. Foliar nutrition and growth in red pine: Distribution of photo-assimilated carbon in seedlings during bud expansion. Can. J. Bot. 50(10):2053–2061. FA34:3233.

952. Rathbun-Gravatt, A.

1931. Germination loss of coniferous seeds due to parasites. J. Agric. Res. 42(2):71-92.

953. Redmond, D.R.

1954. Increased soil temperatures cause rootlet mortality in yellow birch. Bi-mon. Progr. Rep. Div. For. Biol., Dep. Agric. Can. 10(3):1. FA16:740.

954. Redmond, D.R.

1954. Variations in development of yellow birch roots in two soil types. For. Chron. 30(4):401–406. BA29:26970.

955. Redmond, D.R.

1955. Soil temperatures and yellow birch decline. Bi-mon. Progr. Rep. Div. For. Biol, Dep. Agric. Can. 11(5):1. FA17:1825.

956. Redmond, D.R.

1955. Studies in forest pathology. XV. Rootlets, mycorrhiza, and soil temperatures in relation to birch dieback. Can. J. Bot. 33(6):595-627.

957. Redmond, D.R.

1957. Observations on rootlet development in yellow birch. For. Chron. 33(3):208-212. BA32:13776.

958. Redmond, D.R.

1957. The future of birch from the viewpoint of diseases and insects. For. Chron. 33(1):25–30.

959. Redmond, D.R.

1959. Mortality of rootlets in balsam fir defoliated by the spruce budworm. For. Sci. 5(1):64–69. BA33:35286.

960. Reed, J.F.

1939. Root and shoot growth of shortleaf and loblolly pines in relation to certain environmental conditions. Duke Univ. Sch. For. Bull. 4, Durham N.C. BA14:14321.

961. Regehr, D.L., F.A. Bazzaz, and W.R. Boggess.

1975. Photosynthesis, transpiration and leaf conductance of *Populus deltoides* in relation to flooding and drought. Photosynthetica 9(1):52-61.

962. Reid, C.P.P.

1968. Nutrient transfer by mycorrhizae. Diss. Abstr. 29B(2):429. FA30:3695.

963. Reid, C.P.P., and W. Hurtt.

1970. Root exudation of herbicides by woody plants: Allelopathic implications. Nature 225(5229):291. FA31:4710.

964. Reid, C.P.P., and F.W. Woods.

1967. Atmospheric transfer of carbon-14. A problem in fungus translocation studies. Science 157(3789):712-713.

965. Reid, C.P.P., and F.W. Woods.

1969. Translocation of C¹⁴-labeled compounds in mycorrhizae and its implications in interplant nutrient cycling. Ecology 50(2):179–187.

966. Reineke, L.H.

1942. Effect of pregermination and radicle damage on first-year development of red oak. J. For. 40(4):346-347. FA4-102.

967. Reineke, L.H.

1942. Effect of stocking and seed on nursery development of eastern white pine seedlings. J. For. 40(7):577–578. FA4:166.

968. Reines, M., and J.H. Bamping.

1960. Seasonal rooting responses of slash and loblolly pine cuttings. J. For. 58(8):646–647. BA35:64633.

969. Rexrode, C.O., and R.E. Frame.

1977. Root graft incidence at oak wilt sites in West Virginia. Plant Dis. Rep. 61(11):970-971.

970. Richard, C., and J.A. Fortin.

1970a. [The rootlets of *Picea mariana*: Symbiosis and pathology.] (Abstr.) Ann. ACFAS (Assoc. Can. Fr. Av. Sci.) 37(suppl.):9. [In Fr.] FA37:3607.

971. Richard, C., and J.A. Fortin.

1970b. Les mycorrhizes de *Picea mariana* (Mill.) BSP: Aspects morphologiques, anatomiques et systématiques. Nat. Can. 97:163–173. [Engl. summ.]

972. Richard, C., and J.A. Fortin.

1974. Distribution géographique, écologie, physiologie, pathogenécité et sporulation du *Mycelium radicis atrovirens*. [Geographical distribution, ecology, physiology, pathogenicity and sporulation of *M. radicis atrovirens*.] Phytoprotection 55(2):67–88. [Engl. summ.] FA36:4856.

973. Richard, C., and J.A. Fortin.

1975. Rôle protecteur du Suillus granulatus contre le Mycelium radicis atrovirens sur des semis de Pinus resinosa. [The protective role of S. granulatus against M. radicis atrovirens on seedlings of P. resinosa.] Can. J. For. Res. 5(3):452–456. [Engl. summ.]

974. Richard, C., J.A. Fortin, and A. Fortin.

1971. Protective effect of an ectomycorrhizal fungus against the root pathogen *Mycelium radicis atrovirens*. Can. J. For. Res. 1(4):246–251.

975. Richards, B.N.

1961. Soil pH and mycorrhiza development in *Pinus*. Nature 190(4770):105–106.

976. Richards, B.N.

1962. Increased supply of soil nitrogen brought about by *Pinus*. Ecology 43(3):538–541. FA24:2022.

977. Richards, B.N.

1965. Mycorrhiza development of loblolly pine seedlings in relation to soil reaction and the supply of nitrate. Plant and Soil 22(2):187–199. FA27:1862.

978. Richards, N.A., A.L. Leaf, and D.H. Bickelhaupt.

1973. Growth and nutrient uptake of coniferous seedlings: Comparison among ten species at various seedbed densities. Plant and Soil 38(1):125–143. BA56:31324.

979. Richardson, S.D.

1953. A note on some differences in root-hair formation between seedlings of sycamore and American oak. New Phytol. 52(1):80–82. BA28:4208.

980. Richardson, S.D.

1953. Studies of root growth in *Acer saccharinum* L. I. The relation between root growth and photosynthesis. Proc. K. Ned. Akad. Wet. Ser. C. 56(2):185–193. BA28:1884.

981. Richardson, S.D.

1953. Studies on root growth in *Acer saccharinum* L. II. Factors affecting root growth when photosyntheses is curtailed. Proc. K. Ned. Akad. Wet. Ser. C 56(3):346–353. FA15:147.

982. Richardson, S.D.

1954. Root growth of *Acer saccharinum* seedlings during winter. Proc. 8th Int. Bot. Congr., [Paris 1954], Part 5, Section 11, p. 303–304. FA16:233.

983. Richardson, S.D.

1955. The influence of rooting medium on the structure and development of the root-cap in seedlings of *Acer saccharinum* L. New Phytol. 54(3):336–337. BA30:29554.

984. Richardson, S.D.

1956a. On the role of the acorn in root growth of American oak seedlings. Meded. Landbouwhogesch Wageningen. 56(12):1–18. BA32:17509.

985. Richardson, S.D.

1956b. Studies of root growth in *Acer saccharinum* L. III. The influence of seedling age on the short-term relation between photosynthesis and root growth. IV. The effect of differential shoot and root temperatures on root growth. Proc. K. Ned. Akad. Wet. Ser. C 59(3):416–438. FA18:209.

986. Richardson, S.D.

1956c. Studies of root growth in *Acer saccharinum* L. V. The effect of a long-term limitation of photosynthesis on root growth rate in first-year seedlings. Proc. K. Ned. Akad. Wet. Ser. C 59(5):694–701. FA18:3788.

987. Richardson, S.D.

1957. Studies of root growth in *Acer saccharinum* L. VI. Further effects of the shoot system on root growth. Proc. K. Ned. Akad. Wet. Ser. C 60(5):625–629. FA19:2815.

988. Richardson, S.D.

1958a. Bud dormancy and root development in *Acer saccharinum*. *In* The physiology of forest trees. A symposium. K.V. Thimann, ed. Maria Moors Cabot Found. for Bot. Res. Int. Symp. For. Tree Physiol., Harvard Univ., Cambridge, Mass., Ronald Press Co., New York. p. 409–425.

989. Richardson, S.D.

1958b. The effect of IAA on root development of *Acer saccharinum* L. Physiol. Plant. 11(4):698–709. BA33:19383.

990. Richardson, S.D.

1961. Some root-shoot interrelationships. *In* Recent advances in botany. Proc. 9th Int. Bot. Congr., [Montreal 1959], 2:1321–1323.

991. Riedacker, A.

1976. Étude bibliographique: Rythmes de croissance et de régéneration des racines des végétaux ligneaux. [Bibliographic study: Rhythms of growth and of root regeneration of woody plants.] Ann. Sci. For. 33(3):109–138. [Engl. and Ger. summ.] BA64:16914.

992. Riffle, J.W., and F.C. Strong.

1960. Studies in white pine seedling root rot. Mich. Agric. Exp. Stn. Q. Bull. 42(4):845–853. FA22:710.

993. Riley, C.G., W.B. Denyer, and R.D. Whitney.

1952. 'Littleleaf' of white elm. Can. Dep. Agric. Entomol. Branch Annu. Rep. For. Insect and Dis. Surv. 1951:142. FA14:393.

994. Rishbeth, J.

1972. Resistance to fungal pathogens of tree roots. Proc. R. Soc. Lond., B. 181(1064):333–351. FA34:385.

.995. Roberts, B.R., and L.R. Schreiber.

1977. Influence of Dutch elm disease on resistance to water flow through roots of American elm. Phytopathology 67(1):56–59.

996. Robertson, G.I.

1969. Root rot resistant trees and shrubs. N.Z. Gard. 26(9):34–35, 37. FA31:6679.

997. Rodriguez-Barrueco, C.

1969. The occurrence of nitrogen-fixing root nodules on non-leguminous plants. Bot. J. Linn. Soc. 62(1):77–84. FA31:2173.

998. Rodriguez-Barrueco, C.

1971. [The effect of combined nitrogen on the development of nodules and the growth of *Elaeagnus angustifolia*.] An. Edafol. Agrobiol. 30(1/2):149–161. [In Span. with Engl. summ.] FA33:4054.

999. Röhrig, E.

1967. Root development of forest trees in relation to ecological conditions—Part I and II. [Die Wurzelentwichlung der Waldbäume in Abhängigkeit von den ökologischen Verhältnissen—I. and II. Teil.] Transl., Can. Dep. Sec. State Transl. Bur., Foreign Lang. Div. from Forstarchiv 37(10):217–229, 37(11):237–249 (1966). FA29:291 (Transl. Dep. For. Can. 101).

1000. Romberg, L.D., and C.L. Smith.

1938. Effects of indole-3-butyric acid in the rooting of transplanted pecan trees. Proc. Am. Soc. Hort. Sci. 36:161–170. BA13:13987.

1001. Rosendahl, R.O.

1942. The effect of mycorrhizal and non-mycorrhizal fungi on the availability of difficultly soluble potassium and phosphorus. Proc. Soil Sci. Soc. Am. 7:477–479.

1002. Ross, W.D.

1976. Relation of aspen root size to infection by *Ganoderma applanatum*. Can. J. Bot. 54(8):745–751.

1003. Roth, E.B., T.S. Buchanan, and G.H. Hepting.

1948. A five-year record of littleleaf on thirty-one plots. For. Path. Spec. Release U.S. Div. For. Path. 32. FA10:2991.

1004. Roth, E.R.

1952. Roots of living *Pinus rigida* decayed by *Fomes annosus*. Plant Dis. Rep. 36:(8):330. FA14:2386.

1005. Roth, L.F., and A.J. Riker.

1943. Influence of temperature, moisture, and soil reaction on the damping-off of red pine seedlings by *Pythium* and *Rhizoctonia*. J. Agric. Res. 67(7):273–293.

1006. Routien, J.B., and R.F. Dawson.

1943. Some interrelationships of growth, salt absorption, respiration, and mycorrhizal development in *Pinus echinata* Mill. Am. J. Bot. 30(6):440–451.

1007. Rowan, S.J.

1960. The susceptibility of 23 tree species to black root rot. Plant Dis. Rep. 44(8):646–647. BA36:38796.

1008. Rudolf, P.O.

1937. Diagnosing plantation mortality. Pap. Mich. Acad. Sci. Arts and Lett. 23:333–338. BA12:7066.

1009. Rudolf, P.O.

1939. Why forest plantations fail. J. For. 37(5):377-383. FA1:162.

1010. Rudolf, P.O.

1950. Forest plantations in the Lake States. USDA Tech. Bull. 1010.

1011. Ruehle, J.L.

1961. Investigation of the relationships of plant-parasitic nematodes to the stunting of pines in outplantings. Diss. Abstr. 22(1):22–23. FA23:2468.

1012. Ruehle, J.L.

1965. Host range studies of several plant-parasitic nematodes found in southern pine forests. (Abstr.) Nemotologica 11(1):45. FA27:2569.

1013. Ruehle, J.L.

1966. Nematodes parasitic on forest trees. I. Reproduction of ectoparasites on pines. Nematologica 12(3):443–447.

1014. Ruehle, J.L.

1969. Influence of stubby-root nematode on growth of southern pine seedlings. For. Sci. 15(2):130–134. FA31:1093.

1015. Ruehle, J.L.

1969. Nematodes parasitic on forest trees. II. Reproduction of endoparasites on pines. Nematologica 15(1):76–80. FA32:4633.

1016. Ruehle, J.L., and D.H. Marx.

1971. Parasitism of ectomycorrhizae of pine by lance nematode. For. Sci. 17(1):31–34. FA33:1181.

1017. Ruehle, J.L., and D.H. Marx.

1977. Developing ectomycorrhizae on containerized pine seedlings. USDA For. Serv. Res. Note SE-242.

1018. Ruehle, J.L., and J.N. Sasser.

1960. The relationship of plant-parasitic nematodes to the growth of pines in outplantings. (Abstr.) Phytopathology 50(9):652. FA22:2171.

1019. Rushmore, F.M.

1956. Beech root sprouts can be damaged by sodium arsenite treatment of parent tree. USDA For. Serv. Res. Note 57. Northeast. For. Exp. Stn.

1020. Russo, V.M., and W.L. Klarman.

1973. The effect of air and methane gas on root development of *Pinus virginiana* and on mycorrhiza formation with *Amanita rubescens*. (Abstr.) Phytopathology 63(7):805. FA35:1380.

1021. Safford, L.O.

1974. Effect of fertilization on biomass and nutrient content of fine roots in a beech-birch-maple stand. Plant and Soil 40(2):349–363. BA58:54326.

1022. Safford, L.O.

1976. Seasonal variation in the growth and nutrient content of yellow birch replacement roots. Plant and Soil 44(2):439–444. BA62:48431.

1023. Safford, L.O., and S. Bell.

1972. Biomass of fine roots in a white spruce plantation. Can. J. For. Res. 2(3):169–172. BA55:25543.

1024. Saksena, H.K., and O. Vaartaja.

1961. Taxonomy, morphology, and pathogenicity of *Rhizoctonia* species from forest nurseries. Can. J. Bot. 39(3):627–647. FA23:2241.

1025. Salonius, P.O.

1977. Nutrient concentration in black spruce foliage immediately following fertilization. Soil Sci. Soc. Am. J. 41(1):136–139. FA39:48.

1026. Sandburg, D.

1951. The regeneration of quaking aspen by root suckering. M.F. thesis, Univ. Minn.

1027. Sandburg, D., and A.E. Schneider.

1953. The regeneration of aspen by suckering. Minn. For. Notes 24.

1028. Santamour, F.S., Jr.

1960. The relation between anatomy and inherent growth potential of hybrid poplars. Diss. Abstr. 21(4):718. FA22:2777.

1029. Santamour, F.S., Jr.

1965. Rooting of pitch pine stump sprouts. Tree Plant. Notes 70, p. 7–8. FA27:538.

1030. Santantonio, D., R.K. Hermann, and W.S. Overton.

1977. Root biomass studies in forest ecosystems. Pedobiologia 17(1):1–31.

1031. Sasaki, S., and T.T. Kozlowski.

1968. Effects of herbicides on seed germination and early seedling development of *Pinus resinosa*. Bot. Gaz. 129(3):238–246. BA50:32959.

1032. Sasaki, S., and T.T. Kozlowski.

1968. Effects of ipazine, EPTC, and 2,4–D on respiration of root tips of red pine (*Pinus resinosa*) plants of different ages. Adv. Front. Plant Sci. 21:135–140. BA50:89070.

1033. Schantz-Hansen, T.

1945. The effect of planting methods on root development. J. For. 43(6):447-448. BA20:3738.

1034. Schier, G.A.

1972. Apical dominance in multishoot cultures from aspen roots. For. Sci. 18(2):147–149. BA54:54456.

1035. Schier, G.A.

1973. Effects of gibberellic acid and an inhibitor of gibberellin action on suckering from aspen root cuttings. Can. J. For. Res. 3(1):39–44. FA34:6835.

1036. Schier, G.A.

1973. Origin and development of aspen root suckers. Can. J. For. Res. 3(1):45–53. BA56:48894.

1037. Schier, G.A.

1973. Seasonal variation in sucker production from excised roots of *Populus tremuloides* and the role of endogenous auxin. Can. J. For. Res. 3(3):459–461. BA57:54623.

1038. Schier, G.A.

1975. Promotion of sucker development on *Populus tremuloides* root cuttings by an antiauxin. Can. J. For. Res. 5(2):338–340. BA60:48890.

1039. Schier, G.A.

1978. Variation in suckering capacity among and within lateral roots of an aspen clone. USDA For. Serv. Res. Note INT-241.

1040. Schier, G.A., and R.B. Campbell.

1976. Differences among *Populus* species in ability to form adventitious shoots and roots. Can. J. For. Res. 6(3):253–261. BA63:20510.

1041. Schier, G.A., and R.S. Johnston.

1971. Clonal variation in total nonstructural carbohydrates of trembling aspen roots in three Utah areas. Can. J. For. Res. 1(4):252–255. BA54:16568.

1042. Schier, G.A., and J.C. Zasada.

1973. Role of carbohydrate reserves in the development of root suckers in *Populus tremuloides*. Can. J. For. Res. 3(2):243–250. BA56:63206.

1043. Schmiege, D.C.

1959. Pine root collar weevil. USDA For. Serv. For. Pest Leaflet 39.

1044. Schmitt, D.M.

1954. Early root development in black spruce (*Picea mariana*). (Abstr.) For. Chron. 30(1):105. FA15:3348.

1045. Schmitz, H., and L.W.R. Jackson.

1927. Heartrot of aspen with special reference to forest management in Minnesota. Minn. Agric. Exp. Stn. Tech. Bull. 50.

1046. Schoeneweiss, D.F., and A.C. Valcarce.

1966. *Fomes annosus* root rot in Illinois pine plantings. Plant Dis. Rep. 50(6):446–448. BA47:104379.

1047. Scholten, H.

1968. Factors affecting the infection of black spruce (*Picea mariana* Mill.) transplants by *Cylindrocladium scoparium*. Diss. Abstr. 28B(8):3131. FA29:5821.

1048. Scholtes, W.H.

1953. The concentration of forest tree roots in the surface zone of some Piedmont soils. Proc. Ia. Acad. Sci. 60:243–259. FA16:3885.

1049. Schramm, J.R.

1966. Plant colonization studies on black wastes from anthracite mining in Pennsylvania. Trans. Am. Phil. Soc. (N.S.) 56(1).

1050. Schreiber, L.R., and B.R. Roberts.

1971. Intraclonal vs. intraseedling variation in *Ulmus americana* L. and *U. pumila* L. J. Am. Soc. Hort. Sci. 96(1):115–116. BA52:66258.

1051. Schreiner, E.J.

1949. Can black walnut poison pines? Morris Arbor. Bull. 4(11):94–96. FA11:3260.

1052. Schultz, J.D.

1969. The vertical rooting habit in black spruce, white spruce, and balsam fir. Diss. Abstr. Int. 30B(5):1976. FA32:5632.

1053. Schultz, R.P.

1966. The frequency and implications of intraspecific root-grafting in loblolly pine (*Pinus taeda* L.). Diss. Abstr. 26(8):4148. FA28:319.

1054. Schultz, R.P., and F.W. Woods.

1967. The frequency and implications of intraspecific root-grafting in loblolly pine. For. Sci. 13(3):226–239. BA49:16852.

1055. Scott, S.J., and G.W. Hay.

1967. The carbohydrates of the roots of the sugar maple. Can. J. Chem. 45(19):2217–2225. FA29:4582.

1056. Scully, N.J.

1942. Root distribution and environment in a maple-oak forest. Bot. Gaz. 103(3):492–517. BA16:17504.

1057. Secrest, H.C., H.J. MacAloney, and R.C. Iorenz.

1941. Causes of the decadence of hemlock at the Menominee Indian Reservation, Wisconsin. J. For. 39(1):3–12.

1058. Seeliger, I.

1959. Über die Bildung wurzelburtiger Sprosse und das Wachstum isolierter Wurzeln der Robinie (*Robinia pseudoacacia* L.). [On the formation of shoots by roots and the growth of isolated roots in *R. pseudoacacia*.] Flora Allg. Bot. Ztg. 148(2):218–254. BA37:15243.

1059. Seeliger, I.

1967. Versuche zur Knoellchenbildung an isolierten Wurzels der Robinie (*Robinia pseudoacacia* L.) [Nodule formation experiments on isolated roots of R. *pseudoacacia*.] Flora Allg. Bot. Ztg. 157(6):561–565. BA49:41573.

1060. Seidel, K.W.

1972. Drought resistance and internal water balance of oak seedlings. For. Sci. 18(1):34–40. BA54:19478.

1061. Shapiro, S

1949. Reversal of polarity in regenerated roots on stem cuttings of *Populus*. (Abstr.) Am. J. Bot. 36(10):803. BA24:13514.

1062. Shapiro, S.

1957. Auxin control of seasonal polarization of root emergence. (Abstr.) Plant Physiol. 32(suppl.):xlii. FA20:441.

1063. Shapiro, S.

1958. The role of light in the growth of root primordia in the stem of the lombardy poplar. *In* The physiology of forest trees. A symposium. K.V. Thimann, ed., Maria Moors Cabot Found. for Bot. Res. Int. Symp. For. Tree Physiol., Harvard Univ., Ronald Press Co., New York. p. 445–465.

1064. Shaw, K.

1977. Girdling roots common in container-grown plants, wood and forest trees. Arnoldia 37(5):242–247.

1065. Shea, S.R.

1975. Growth and development of jack pine (*Pinus banksiana* Lamb.) in relation to edaphic factors in northeastern Ontario. Diss. Abstr. Int. 36B(5):2048–2049. FA37:4672.

1066. Shenefelt, R.D.

1956. A further note on protecting machine transplanted trees from white grubs. Univ. Wis. For. Res. Notes 25.

1067. Shigo, A.L.

1972. Successions of microorganisms and pattern of discoloration and decay after wounding in red oak and white oak. Phytopathology 62(2):256–259.

1068. Shigo, A.L.

1975. Compartmentalization of decay associated with *Fomes annosus* in trunks of *Pinus resinosa*. Phytopathology 65(9):1038–1039. FA37:2989.

1069. Shirley, H.L.

1932. Does light burning stimulate aspen suckers? II. J. For. 30(4):419-420.

1070. Shirley, H.L.

1941. Restoring conifers to aspen lands in the Lake States. USDA Tech. Bull. 763.

1071. Shirley, H.L.

1945. Reproduction of upland conifers in the Lake States as affected by root competition and light. Am. Midl. Nat. 33(3):537–612. BA19:15518.

1072. Shirley, H.L., and L.J. Meuli.

1939. Influence of moisture supply on drought resistance of conifers. J. Agric. Res. 59(1):1–21. FA1:157.

Shiroya, T., G.R. Lister, V. Slankis, G. Krotkov, and C.D. Nelson.
 1962. Translocation of the products of photosynthesis to roots of pine seedlings. Can. J. Bot. 40(8):1125–1135. BA40:16467.

074. Shoulders, E., and J.R. Jorgensen.

1969. Mycorrhizae increase field survival of planted loblolly pine. Tree Plant. Notes 20(1):14–17. FA31:374.

1075. Šika, A.

1957. Vývoj kořenú ořešáku černého a vlašského v mládi. [Root development in young plants of *Juglans nigra* and *J. regia*.] Sborn. csl. Akad. zemed. (Lesn.) 3(11):769–784. [In Czech. with Russ. summ.] FA19:3949.

1076. Šika, A.

1959. Korenovy system ořešáku černého. [The root system of *Juglans nigra*.] Sborn. csl. Akad. zemed. (Lesn.) 5(3):267–290. [In Czech. with Russ. and Ger. summ.] FA20:4281.

1077. Silverborg, S.B., and R.L. Gilbertson.

1961. *Armillaria mellea* root rot in a northern white pine plantation. Plant Dis. Rep. 45(5):389. BA36:73962.

1078. Silverborg, S.B., J.R. Risley, and A.L. Leaf.

1962. A basal canker disease of eastern white pine. Plant Dis. Rep. 46(4):285–286. FA24:2498.

1079. Sims, H.P.

1964. Root development of jack pine seedlings on burned-over dry sites in southeastern Manitoba. Can. Dep. For. Publ. 1061. FA26:314.

1080. Sims, H.P., and D. Mueller-Dombois.

1968. Effect of grass competition and depth to water table on height growth of coniferous tree seedlings. Ecology 49(4):597–603. FA30:2012.

1081. Sinclair, W.A.

1963. The root- and butt-rot of conifers cuased by *Fomes annosus* with special reference to inoculum dispersal and control of the disease in New York. Diss. Abstr. 23(8):2655. BA43:25028.

1082. Sinclair, W.A., E.L. Stone, and C.F. Scheer, Jr.

1975. Toxicity to hemlocks grown in arsenic-contaminated soil previously used for potato production. HortScience 10(1):35–36. FA37:359.

1083. Singer, F.P., and R.J. Hutnik.

1965. Excavating roots with water pressure. J. For. 63(1):37–38. FA26:4876.

1084. Singh, P.

1970. Armillaria root rot in a coniferous plantation in Newfoundland. Bi-mon. Res. Notes 26(1):5–6.

1085. Singh, P.

1975. Armillaria root rot: Distribution and severity in softwood plantations in Newfoundland. Acta Phytopathol. Acad. Sci. Hung. 10(3–4):389–406.

1086. Singh, P., and N.D. Bhure.

1974. Influence of *Armillaria* root rot on the foliar nutrients and growth of some coniferous species. Eur. J. For. Pathol. 4(1):20–26. FA36:958.

1087. Singh, P., and J. Richardson.

1973. Armillaria root rot in seeded and planted areas in Newfoundland. For. Chron. 49(4):180–182.

1088. Skelly, J.M., S.A. Alexander, and C.L. Morris.

1974. Excavation of entire tree root systems reveals higher incidences of *Fomes annosus*. Proc. Am. Phytopath. Soc. 1:155. (Abstr.) FA37:1709.

1089. Skilling, D.

1965. Root rot on conifers in the Lake States. USDA For. Serv., Region 9 Nurserymen's Conf. p. 4-11.

1090. Slankis, V.

1958. The role of auxin and other exudates in mycorrhizal symbiosis of forest trees. *In* The physiology of forest trees. A symposium. K.V. Thimann, ed. Maria Moors Cabot Found. for Bot. Res. Int. Symp. For. Tree Physiol., Harvard Univ., Ronald Press Co., New York. p. 427–443.

1091. Slankis, V.

1961. On the factors determining the establishment of ectotrophic mycorrhiza of forest trees. *In* Recent advances in botany, 9th Int. Bot. Congr. [Montreal 1959] 2:1738–1742.

1092. Slankis, V

1967. Renewed growth of ectotrophic mycorrhizae as an indication of an unstable symbiotic relationship. 14th Int. Union For. Res. Organ. Congr. Proc. [Munich 1967] Part V, Section 24, p. 84–99. FA29:1989.

Slankis, V., V.C. Runeckles, and G. Krotkov.
1964. Metabolites liberated by roots of white pine (*Pinus strobus* L.) seedlings. Physiol. Plant. 17(2):301–313. FA26:126.

1094. Slocum, G.K., and T.E. Maki.

1956a. Exposure of loblolly pine planting stock. J. For. 54(5):313–315. FA17:3899.

1095. Slocum, G.K., and T.E. Maki.

1956b. Some effects of depth of planting upon loblolly pine in the North Carolina Piedmont. J. For. 54(1):21–25. FA17:2748.

1096. Smerlis, E.

1961. Pathological condition of immature balsam fir stands of *Hylocomium-Oxalis* type in the Laurentide Park, Quebec. For. Chron. 37(2):109–115.

1097. Smith, C.L., J. Hamilton, C.J.B. Thor, and L.D. Romberg. 1939. Root composition and top development in large pecan trees headed to various degrees of severity in top working. J. Agric. Res. 58:821–842.

1098. Smith, C.L., and J.G. Waugh.

1938. Seasonal variations in the carbohydrate and nitrogen content of roots of bearing pecan trees. J. Agric. Res. 57(6):449–460. BA12:14371.

1099. Smith, D.M.

1951. The influence of seedbed conditions on the regeneration of eastern white pine. Conn. Agric. Exp. Stn. Bull. 545.

1100. Smith, W.H.

1969. Release of organic materials from the roots of tree seedlings. For. Sci. 15(2):138–143. FA31:170.

1101. Smith, W.H.

1970. Root exudates of seedling and mature sugar maple. Phytopathology 60(4):701–703. BA51:104146.

1102. Smith, W.H

1972. Influence of artificial defoliation on exudates of sugar maple. Soil. Biol. and Biochem. 4(1):111–113. FA33:3813.

1103. Smith, W.H.

1976. Character and significance of forest tree root exudates. Ecology 57(2):324–331.

1104. Smyly, W.B., and T.H. Filer, Jr.

1977. Cylindrocladium scoparium associated with root rot and mortality of cherrybark oak seedlings. Plant Dis. Rep. 61(7):577–579. BA64:59156.

1105. Snow, A.G., Jr.

1939. Clonal variation in rooting response of maple cuttings. USDA For. Serv. Tech. Note 29. Northeast. For. Exp. Stn.

1106. Spaulding, P

1952. Root rots of conifers. *In* Important tree pests of the Northeast. New Engl. Sect. Soc. Amer. For. Evans Printing Co., Concord, N.H. p. 138–140.

1107. Spaulding. P., and H.J. MacAloney.

1931. A study of organic factors concerned in the decadence of birch on cut-over lands in northern New England. J. For. 29(8):1134–1149.

1108. Speers, C.F.

1974. Pales and pitcheating weevils: Development in relation to time pines are cut in the Southeast. USDA Serv. Res. Note SE–207. FA37:1786.

1109. Spurr, A.R.

1949. Histogenesis and organization of the embryo in *Pinus strobus* L. Am. J. Bot. 36(9):629–641. BA24:24778.

1110. Spyridakis, D.E., G. Chesters, and S.A. Wilde.

1967. Kaolinization of biotite as a result of coniferous and deciduous seedling growth. Proc. Soil Sci. Soc. Am. 31:203–210.

1111. Stanek, W.

1967. Natural layering of red spruce in Quebec. Bi-mon. Res. Notes 23(6):42.

1112. Starikov, R.

1969. [Features of the growth of the root systems of certain woody species in nurseries.] Lesn. Hozjajstvo 22(12):28–30. [In Russ.] FA31:4107.

1113. Stark, R.W.

1959. Studies on the pine root weevil, *Hylobius warreni* Wood, in Alberta (1957). Can. Dep. Agric. Sci. Serv. Interim Rep., For. Biol. Div. For Biol. Lab. Calgary, Alberta.

1114. Steinbeck, K., and P.P. Kormanik.

1968. First-year root system development of two clones of yellow-poplar. Ga. For. Res. Pap. 55. BA51:63428.

1115. Steinbeck, K., and R.G. McAlpine.

1966. Inter- and intra-specific differences in the root respiration rates of four hardwood species. For. Sci. 12(4):473-476. BA48:40797.

1116. Sterner, T.E.

Inhibition of root- and butt-decay fungi by extractives of balsam fir root wood. Can. J. For. Res. 4(2):213–221. BA58:33778.

1117. Sterrett, J.P., W.E. Chappell, and G.M. Shear.

1968. Temperature and annual growth cycle effects on root suckering in black locust. Weed Sci. 16(2):250–251. BA49:85782.

1118. Sterrett, W.D.

1920. Jack pine. USDA Bull. 820.

1119. Stevens, C.L.

1931. Root growth of white pine (*Pinus strobus* L.). Yale Univ. Sch. For. Bull. 32. BA9:5697.

1120. Stiell, W.M.

1970. Some competitive relations in a red pine plantation. Can. Dep. For. Publ. 1275. BA52:66260.

1121. Stillwell, M.A.

1960. Rootlet recovery in balsam fir defoliated by the spruce budworm. Bi-mon. Progr. Rep. Div. For. Biol., Dep. Agric. Can. 16(5):1. FA22(2117).

1122. Stoddard, E.M., A.D. McDonnell, and H.W. Hicock.

1939. Fomes annosus on conifers in Connecticut. Plant Dis. Rep. 23:385–386.

1123. Stoeckeler, J.H., and C. Arbogast, Jr.

1955. Forest management lessons from a 1949 windstorm in northern Wisconsin and Upper Michigan. USDA For. Serv. Stn. Pap. 34, Lake States For. Exp. Stn.

1124. Stoeckeler, J.H., and G.A. Limstrom.

1942. Ecological factors influencing reforestation in northern Wisconsin. Ecol. Monogr. 12(2):191–212.

1125. Stone, E.L., Jr.

1974. The communal root system of red pine: Growth of girdled trees. For. Sci. 20(4):294–305. BA60:2052.

1126. Stone, E.L., Jr.

1977. Abrasion of tree roots by rock during wind stress. For. Sci. 23(3):333–336.

1127. Stone, E.L., Jr., and T. Greweling.
1971. Arsenic toxicity in red pine and the persistence of arsenic in nursery soils. Tree Plant. Notes 22(1):5-7.

1128. Stone, E.L., Jr., and R.C. McKittrick.

1976. On the layering of white spruce. Tree Plant. Notes 27(1):14.

1129. Stone, E.L., Jr., and M.H. Stone. 1954. Root collar sprouts in pine. J. For. 52(7):487–491.

1130. Stone, E.L., Jr., and V.R. Timmer.
1975. On the copper content of some northern conifers. Can. J.
Bot. 53(15):1453-1456. BA61:2162.

1131. Stone, E.L., Jr., R.R. Morrow, and D.S. Welch.
1954. A malady of red pine on poorly drained sites. J. For.
52(2):104-114.

1132. Stone, J.E., and E.L. Stone, Jr.
1975. Water conduction in lateral roots of red pine. For. Sci.
21(1):53-60. BA60:34161.

1133. Stone, J.E., and E.L. Stone, Jr.
1975. The communal root system of red pine: Water conduction through root grafts. For. Sci. 21(3):255-261. BA61:24909.

1134. Stonecypher, R.W., F.C. Cech, and B.J. Zobel.
1966. Estimates of components of variance and covariance in root and shoot characteristics of loblolly pine after one growing season.
Proc. 8th South. For. Tree Improv. Conf. 1965, p. 86–95.

Stout, B.B.1956. Studies of the root systems of deciduous trees. Black RockFor. Bull. 15. BA32:17462.

1136. Stout, B.B.
1961. Season influences the amount of backflash in a red pine plantation. J. For. 59(12):897-898.

1137. Stout, B.B.
1968. Root systems of deciduous trees. Proc. 43rd Int. Shade Tree
Conf. 1967, p. 30–39. FA30:3705.

1138. Stoutmeyer, V.T., F.L. O'Rourke, and W.W. Steiner.
1944. Some observations on the vegetative propagation of honey-locust. J. For. 42(1):32–36.

1139. Strang, R.M.
1973. The rate of silt accumulation in the lower Peel River, Northwest Territories. Can. J. For. Res. 3(3):457–458. FA35:2132.

Strong, F.C.1941. Root and butt rot in the pinetum at Michigan State College.Mich. Agric. Exp. Stn. Q. Bull. 23(3):159-163. FA3:161.

Strong, F.C.1944. A study of calcium chloride injury to roadside trees. Mich.Agric. Exp. Stn. Q. Bull. 27(2):209-224. FA7:390.

Suggs, E.G., and L.F. Grand.
1972. Formation of mycorrhizae in monoxenic culture by pond pine (*Pinus serotina*). Can. J. Bot. 50(5):1003-1007. FA34:828.
Sutherland, J.R.

1967. Parasitism of *Tylenchus emarginatus* on conifer seedling roots and some observations on the biology of the nematode. Nematologica 13(2):191–196. BA49:10258. FA29:1070.

1144. Sutherland, J.R.
1967. Host range and reproduction of the nematodes *Pratylenchus*projectus, *Pratylenchus* penetrans, and *Tylenchus* emarginatus on
some forest nursery seedlings. Plant. Dis. Rep. 51(2):91-93.
FA29:6060.

1145. Sutherland, J.R.
1967. Failure of the nematode Aphelenchus avenae to parasitize conifer seedling roots. Plant Dis. Rep. 51(5):367-370.
BA48:77293.

1146. Sutherland, J.R.
1969. Feeding of Xiphinema bakeri. Phytopathology 59: 1963–1965.

1147. Sutherland, J.R., and R.E. Adams.
1964. Host range of Tylenchorhynchus claytoni on some forest nursery seedlings. (Abstr.) Phytopathology 54(7):749.
FA26:1002.

1148. Sutherland, J.R., and R.E. Adams.
1965. The parasitism of red pine and other forest nursery crops by Tylenchorhynchus claytoni Steiner. Nematologica 10(4):637-643.
FA26:5547.

1149. Sutherland, J.R., and J.A. Fortin. 1968. Effect of the nematode Aphelenchus avenae on some ectotrophic, mycorrhizal fungi and on a red pine mycorrhizal relationship. Phytopathology 58(4):519-523.

1150. Sutton, R.F.
1967. Influence of planting depth on early growth of conifers.
Commonw. For. Rev. 46(4):282-295. FA29:3821.

1151. Sutton, R.F.
1967. Influence of root pruning on height increment and root development of outplanted spruce. Can. J. Bot. 45(9):1671-1682.
BA49:10174.

1152. Sutton, R.F.
1968. Ecology of young white spruce (*Picea glauca* (Moench) Voss). Diss. Abstr. 29(2):430.

Sutton, R.F.1969. Form and development of conifer root systems. Commonw.For. Bur. Tech. Commun. 7. BA53:57316.

1154. Sutton, R.F., and E.L. Stone, Jr.
1975. White grubs and white spruce establishment. Tree Plant.
Notes. 26(2):9-11.

1155. Swingle, C.F.
1937. Experiments in propagating shipmast locust. J. For.
35(8):713-720.

1156. Szaniawski, R.K., and M.S. Adams.
1974. Root respiration of *Tsuga canadensis* seedlings as influenced by intensity of net photosynthesis and dark respiration of shoots.
Am. Midl. Nat. 91(2):464-468. BA58:22330. FA37:3466.

1157. Tabor, C.A., and C.B. Davey.
1964. Effects of clay-suspension root-coatings upon the survival and growth of tree seedlings. Agron. Abstr. 1964:55.

1158. Tabor, C.A., and C.B. Davey.
1966. Clay-suspension root-coating as antidessicants and rhizo-spheric nutrient sources. Proc. Soil. Sci. Soc. Am. 30(4):516-520.
FA28:2184.

1159. Tainter, F.H., and J.D. Walstad.
1977. Colonization of outplanted loblolly pines by native ectomy-corrhizal fungi. For. Sci. 23(1):77-79.

1160. Taubenhaus, J.J., and W.N. Ezekiel.
1936. A rating of plants with reference to their relative susceptibility to *Phymatotrichum* root rot. Texas Agric. Exp. Stn. Bull. 527.

1161. Taylor, J.S., and E.B. Dumbroff. 1975. Bud, root, and growth-regulator activity in Acer saccharum during the dormant season. Can. J. Bot. 53(4):321-331. FA36:6882.

1162. Tellefsen, M.A.
1922. The relation of age to size in certain root cells and in vein-islets of the leaves of Salix nigra Marsh. Am. J. Bot. 9(3):121-139.

1163. Tew, R.K. 1970. Root carbohydrate reserves in vegetative reproduction of aspen. For. Sci. 16(3):318–320. BA52:1918.

1164. Tew, R.K., N.V. DeByle, and J.D. Schultz.
1969. Intraclonal root connections among quaking aspen trees.
Ecology 50(5):920–921. BA51:74825.

1165. Theobald, W.F., and W.H. Smith. 1974. Nitrate production in two forest soils and nitrate reduction in pine. Proc. Soil Sci. Soc. Am. 38(4):668-672. FA36:1304.

1166. Thimann, K.V., ed. 1958. The physiology of forest trees. A symposium. Maria Moors Cabot Found. for Bot. Res. Int. Symp. For. Tree Physiol., Harvard Univ., Ronald Press Co., N.Y. BA33:19470. 1167. Thomas, H.E.

1934. Studies on Armillaria mellea (Vahl) Quel. Infection, parasitism, and host resistance. J. Agric. Res. 48(3):187–218.

1168. Thomas, W.D., Jr.

1941. The mycorrhizal fungi and mycorrhizae of four coniferous plantations in the Rhine Valley. Phytopathology 31:567–569. FA3:103.

1169. Tint, H.

1945. Studies in the *Fusarium* damping-off of conifers. I. The comparative virulence of certain Fusaria. Phytopathology 35:421–439. FA7:930.

1170. Tint, H.

1945. Studies in the *Fusarium* damping-off of conifers. II. Relation of age of host, pH, and some nutritional factors to the pathogenicity of *Fusarium*. Phytopathology 35:440–457. FA7:930.

1171. Tognoni, F., M. Kawase, and A. Alpi.

1977. Seasonal changes in rootability and rooting substances of *Picea glauca* cuttings. J. Am. Soc. Hort. Sci. 102(6):718–720.

1172. Toole, E.R.

1949. White pine blight in the Southeast. J. For. 47(5):378–382.

1173. Toole, E.R.

1960. Root rot of white oak in Arkansas. Plant Dis. Rep. 44(10):783. BA36:50364.

1174. Toole, E.R.

1966. Root rot caused by *Polyporous lucidus (Ganoderma lucidum*). Plant Dis. Rep. 50(12):945–946. FA28:5868.

1175. Toole, E.R.

1967. Root rot of southern hardwood trees caused by *Corticum galactinum*. Plant Dis. Rep. 51(6):500-501. BA48:103747.

1176. Toronto, University of.

1961. Effect of substrate depth on the growth of seedlings of *Picea mariana*, *Pinus banksiana*, *Larix laricina*, and *Betula lutea*. Rep. For. Res. Glendon Hall Fac. For. Univ. Toronto 1959/60:7. FA22:2871.

1177. Toronto, University of.

1965. Seedling growth in relation to aeration of the root system. Rep. For. Res. Glendon Hall Fac. For. Univ. Toronto 1964/65:5-6. FA27:5526.

1178. Toumey, J.W.

1929. Initial root habit in American trees and its bearing on regeneration. Proc. Int. Congr. Plant Sci. 1:713–728. BA4:9447.

1179. Towers, E

1964. Root-contact infections by *Fomes annosus* in a thinned loblolly pine plantation. Plant Dis. Rep. 48(10):767-769. BA46:95920.

1180. Towers, B., and W.J. Stambaugh.

1968. The influence of induced soil moisture stress upon *Fomes annosus* root rot of loblolly pine. Phytopathology 58 (3):269–272. BA49:69698.

1181. Trappe, J.M.

1962. Fungus associates of ectotrophic mycorrhizae. Bot. Rev. 28(4):538–606.

1182. True, R.P., T.M. Judy, and E. Ross.

1955. The intake of solutions by tops of freshly cut oak stumps. Phytopathology 45(8):466. FA17:1626.

1183. Tubbs, C.H.

1963. Root development of yellow birch in humus and a sandy loam. USDA For. Serv. Res. Note LS-33. FA25:3318.

1184. Tubbs, C.H.

1973. Allelopathic relationship between yellow birch and sugar maple seedlings. For. Sci. 19(2):139–145. BA56:66322.

1185. Tubbs, C.H.

1976. Effect of sugar maple root exudate on seedlings of northern conifer species. USDA For. Serv. Res. Note NC-213.

1186. Tubbs, C.H.

1977. Root-crown relations of young sugar maple and yellow birch. USDA For. Serv. Res. Note NC-225.

1187. Turner, L.M.

1936. Root growth of seedlings of *Pinus echinata* and *Pinus taeda*. J. Agric. Res. 53(2):145–149. BA11:16898.

1188. Turner, L.M.

1936. A comparison of roots of southern shortleaf pine in three soils. Ecology 17(4):649–658. BA11:14604.

1189. Turner, P.D.

1962. Morphological influence of exudates of mycorrhizal and non-mycorrhizal fungi on excised root cultures of *Pinus sylvestris* L. Nature 194(4828):551–552. BA40:3445.

1190. United Kingdom Forestry Commission.

1971. Select annotated bibliography on tree root systems, including damage by tree roots. London, UK. For. Comm. Libr. FA34:4460.

1191. U.S. Forest Service.

1908. Tamarack (*Larix laricina* (DuRoi) Koch). USDA For. Serv. Silvical Leaflet 32.

1192. U.S. Forest Service.

1909. Pignut hickory: *Hicoria glabra* (Mill.) Britton. USDA For. Serv. Silvical Leaflet 48.

1193. U.S. Forest Service.

1935. Prolonged exposure of roots causes death of planting stock. USDA For. Serv. Tech. Note 96, Lake States For. Exp. Stn.

1194. U.S. Forest Service.

1951. Protection from fungi: The littleleaf disease of pine. USDA For. Serv. Southeast. For. Exp. Stn. Rep. 1949/50:11-14. FA13:1411.

1195. U.S. Forest Service.

1959. Root grafts in oak. USDA For. Serv. Central States For. Exp. Stn. Annu. Rep. 1958:6–7. FA20:4791.

1196. U.S. Forest Service.

1960. Hybrid poplar growth rates correlated with root anatomy. USDA For. Serv. Northeast. For. Exp. Stn. Annu. Rep. 1959:18. FA21:4069.

1197. U.S. Forest Service.

1960. Root grafts in oaks. USDA For. Ser. Central States For. Exp. Stn. Annu. Rep. 1959:2–3. FA21:4584.

1198. U.S. Forest Service.

1960. Root rots. USDA For. Serv. Northeast. For. Exp. Stn. Annu. Rep. 1959:45–48. FA21:4571.

1199. U.S. Forest Service.

1964. Backflash in sweetgum. USDA For. Serv. Northeast. For. Exp. Stn. Annu. Rep. 1963:22–23. FA26:619.

1200. Urie, D.H.

1966. An analysis of the hydrologic budget in glacial sands under pine and hardwood forests. Diss. Abstr. 26(8):4149-4150. FA28:74.

1201. Ursic, S.J.

1963. Modifications of planting technique not recommended for loblolly on eroded soils. Tree Plant. Notes 57, p. 13-17. FA24:3596.

1202. Ursino, D.J., and G. Krotkov.

1968. The effect of water stress on the translocation of recent photosynthate to the roots of young plants of *Pinus strobus*. Can. J. Bot. 46(10):1327–1329. BA50:32962.

1203. Vaartaja, C

1957. Effect of *Trichoderma* on tree seedlings and on their pathogens. Bi-mon. Progr. Rep. Div. For. Biol., Dep. Agric. Can. 13(5):1. FA19:1926.

1204. Vaartaja, O., and W.H. Cram.

1956. Damping-off pathogens of conifers and of *Caragana* in Saskatchewan. Phytopathology 46(7):391–397. FA18:367.

1205. Vaartaja, O., and P.J. Salisbury.

1961. Potential pathogenicity of *Pythium* isolates from forest nurseries. Phytopathology 51(8):505-507.

1206. Vaartaja, O., W.H. Cram, and G.A. Morgan.
1961. Damping-off etiology especially in forest nurseries. Phytopathology 51(1):35–42.

1207. Van Eck, W.A.
1959. Site and root studies of red pine (*Pinus resinosa* Ait.) plantations in Lower Michigan. Ph.D. thesis, Mich. State Coll. BA35:17177.

1208. Van Groenewood, H.
1956. A root disease complex in Saskatchewan white spruce. For. Chron. 32(1):11–13.

1209. Van Wormer, H.M.
1937. Effect and treatment of girdling roots. Natl. Shade Tree
Conf. Proc. 13:30–36.

1210. Van Wormer, H.M.
1951. Basic tree ailments augmented by girdling the roots. Arborist's News 16(1):2–6. BA25:25246.

1211. Vatkovskii, O.S., T.L. Bystritskaya, and L.A. Grishina.
1971. Biomassa kornevykh sistem eli, duba i gledichii v europeiskoi chasti SSSR. [Biomass of the root systems of the spruce, oak, and honeylocust in the European part of the USSR.] Biol. Nauk. 14(9):108–111. BA54:18883.

Vecchierello, H.
1928. A study of the origin and development of the radicle histogens of *Quercus prinus* L. Cathol. Univ. Am. Contrib. 8.

1213. Vimmerstedt, J.P.
1968. Root cation-exchange capacity and the mineral nutrition of eastern white pine and eastern redcedar. Proc. Soil Sci. Soc. Am. 32(2):289–292. BA49:102122.

1214. Vogt, A.R.
1970. Effect of gibberellic acid on germination and initial seedling growth of northern red oak. For. Sci. 16(4):453-459.
BA52:36839.

1215. Voigt, G.K.
1953. The effects of fungicides, insecticides, herbicides, and fertilizer salts on the respiration of root tips of tree seedlings. Proc. Soil Sci. Soc. Am. 17(2):150–152. BA28:11944.

1216. Voight, G.K.
1955. The effect of applied fungicides, herbicides and insecticides on the content of nutrient elements in tissue of coniferous seedlings. Proc. Soil Sci. Soc. Am. 19(2):237–239. FA17:502.

1217. Voigt, G.K.
1966. Phosphorus uptake in young pitch pine (*Pinus rigida* Mill.).
Proc. Soil Sci. Soc. Am. 30(3):403–406.

1218. Voigt, G.K.
1970. Ion source and ion uptake by pine seedlings. *In* Tree growth and forest soils. C.T. Youngberg, and C.B. Davey, eds., 3rd North Am. For. Soils Conf. Proc., N.C. State Univ., Raleigh, [August 1968], Oreg. State Univ. Press, Corvallis, p. 181–191.

1219. Voight, G.K., anf G.L. Steucek.
1969. Nitrogen distribution and accretion in an alder ecosystem.
Proc. Soil Sci. Soc. Am. 33(6):946–949.

1220. Voigt, G.K., B.N. Richards, and E.C. Mannion.
1964. Nutrient utilization by young pitch pine. Proc. Soil Sci. Soc.
Am. 28(5):707-709. FA26:3425.

1221. Vozzo, J.A.
1969. Endotrophic mycorrhizae found on *Populus deltoides*. For. Sci. 15(2):158. FA31:379.

1222. Vozzo, J.A., and E. Hacskaylo.
1964. Anatomy of mycorrhizae of selected eastern forest trees.
Bull. Torrey Bot. Club 91(5):378-387. FA26:3434.

1223. Vozzo, J.A., and E. Hacskaylo.
1974. Endo- and ectomycorrhizal associations in five *Populus* species. Bull. Torrey Bot. Club 101(4):182–186. FA36:3831.

1224. Wagg, J.W.B.1964. White spruce regeneration on the Peace and Slave River lowlands. Can. Dep. For. Publ. 1069.

Wagg, J.W.B.1967. Origin and development of white spruce root-forms. Can.Dep. For. Publ. 1192. BA49:64706.

1226. Wagner, K., and H. Kurz.
1954. Cypress: Root and stem modifications in relation to water.
Fla. State Univ. Stud. 13(2):18-47.

1227. Wahlenberg, W.G.1960. Loblolly pine: Its use, ecology, regeneration, protection, growth and management. Duke Univ. Sch. For.

1228. Wakeley, P.C.
1953. Progress in the study of pine races. South. Lumberman 187(2345):137–140.

1229. Walker, L.C.1962. The effects of water and fertilizer on loblolly and slash pine seedlings. Proc. Soil Sci. Am. 26(2):197-200.

1230. Walker, L.C.
1967. Effects of water level and fertilizer combinations on loblolly and slash pine seedlings. Tree Plant. Notes 18(1):10-12. FA28:5234.

1231. Walker, L.C., R.L. Green, and J.M. Daniels.
1961. Flooding and drainage effects on slash pine and loblolly pine seedlings. For. Sci. 7(1):2–15. FA22:4224.

1232. Wallis, G.W.
1959. Effect of biocides on the growth of seedlings of monterey pine, black locust, and European alder, and the development of their symbiotrophic organs. Univ. Wis. For. Res. Notes 45.

1233. Wallis, G.W.
1961. Infection of Scots pine roots by *Fomes annosus*. Can. J. Bot. 39(1):109–121. BA36:28659.

1234. Walstad, J.D., J.B. Hart, Jr., and S.C. Cade. 1973. Carbofuran-clay root dip protects loblolly pine seedlings from debarking weevils. J. Econ. Entomol. 66(5):1219–1220. FA35:2346.

1235. Ward, J.C., J.E. Kuntz, and E.M. McCoy.
1969. Bacteria associated with 'shake' in broadleaf trees. (Abstr.)
Phytopathology 59(8):1056. FA31:2763.

1236. Ware, G.H., and W.T. Penfound.
1949. The vegetation of the lower levels of the floodplain of the South Canadian River in central Oklahoma. Ecology 30(4):478–484. FA11:1673.

1237. Wargo, P.M.
1971. Enhanced growth of *Armillaria mellea* on extracts from roots of defoliated sugar maple trees. (Abstr.) Phytopathology 61(8):915. FA33:2772.

1238. Wargo, P.M.
1971. Seasonal changes in carbohydrate levels in roots of sugar
maple. USDA For. Serv. Res. Pap. NE-213. BA55:4731.

1239. Wargo, P.M.
1972. Defoliation-induced chemical changes in sugar maple roots stimulate growth of *Armillaria mellea*. Phytopathology 62(11):1278-1283. BA55:51303.

1240. Wargo, P.M.
1974. Lysis of the cell wall of *Armillaria mellea* by enzymes from forest trees. (Abstr.) Phytopathology 64(5):588. FA35:7693.

1241. Wargo, P.M.1975. Estimating starch content in roots of deciduous trees—a visual technique. USDA For. Serv. Res. Pap. NE-313. FA37:797.

1242. Wargo, P.M.
1976. Variation of starch content among and within roots of red and white oak trees. For. Sci. 22(4):468–472.

1243. Wargo, P.M., J. Parker, and D.R. Houston.
1972. Starch content in roots of defoliated sugar maple. For. Sci.
18(3):203–204. BA55:45794.

1244. Warren, G.L.
1956. Root injury to conifers in Canada by species of *Hylobius* and *Hypomolyx* (Coleoptera: Curculionidae). For. Chron. 32(1):7–10.

1245. Warren, G.L.

1956. The effect of some site factors on the abundance of *Hypomolyx piceus* (Coleoptera: Curculionidae). Ecology 37(1):132–139.

1246. Warren, G.L., and P. Singh.
1970. *Hylobius* weevils and *Armillaria* root rot in a coniferous plantation in Newfoundland. Bi-mon. Res. Notes 26(6):55.

FA32:6471.

1247. Warren, G.L., and R.D. Whitney.

1951. Spruce root borer (*Hypomolyx* sp.), root wounds, and root diseases of white spruce. Bi-mon. Progr. Rep. Div. For. Biol., Dep. Agric. Can. 7(4):2–3. FA13:1386.

1248. Wassink, E.C., and S.D. Richardson.
1951. Observations on the connection between root growth and shoot illumination in first-year seedlings of *Acer pseudoplatanus* L. and *Quercus borealis maxima* (Marsh) Ashe. Proc. K. Ned. Akad. Wet. Ser. C 54(5):503–510. FA13:3682.

1249. Watt, A.S., and G.K. Fraser.
1933. Tree roots and the field layer. J. Ecol. 21(2):404–414.
BA10:128.

1250. Watts, J.G., and J.B. Hatcher.
1954. White grub damage to young pine plantations. J. Econ.
Entomol. 47(4):710–711. FA16:2009.

1251. Wean, R.E.
1937. The parasitism of *Polyporus schweinitzii* on seedling *Pinus strobus*. Phytopathology 27(12):1124–1142.

1252. Weaver, J.E.
1925. Investigations on the root habits of plants. Am. J. Bot.
12:502–509.

1253. Weaver, J.E., and J. Kramer.
1932. Root system of *Quercus macrocarpa* in relation to the invasion of prairie. Bot. Gaz. 94(1):51–85. BA7:17934.

1254. Webb, D.P. 1976. Effects of cold storage duration on bud dormancy and root regeneration of white ash (*Fraxinus americana* L.) seedlings. Hort-Science 11(2):155–157.

1255. Webb, D.P. 1976. Root growth in *Acer saccharum* Marsh. seedlings: Effects of light intensity and photoperiod on root elongation rates. Bot. Gaz. 137(3):211–217.

1256. Webb, D.P. 1977. Root regeneration and bud dormancy of sugar maple, silver maple, and white ash seedlings: Effects of chilling. For. Sci. 23(4):474–483.

1257. Weidensaul, T.C., and J.R. McClenahen.
1977. Influences of rooting media on containerized tree growth.
Ohio Rep. Res. and Devel. 62(6):83–86. Ohio Agric. Res. and Devel. Cent. Wooster.

1258. Wells, C.G., J.R. Jorgensen, and C.E. Burnette. 1975. Biomass and mineral elements in a thinned loblolly pine plantation at age 16. USDA For. Serv. Res. Pap. SE-126. FA37:6932.

1259. Wenger, K.F.
1952. Effect of moisture supply and soil texture on the growth of sweetgum and pine seedlings. J. For. 50(11):862-864. FA14:2027.

1260. Wenger, K.F. 1955. Light and mycorrhiza development. Ecology 36(3):518– 520. FA17:228.

1261. Werner, R.A.
1973. Absorption, translocation, and metabolism of root-absorbed

14C-Monitor in loblolly pine seedlings. J. Econ. Entomol.
66(4):867–872. FA35:1751.

1262. Werner, R.A.
1974. Distribution and toxicity of root-absorbed ¹⁴C-Orthene and its metabolites in loblolly pine seedlings. J. Econ. Entomol. 67(5):588–591. FA36:4907.

1263. Werner, R.A.
1975. Bioactivity of Orthene in loblolly pine seedlings. J. Ga.
Entomol. Soc. 10(2):156–162. FA37:430.

1264. Westing, A.H.
1959. A curious example of natural layering in maple. Turtox News
37(2):68. BA33:30926.

1265. Wheeler, C.T.
1971. The causation of the diurnal changes in nitrogen fixation in the nodules of *Alnus glutinosa*. New Phytol. 70(3):487–495. FA33:2205.

1266. White, D.P., and B.G. Ellis.
1965. Nature and action of slow release fertilizers as nutrient sources for forest tree seedlings. Mich. Agric. Exp. Stn. Q. Bull. 47(4):606-614.

White, D.P., and R.S. Wood.
1958. Growth variations in a red pine plantation influenced by a deep-lying fine soil layer. Proc. Soil Sci. Soc. Am. 22:174–177.

1268. White, D.P., G. Schneider, and W. Lemmien.
1975. Regeneration and growth of quality hardwoods on modified sites. *In* Forest soils and forest land management. B. Bernier and C.H. Winget, eds., 4th North Am. For. Soils Conf. Proc., Laval Univ., 1973. Press l'Univ. Laval, Québec. p. 397–415.

1269. White, L.T.
1951. Studies of Canadian Thelephoraceae. VIII. Corticum galactinum (Fr.) Burt. Can. J. Bot. 29(3):279–296. FA13:1331.

1270. Whitford, L.A.
1956. A theory on the formation of cypress knees. J. Elisha Mitchell Sci. Soc. 72(1):80–83. BA31:2269.

1271. Whitney, R.D.
1952. Relationship between entry of root-rotting fungi and root-wounding by *Hypomolyx* and other factors in white spruce. Bimon. Progr. Rep. Div. For. Biol. Dep. Agric. Can. 8(1):2–3.

1272. Whitney, R.D.1952. Root disease of balsam fir. Bi-mon. Progr. Rep. Div. For. Biol. Dep. Agric. Can. 8(5):2.

1273. Whitney, R.D.
1961. Root wounds and associated root rots of white spruce. For.
Chron. 37(4):401–411. BA38:11599.

Whitney, R.D.
1962. Studies in forest pathology. XXIV. *Polyporus tomentosus*Fr. as a major factor in stand-opening disease of white spruce. Can.
J. Bot. 40(12):1631–1658. FA24:5180.

1275. Whitney, R.D. 1963a. Artificial infection of small spruce roots with *Polyporus tomentosus*. Phytopathology 53(4):441–443. BA44:8242.

Whitney, R.D.
1963b. White birch mortality at Candle Lake, Saskatchewan. Bimon. Progr. Rep., For. Entom. Path. Branch, Dep. For. Can. 19(1):3. BA45:26215.

1277. Whitney, R.D.
1965. Mycorrhiza-infection trials with *Polyporus tomentosus* on spruce and pine. For. Sci. 11(3):265–270. FA27:3552.

1278. Whitney, R.D.
1966. Germination and inoculation tests with basidiospores of *Polyporus tomentosus*. Can. J. Bot. 44(10):1333-1343.
BA48:51654.

1279. Whitney, R.D.
1967. Comparative susceptibility of large and small spruce roots to
Polyporus tomentosus. Can. J. Bot. 45(11):2227-2229.
BA49:31630.

1280. Whitney, R.D.

1972a. Mortality of spruce in Ontario caused by *Polyporus tomentosus* root rot. Bi-mon. Res. Notes 28(6):39–40.

1281. Whitney, R.D.

1972b. Root rot in white spruce planted in areas formerly heavily attacked by *Polyporus tomentosus* in Saskatchewan. Bi-mon. Res. Notes 28(4):24. FA34:1731.

1282. Whitney, R.D.
1977. Polyporus tomentosus root rot of conifers. Fish. and Environ. Can. For. Serv. For. Tech. Rep. 18, Great Lakes For. Res. Centre.

1283. Whitney, R.D., W.P. Bohaychuk, and M.A. Briant.
1972. Mycorrhizae of jack pine seedlings in Saskatchewan and Manitoba. Can. J. For. Res. 2(3):228–235. FA34:5048.

1284. Whittaker, R.H.
1961. Estimation of net primary production of forest and shrub communities. Ecology 42(1):177–180.

1285. Whittaker, R.H., and G.M. Woodwell.
1968. Dimension and production relations of trees and shrubs in the
Brookhaven Forest, N.Y. J. Ecol. 56:1–25.

1286. Wiant, H.V., Jr.
1966. The concentration of carbon dioxide near the ground under eastern white pine and black spruce trees. Adv. Front. Plant Sci. 16:197–203. FA29:3297.

1287. Wiant, H.V., Jr., and M.A. Ramirez.
1974. Don't plant white pine near walnut! Tree Plant. Notes 25(4):30.

1288. Wichman, J.R.
1973. Uptake, distribution, and degradation of simazine by black walnut and yellow poplar seedlings. Diss. Abstr. Int. 34B(1):13–14. FA36:937.

1289. Wight, W.
1933. Radial growth of the xylem and the starch reserves of *Pinus sylvestris*. A preliminary survey. New Phytol. 32(2):77–96. BA8:6956.

1290. Wilcox, H.E.
1962. Morphological features and growth behavior of the root of red pine, *Pinus resinosa* Ait. (Abstr.) Am. J. Bot. 49(6, part 2):658. BA41:3388.

1291. Wilcox, H.E.
1964. Xylem in roots of *Pinus resinosa* Ait. in relation to heterorhizy and growth activity. *In* The formation of wood in forest trees. A symposium. M.H. Zimmermann, ed., Academic Press, Inc., N.Y. p. 459–478. BA46:49380.

1292. Wilcox, H.E. 1967. Seasonal patterns of root initiation and mycorrhizal development in *Pinus resinosa* Ait. Proc. 14th Int. Union For. Res. Organ. Congr. [Munich 1967] Part V, Section 24, p. 29–39. FA29:2003.

1293. Wilcox, H.E.
1968. Morphological studies of the root of red pine, *Pinus resinosa*. I. Growth characteristics and patterns of branching. Am. J. Bot. 55(2):247–254. BA49:85785.

1294. Wilcox, H.E.
1968. Morphological studies of the roots of red pine, *Pinus resinosa*. II. Fungal colonization of roots and the development of mycorrhizae. Am. J. Bot. 55(6, part 1):688–700. BA49:112949.

1295. Wilcox, H.E.
1971. Morphology of ectendomycorrhizae in *Pinus resinosa*. *In*Mycorrhizae. E. Hacskaylo, ed., First North Am. Conf. on Mycorrhizae, Univ. Ill. [Urbana, April 1969] USDA For. Serv. Misc. Publ., p. 54–68.

1296. Wilcox, H.E., and R. Ganmore-Neumann.
1974. Ectendomycorrhizae in *Pinus resinosa* seedlings. I. Characteristics of mycorrhizae produced by a black imperfect fungus.
Can. J. Bot. 52(10):2145–2155. FA36:4537.

1297. Wilcox, H.E., and R. Ganmore-Neumann.
1975. Effects of temperature on root morphology and ectendomy-corrhizal development in *Pinus resinosa* Ait. Can. J. For. Res. 5(2):171–175. BA60:48874.

1298. Wilcox, H.E., R. Ganmore-Neumann, and C.J.K. Wang. 1974. Characteristics of two fungi producing ectendomycorrhizae in *Pinus resinosa*. Can. J. Bot. 52(11):2279–2282. FA36:5500.

1299. Wilcox, J.R., and R.E. Farmer, Jr.
1968. Heritability and *C* effects in early root growth of eastern cottonwood cuttings. Heredity 23(2):239–245. BA49:19477.

1300. Wilde, S.A.1946. Forest soils and forest growth. Chron. Bot. Co., Waltham, Mass.

1301. Wilde, S.A., and R.O. Rosendahl.1945. Value of potassium feldspar as a fertilizer in forest nurseries.J. For. 43(5):366-367.

1302. Wilde, S.A., and G.K. Voigt.
1949. Absorption-transpiration quotient of nursery stock. J. For.
47(8):643–645. FA11:865.

1303. Williams, J.R.M.1958. Root development and height growth of red pine in relation to soil conditions. M.Sc.F. thesis, Univ. Toronto.

Williams, R.D.1972. Root fibrosity proves insignificant in survival, growth of black walnut seedlings. Tree Plant. Notes 23(2):22-25.

1305. Williamson, A.W.1913. Cottonwood in the Mississippi Valley. USDA Bull. 24.

1306. Williston, H.L.
1962. Pine planting in a water impoundment area. USDA For. Serv. South. For. Note 137, South. For. Exp. Stn. FA24:5098.

1307. Wilner, J., and E.J. Brach.
1974. Hardiness of roots in relation to shoots of container-grown plants by an electric method. Can. J. Plant Sci. 54(2):281–289. FA36:6118.

1308. Wilson, B.F.
1964. Structure and growth of woody roots of *Acer rubrum* L.
Harvard For. Pap. 11. BA47:9524.

1309. Wilson, B.F.1967. Root growth around barriers. Bot. Gaz. 128(2):79–82.BA48:123727.

Wilson, B.F.1970. Evidence for injury as a cause of tree root branching. Can. J.Bot. 48(8):1497–1498. FA32:2217.

1311. Wilson, B.F.1971. Vertical orientation of red maple (*Acer rubrum L.*) roots.Can. J. For. Res. 1(3):147–150. BA53:62982.

1312. Wilson, B.F.
1975. Distribution of secondary thickening in tree root systems. *In*The development and function of roots. 3rd Cabot symposium.
J.G. Torrey and D.T. Clarkson, eds., Academic Press, London, p.
197–219.

1313. Wilson, B.F., and S.B. Horsley.
1970. Ontogenetic analysis of tree roots in *Acer rubrum* and *Betula papyrifera*. Am. J. Bot. 57(2):161–164. BA51:80539.

1314. Wilson, L.F.1977. A guide to insect injury of conifers in the Lake States. USDAFor. Serv. Agric. Handbk. 501.

1315. Wingfield, E.B. 1969. Mycotrophy in loblolly pine. I. The role of *Pisolithus tinctorius* and *Rhizoctonia solani* in survival of seedlings. II. Mycorrhiza formation after fungicide treatment. Diss. Abstr. 29B(8):2707. FA31:373.

1316. Winston, D.A., and G. Schneider. 1977. Conifer establishment by hand seeding on sites prepared with the Bräckekultivatorn. Can. For. Serv. Pap. O-X-255, Great Lakes For. Res. Centre. 1317. Wong, T.L., R.W. Harris, and R.E. Fissell.
1971. Influence of high soil temperatures on five woody-plant species. J. Am. Soc. Hort. Sci. 96(1):80–83. FA33:2495.

1318. Wood, O.M.

1934. The root system of a chestnut oak. (*Quercus montana* Willd.). Natl. Shade Tree Conf. Proc. 10:95–99. BA9:14506.

1319. Wood, O.M.

1939. Relation of the root system of a sprouting stump in *Quercus montana* Willd. to that of an undisturbed tree. J. For. 37(4):309–312. BA14:5587.

1320. Woodroof, J.G.

1933. Relation of the root system of pecan trees to nursery and orchard practices. Ga. Agric. Exp. Stn. Bull. 176. BA10:14173.

1321. Woodroof, J.G., and N.C. Woodroof.
1934. Pecan root growth and development. J. Agric. Res.
49(6):511-530. BA9:19480.

1322. Woodroof, N.C.

1933. Pecan mycorrhizas. Ga. Agric. Exp. Stn. Bull. 178.

1323. Woods, F.W.

1960. Biological antagonisms due to phytotoxic root exudates. Bot. Rev. 26(4):546–569. FA22:2874.

1324. Woods, F.W.

1962. Root exudates of higher plants: Ecological considerations. Proc. 13th Int. Union For. Res. Organ. Congr. [Vienna 1961] Part 2(1), Section 21–4/4. FA24:124.

1325. Woods, F.W.

1970. Interspecific transfer of inorganic materials by root systems of woody plants. J. Appl. Ecol. 7(3):481–486. BA52:48136.

1326. Woods, F.W., and K. Brock. 1964. Interspecific transfer of Ca⁴⁵ and P³² by root systems. Ecology 45(4):886–889. BA46:55571.

1327. Woods, F.W., and R. Martin.

1973. Loblolly pine mycorrhizae in east Tennessee. Tree Plant. Notes 24(4):27. FA36:5503.

1328. Woods, F.W., and D. O'Neal.

1965. Tritiated water as a tool for ecological field studies. Science 147(3654):148–149.

1329. Woodwell, G.M., R.H. Whittaker, and R.A. Houghton. 1975. Nutrient concentrations in plants in the Brookhaven oak-pine forest. Ecology 56(2):318–332. FA37:1483.

1330. Woolsey, T.S., Jr., and H.H. Chapman.

1914. Norway pine in the Lake States. USDA Bull. 139.

1331. Worley, J.F., and E. Hacskaylo.

1959. The effect of available soil moisture on the mycorrhizal association of Virginia pine. For. Sci. 5(3):267–268. FA21:1549.

1332. Yeager, A.F.

1935. Root systems of certain trees and shrubs grown on prairie soils. J. Agric. Res. 51(12):1085–1092. BA10:20447.

1333. Yeager, L.E.

1949. Effect of permanent flooding in a river-bottom timber area. Ill. Nat. Hist. Surv. Bull. 25(2). FA12:1543.

1334. Yeatman, C.W.

1955. Tree root development on upland heaths. (G.B.) For. Comm. Bull. 21. BA31:12058.

1335. Yelenosky, G.

1964. Tolerance of trees to deficiencies of soil aeration. Int. Shade Tree Conf. Proc. 40:127–147.

1336. Yelenosky, G., and C.L. Fergus.

1959. Absorption and translocation of solutions by healthy and wilt-diseased red oaks. Pa. Agric. Exp. Stn. Bull. 657. FA23:787.

1337. Yim, K.B.

1962. Physiological studies on rooting of pitch pine (*Pinus rigida* Mill.) cuttings. Res. Rep. Inst. For. Genet., Suwon. 2, p. 20–56. FA24:1967.

1338. York, H.H.

1941. Relative susceptibility of young pine trees in artificial and natural stands to various root, root-crown, and butt fungus parasites. (Abstr.) Phytopathology 31:25.

1339. York, H.H., R.E. Wean, and T.W. Childs.

1936. Some results of investigations on *Polyporus schweinitzii* Fr. Science 84:160–161.

1340. Yorke, J.S., and F.C. Pollett.

1969. Root systems of tamarack growing on peatland. Bi-mon. Res. Notes 25(1):2. FA30:5400.

1341. Yotov, T., I. Palashev. V. Nikolov, and Y. Lyapova.
1975. [Planting depth for poplars on sandy strips on the Black Sea coast in south Bulgaria.] Gorskostop. Nauka 12(2):23–33. [In. Bulg. with Russ. and Engl. summ.] FA36:7668.

1342. Young, H.E.

1947. Carbohydrate absorption by roots of *Pinus taeda*. Queensland J. Agric. Sci. 4(1/2):1-6. FA11:2513.

1343. Young, Harold E., L. Strand, and R. Altenberger. 1964. Preliminary fresh and dry weight tables for seven tree species in Maine. Maine Agric. Exp. Stn. Tech. Bull. 12.

1344. Youngman, A.L.

1967. An ecotypic differentiation approach to the study of isolated populations of *Pinus taeda* in south central Texas. Diss. Abstr. 27B(9):3006. FA29:173.

1345. Yount, W.L.

1955. Longevity of the oak wilt fungus in oak roots as related to spread through root grafts. Plant Dis. Rep. 39(3):256–257. FA16:4247.

1346. Yount, W.L.

1958. Results of root inoculations with the oak wilt fungus in Pennsylvania. Plant Dis. Rep. 42(4):548–551. BA33:19520.

1347. Young, W.L., and J.R. Bloom.

1963. Nematodes in nursery plants in Pennsylvania. Plant Dis. Rep. 47(5):405–407. FA25:5415.

1348. Zahner, R.

1956. Root development of thinned pines. USDA For. Serv. South. For. Notes 101, p. 1–2. South. For. Exp. Stn. FA17:3717.

1349. Zahner, R., and N.V. DeByle.

1965. Effect of pruning the parent root on growth of aspen suckers. Ecology 46(3):373–375. FA27:448.

1350. Zahner, R., and F.W. Whitmore.

1960. Early growth of radically thinned loblolly pine. J. For. 58(8):628–634.

1351. Zak, B.

1955. Inheritance of resistance to littleleaf in shortleaf pine. USDA For. Serv. Res. Note 88, Southeast. For. Exp. Stn. FA17:1238.

1352. Zak, B.

1956. Experimental air-layering of shortleaf and loblolly pine. USDA For. Serv. Stn. Pap. 69, Southeast. For. Exp. Stn.

1353. Zak, B

1957. Resistance to littleleaf in shortleaf pine. 4th Conf. For. Tree Improv. Proc., Athens, Ga. [8-9 January 1957] p. 41-43.

1354. Zak, B.

1961. Aeration and other soil factors affecting southern pines as related to littleleaf disease. USDA Tech. Bull. 1248. FA23:3294.

1355. Zak, B.

1964. Role of mycorrhizae in root disease. Annu. Rev. Phytopathol. 2:377–392.

1356. Zak, B., and W.A. Campbell.

1958. Susceptibility of southern pines and other species to the littleleaf pathogen in liquid culture. For. Sci. 4(2):156–161. FA20:723.

1357. Zalasky, H.

1974. Keissleriella emergens, a perfect state of Diplodia tumefaciens in roots of poplar. Can. J. Bot. 52(1):11-13. BA58:6478.

1358. Ziegler, H.

1958. Die Bedeutung der Knöllchenbakterien für den Stickstoffhaushalt der Robinie (*Robinia pseudoacacia* L.) [The importance of root nodules in the N economy of *R. pseudoacacia*.] Arch. Forstw. 7(4/5):352–369. [Engl. and Russ. summ.] FA20:1556.

- 1359. Zinkan, C.G., J.K. Jeglum, and D.E. Harvey. 1974. Oxygen in water culture influences growth and nutrient uptake of jack pine, black spruce and white spruce seedlings. Can. J. Plant Sci. 54(3):553-558. BA59:8188.
- 1360. Zon, R.1905. Loblolly pine in eastern Texas, with special reference to the production of cross-ties. USDA For. Serv. Bull. 64.

Appendix

Key

| | Subject |
|------------|--|
| IA,B | Root form and relation to vegetative growth |
| IC | Genetic variation in root growth and development |
| ID | Periodic and seasonal aspects of root growth and development |
| IE-1 | Influences of moisture, temperature, and soil on root growth and development |
| IE-1a | Influence of moisture extremes on root growth and development |
| IE-2 | Influence of the shoot on root growth and development |
| IIA | Uptake and upward translocation |
| IIB-1 | Translocation downward—carbohydrate reservation |
| IIB-2 | Translocation downward—exudation |
| IIC | Root respiration |
| IIIA-1 | Competition with woody and herbaceous plants |
| IIIA-2 | Root grafting |
| IIIA-3 | Effects of roots on soil |
| IIIB-1 | Symbiotic relations with other than woody and herbaceous plants |
| IIIB-2a | Diseases of mechanical or undetermined cause |
| IIIB-2bi | Insect disease |
| IIIB–2bii | Nematode disease |
| IIIB–2biii | Fungus disease |
| IIIB-2biv | Bacteria disease |
| IV | Root response to nursery and forestry practices |

See page iii of the introduction for instructions on the use of the key and the subject index.

Subject Index

| | | | | | | | | | | | Subject | | | | | | | | |
|---------------------|--|------|---|---|-----------|---------------------------------------|-----|------|-----------|-----|----------------------------------|------------|-------|--------------------|-----------------------------------|--------------|---------------|---|--------------|
| Species | IA,B | IC I | | IE– 1 | IE– 1a | IE- 2 | IIA | IIB- | IIB- 2 | IIC | IIIA- | IIIA- 2 | IIIA- | IIIB- | IIIB– 2a | IIIB- 2bi | IIIB- 2bii | IIIB- 2biii | IV |
| Abies spp. | | | | | | 253 | | | | | 150 | | | 554 1181 | | | | | |
| Abies balsamea | 859 559 67 382 68 71 1052 1343 86 978 | | 68 | 349 12 859 559 1052 | | 747 740 746 253 939 71 | 20 | |) | | 690 446 659 658 | 86 978 | | 859 783 1181 | 1123 788 871 923 1126 | 1121 | 792 | 1122 1269 1272 310 1096 32 579 480 1087 1116 695 1085 696 | 11 |
| Acer spp. | 830 | | | | | 1210 | 852 | | | | 150 | | 601 | 1181 | 1318 1209 1141 | 448 | 792 | 207 | 1209 1210 |
| Acer negundo | 128 1332 830 211 465 509 935 238 | | | 1332 128 481 566 568 174 104 755 | | 128 935 15 | 568 | | | | 128 446 15 238 | | | 809 | 871 | | 792 | 996 | |
| Acer nigrum | 830 | | | | | | | | | | | | | | | | | | |
| Acer pennsylvanicum | 830 | | 72 | | | | | | | | | | | | | | | • | |
| Acer rubrum | 856 781 763 1178 349 1310 1313 762 830 1130 826 934 828 827 829 114 1200 1309 1343 1308 1135 336 382 | 1. | 72 326:891 891 892 890:763 313 | 1311 828 114 1309 336 | | 1226 762 605 763 | | | | | 762 1135 446 658 659 | | | 114 | | | | 207 | 934 963 |

| | | | | | | | | | | | Subject | | | | | | | | | |
|-------------------|--|----|-----------------------------------|---|-------------------------|---|------------|------------------------------------|------|------|---|------------|--------------|---|-----------------------------------|--------------|-----------------|----------------------|---------------|-----------------------------------|
| Species | IA,B | IC | ID | IE– 1 | IE- 1a | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- | IIIA- 2 | - IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | - IIIB- 2bii | | IIIB- 2biv | IV |
| Acer saccharinum | 1332 830 983 989 238 781 | | 1256 780 982 988 1254 | | | 1333 423 988 989 743 747 1254 | 568 | | | 1177 | 238 446 710 15 | | | 779 | | | | 3% | 2 | 1256 |
| Acer saccharum | 908 372 128 1186 1184 830 1101 375 1135 382 1056 13 211 559 934 859 781 608 | | | 372 872 128 859 174 909 908 372 872 867 481 1161 1254 | | 372 371 416 909 743 747 1255 334 | 467 891 | 910 1243 912 1241 1238 | | | 372 1135 13 446 690 658 659 | | 1184 1185 | | 871 131 1210 1209 839 | 123 | | 124(123° 1239 | 7 | 416 608 1241 1246 334 |
| Aesculus spp. | 547 | | | | | | | | | | | | | | | | | | | |
| Aesculus glabra | 781 128 | | | 128 | | 128 | · | | | · | | | | | | | | | | |
| Aesculus octandra | | | | | 481 174 | | | | | | | | | | | | | | | |
| Alnus spp. | 106 | | | 135 | | | 1300 | | | | | | | 631 135 137 1181 106 138 139 | | | | | | |
| Alnus glutinosa | 791 117 | | 999 462 | 791 790 117 777 999 144 140 | 52 790 427 777 | 791 427 | 521 | 1265 | | | 790 | | | 777 790 791 141 144 1232 1265 141 96 142 139 1181 997 | | | | | | 1232 |

| | | | | | | | | | | 5 | Subject | | | | | | | | | |
|-------------------------|---|----|------------|---|---|-------------------|---------------------------|------|------|------------|------------|---------------------------------|-------|---------------------------------------|---|--------------|---------------|---------------------|---------------|--------------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- 1 | IIIA- ² | IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | IIIB- 2bii | | IIIB- 2biv | IV |
| Alnus glutinosa (con't) | | | | | | | | | | | | | | 931 858 521 332 136 | | | | | | |
| Alnus rugosa | | | | | 778 481 482 | | | | | | | | | 997 1219 858 281 139 | 871 | | | | | |
| Alnus serrulata | | | | | 174 | | | | | | | | | 139 631 997 | | | | | | |
| Asimina triloba | 781 | | | | 482 | | | | | | | | | | | | | | | - |
| Betula spp. | 755 | | | 698 1300 | | | 698 1300 338 890 | | | | 755 | 150 | | 1181 807 | 871 1210 839 | | | 207 | 7 | |
| Betula alleghaniensis | 372 1178 934 1107 957 1186 762 486 1184 559 382 859 608 | | 1022 | 372 956 957 486 559 954 1176 1183 575 576 577 | 372 | 372 743 747 | 576 577 1022 72 | | 1103 | | 762 | 372 710 658 659 446 | 1184 | 372 953 956 859 33 622 | 372 1107 889 503 209 456 953 955 956 957 958 237 | | | 1269 1203 310 | } | 934 1176 608 |
| Betula lenta | 1135 | | | 116 | | | 226 72 | | | | | 1135 446 | | 1181 1049 807 810 | 116 | | | | | |
| Betula nigra | | | 163 328 | 163 | 481 455 1333 778 482 174 | 328 | | | | 163 724 | | 446 | | 1181 810 | | | | | | |
| Betula papyrifera | 1107 1313 562 561 946 934 1343 752 1200 | | 1313 | 956 937 752 585 516 1268 | 12 | 743 747 562 | | | | | 86 | 561 690 658 659 446 | | 956 779 1181 | 1107 209 956 958 237 1276 1313 410 | | 1347 | 272 900 | | 934 796 1268 |

| | | | | | | | | | | | Subject | | | | | | | | | |
|----------------------|---|----|--------------|--------------|----------------------------------|-------------------|------------|--------------|---------------------------------------|----------------------|-----------------------|------------|---------|---|---------------------------------------|--------------|-----------------|---------|----------------------|----------------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- 1 | IIB- 2 | IIC | IIIA- 1 | IIIA- 2 | - IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | - IIIB- 2bii | | – IIIB ii 2biy | |
| Betula populifolia | 1178 762 | | | 1049 | | | 892 891 | | | | 762 | | | 1049 807 810 | 410 |) | 79: | 2 | | |
| Carpinus caroliniana | 781 | | | | 481 482 | | 72 | | | | | | | 1181 | | | | 12 | 69 | |
| Carya spp. | 134 394 1178 413 182 184 | | , | 1178 182 | | 134 733 605 | | | | | 184 | 15 | 50 | 1181 94 | 839 1210 | | | | | |
| Carya cordiformis | 781 1178 | | | 1178 | | 134 373 | 226 72 | | | | | | | *************************************** | | | | • | | |
| Carya glabra | 1192 547 1135 | | | 1192 | 482 | | 72 | 462 | , | | | 113 44 | | | | | | - | | |
| Carya illinoensis | 1178 1320 1321 | | 1320 1321 | 1320 1321 | | 1097 1320 | | 1098 1097 | • • • • • • • • • • • • • • • • • • • | alitaka di waliosh B | and the second second | 75 | 53 | 1320 1322 1321 1181 810 | | | 1320 514 | 0 5 | 14 | 1320 1321 1000 |
| Carya lacinosa | | | 780 | | | | | | | | | | | 780 1181 | | | | | | |
| Carya ovata | 781 544 128 211 375 | | | 544 128 | 481 482 174 104 | 128 134 | 72 | | | | , | | | 779 | | | | | | |
| Carya tomentosa | 547 | | | | 481 174 104 | | | . , . | | | | | | | , , , , , , , , , , , , , , , , , , , | | | | | |
| Castanea spp. | 547 | | | | | | | | | | | | | 631 | | | | 2 | 07 | |
| Castanea dentata | 1178 | | | 1178 | | | 72 | 1,0 | | | | | | 1181 | | | | 2 13 | 53 72 56 07 | |
| Catalpa spp. | 509 610 | | | | 509 | | | | | | | | | | | | | ***** | | 171 |
| Catalpa speciosa | 465 | | | | | 416 | | | | | | | | | | | | | | 416 |
| Celtis spp. | 547 509 | | | | 482 | | | | | | | 75 | 3 | | | | | | | |
| Celtis laevigata | | | | | 481 1333 568 566 174 | | | | | | | | | | | | | | | |

| | | | | | | | | | | | Subjec | t | | | | | | | | |
|------------------------|---------------------------------------|----|-------------------|--------------------------------|--|------------|-------------------|------|------|-----|--------|------------|---------|---------------------------------|-------------|--------------|---------------|---|---|------------|
| Species | IA,B | IC | ID | IE- 1 | IE- la | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- | IIIA- | - IIIA- | IIIB- | IIIB– 2a | IIIB- 2bi | IIIB– 2bii | | - IIIB- 2biv | IV |
| Celtis occidentalis | 781 1332 465 321 | | | 1332 | 455 565 568 566 174 104 | | 568 | | | | | | | 321 | | | 767 | 7 3 | 2 | 171 |
| Cercis canadensis | 781 782 | | | | 481 174 104 | | | | | | | | | 782 | 871 131 | | • | 135 | 6 | |
| CONIFER | 382 68 401 608 375 374 | | 439 462 991 | 90 301 401 375 991 | 263 | 374 991 | 1130 | 462 | | | | | | 324 521 94 | | | 3 | 23 15 71 116 10 108 91 53 86 1 | 8 8 2 1 9 4 7 6 8 | 839 608 |
| Cornus spp. | | | | | | 190 | 666 | | _ | | | | 601 | 631 | | 448 | 3 792 1347 | | | 190 |
| Cornus florida | 546 1178 660 184 | | | 1335 502 | 481 906 482 1335 174 | | 72 906 | | | | | | | 155 | 906 | | 603 | 8 | 9 | 502 |
| Cornus stolonifera | | | | 786 | 455 | | 786 | | | | | | | | 871 | | | | | |
| Corylus spp. | 228 | | | 228 | | 228 190 | | | | | | | | 1181 | | | | | | 190 |
| Crataegus spp. | | | | | 481 1333 482 174 104 | | | | | | | | | 1181 | 871 | | 792 | | | |
| Diospyros virginiana | 1049 | | | | 481 1333 482 174 104 | | | | | | | | | | 1049 | | | | | |
| Elaeagnus spp. | | | - | | | , , | | | • | | | | · | 137 138 | | | | = | | |
| Elaeagnus angustifolia | 509 1332 | | | 755 1332 | | | | | | | 755 | 5 | | 997 139 | 857 871 | | 797 | | | 171 |
| Fagus spp. | 524 490 | | | 524 1249 490 | | | 491 492 339 | | | | | 650 150 | | 324 490 491 492 221 | 839 | | | | | 650 |

| | | | | | | | | | | | Subject | | | | | | | | |
|------------------------|--|----|---------------------|----------------------------------|---------------------------|----------------------------------|-------------------|-------------|-----------|------|------------------|--------------------|-------------|----------------------------|---------------------------|--------------|---------------|----------------|--|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- 1 | IIB- 2 | IIC | IIIA- 1 | IIIA- | IIIA- | IIIB- 1 | IIIB- 2a | IIIB- 2bi | IIIB– 2bii | IIIB- 2biii | IV |
| Fagus spp. (con't) | | | | | | | | | | | | | | 94 339 | | | | | |
| Fagus grandifolia | 859 934 164 559 382 484 1135 | | | 859 1135 | 164 481 482 174 | 934 164 | 72 | | 1103 | | | 484 1135 446 | | 859 1181 1222 545 | 1209 484 | | 3 | 272 207 | 934 1019 |
| Fraxinus spp. | 195 | | | 195 | 482 | | 851 | | | | 195 | 150 | | 324 631 1181 94 | | 606 | 5 | | |
| Fraxinus americana | 781 741 382 417 13 1135 908 608 | | 741 1254 1256 | 755 417 1135 908 236 | 1333 | 1333 747 | 226 72 | | | | 13 | 1135 446 | | 1181 809 | | | | | 796 608 1268 1254 1257 1256 |
| Fraxinus nigra | 382 | | | | 12 | | | | | | | 710 446 | | | | | | | |
| Fraxinus pennsylvanica | 394 238 509 1332 465 321 | | | 509 1332 841 | | 1226 306 795 605 558 | 703 568 569 | | 963 | | 238 15 703 | 650 | 1323 963 | 748 321 809 | 857 871 131 | | 797 | 1175 | 171 795 650 963 |
| Gleditsia triacanthos | 238 509 465 781 782 1252 195 321 211 | | | 128 195 15 841 | 482 1335 174 104 | | 131 | | | 1335 | 238 195 15 | | | 782 321 | 857 1335 871 318 | | 797 | | |
| Gymnocladus dioicus | 781 782 | | | | | | | | | | | | | 782 | | | | | |
| Hamamelis virginiana | 548 | | | | | | 72 | | | | | | | | | 448 | 792 | | |
| HARDWOOD | 460 201 | | 439 462 | 460 375 | 263 | 991 | | 462 1241 | | | | | | | 839 1209 | 91 | | 615 718 | 1209 1241 |

| | | | | | | | | | | : | Subject | | | | | | | | | |
|-------------------------|---|----|-----|---|------------|--|-------------|------|------|-----|---------|-------------------|--|-----------------------|--------------------------|--------------|----------------------|---|----------------------------|---------------------|
| Species | IA,B | IC | ID | IE- | IE- 1a | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- | IIIA- 2 | IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | IIIB– 2bii | | IIIB- 2biv | IV |
| HARDWOOD (con't) | 382 375 646 | | 991 | 646 1021 991 | | | | 911 | | | | | | | 1126 | | | 10 91 | | |
| Ilex opaca | | | | 502 | 482 | 416 605 | | | | | | | | 1222 | | | 792 | | | 416 502 |
| Juglans spp. | | | | | | | | | | | | - | 6 | 631 1181 | | | | | | |
| Juglans cinerea | 781 1332 465 | | | 1332 | | | | | | | | | 611 824 458 1323 | . | | | 792 | | | |
| Juglans nigra | 781 547 394 1178 128 1332 465 15 413 211 235 924 1304 | | | 544 128 1332 90 15 235 1304 | 174 | 128 1268 | 1288 | 462 | | | 924 | 753 | 3 824 291 925 90 458 1051 175 1323 25 1287 410 | | 857 871 319 | | 792 | 27 45 25 | 4 | 235 1304 1268 |
| Juniperus spp. | 610 | | | | _ | | | | | | | | | 1181 | | | 792 1347 | | 6 | |
| Juniperus virginiana | 1178 509 1332 195 336 71 44 184 | | | 1178 1332 195 336 43 | 906 482 | 71 44 | 906 1213 | | | | 195 | • | | | 906 857 871 923 | | | 100 43 93 93 68 68 57 53 | 5 2 3 1 5 4 | |
| Kalmia latifolia | 711 726 | | | | 711 | | | | | | | | | 155 | | | 792 | . 25 | 9 | |
| Larix spp. | 401 978 | | | 401 | | | 1300 | | | 724 | 978 | 3 252 150 | |) 631 3 1181 94 | | 1245 1154 | | 25 99 | | |
| Larix laricina | 946 349 67 382 68 71 297 99 | | | 946 90 68 297 99 1176 1340 | 730 311 | 744 747 253 730 421 71 99 311 | | | | | | 730 710 446 |) | 779 1181 | | 708 | 1147 1148 1144 | 119 | | 1176 |
| Liquidambar styraciflua | 244 336 | | | 244 336 | | 558 1226 | | 926 | | | | 150 1199 | | 1222 735 | 1210 | | 792 | 135 | | 795 1199 |

| | | | | | | | | | S | ubjec | t | | | | | | | | | | |
|------------------------------------|---|--|--|--|--------------------------|---|-----------|------|---|-----------|-----------|------|-------------------|----------------------------------|-------------------|--------------|---------------|------|---------------------------------------|--------------|----------------------------|
| Species | IA,B IC | C ID | IE 1 | IE- la | IE- 2 | IIA | IIB- 1 | IIB- | IIC | IIIA 1 | IIIA 2 | – II | IIA- 3 | IIIB 1 | IIIB- 2a | IIIB- 2bi | IIIB- 2bii | | | IIIB 2biv | IV |
| Liquidambar styraciflua (con't) | 6 178 651 | | 1259 555 | 1226 482 769 566 568 551 557 558 174 553 552 | 178 651 555 605 | 521 735 | | | | | 38 42 | | | 187 479 809 186 | | | | | 175 207 996 258 | | |
| Liriodendron tulipifera | 394 1178 244 6 734 669 1114 | 669 163 558 267 268 269 | 8 872 8 1335 7 236 8 567 742 | 667 482 768 769 1335 1115 | 268 | 664 851 226 6 72 870 892 890 1266 891 669 1288 | 268 | | 1115 163 267 268 269 724 | | | | 601 567 268 | 631 324 155 1222 642 | 634 871 131 | 60 | 6 79 | | 594 633 681 685 256 54 | | 796 734 1266 1257 |
| Maclura pomifera | 509 195 | | 174 | 481 174 | | | | | | 195 | 5 | | | | | | | | | | 171 |
| Magnolia spp. | | | | | | 1300 | | | | | | | | | | | | | | | |
| Magnolia acuminata | | | | | | 266 | | | | | | | | | | | | | | | |
| Morus spp. | | | | | | | | | | | | | | | 871 | | | | | | 171 |
| Morus alba | 781 509 195 | | 195 | | | | | | | 195 | 5 | • | | | | | | ., 4 | | | |
| Morus rubra | | | | 481 174 | 605 | 1300 | | | | | | | | | | | | | | | |
| Nyssa aquatica | | | | 482 566 568 558 556 314 174 553 552 | 558 | 568 314 | | | | | | | | | | | | | | | |
| Nyssa sylvatica | 546 727 | | 727 | 482 569 315 551 556 554 557 558 | | 557 558 | | | | | | | | | | | | | | | |

| | | | | | | | | | | | Subject | | | | | | | | | |
|-------------------------|--|-----|--------------------------|--|--------------------------------|---------------------------------|-----------------------------------|------|------|-----|----------------------------------|------------|--------------|--|--|--|--|----------------------|--|--|
| Species | IA,B | IC | ID | IE- 1 | IE- 1a | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- 1 | IIIA- | IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | IIIB– 2bii | | IIIB- 2biv | IV |
| Nyssa sylvatica (con't) | | | | | 174 552 | | | | | | | | , | | | | | | | |
| Ostraya virginiana | | | | | 482 | - | 226 72 | | | | | · | | 748 | | | | | | |
| Picea spp. | 1178 524 559 437 | | | 524 212 90 437 815 | | 253 212 | 1300 | | | | 429 | 150 |) | 922 1181 815 | | 1245 708 | | 192 | | 429 108 |
| Picea abies | 999 375 349 649 755 700 509 417 619 930 1083 401 117 543 1023 978 592 | 627 | 999 462 700 543 | 999 450 700 90 417 301 619 301 401 627 117 799 | 349 700 731 558 53 | 700 939 | 1 1302 712 | 700 | | 627 | 1 755 619 978 | | 700 | 7 512 450 812 700 771 324 1168 922 1181 680 | 834 871 923 131 | 1246 340 1085 | 792 1147 | 594 1122 293 | | 1151 416 755 352 1266 1150 483 |
| Picea glauca | 946 229 67 68 1071 600 874 873 86 1225 1151 375 1152 608 343 46 1023 978 344 | | | 29 1208 386 1224 873 1225 1080 1152 343 1139 516 68 299 344 | 600 | 375 746 864 747 214 | 891 978 1086 1359 593 | | | | 1071 86 1080 978 344 | 446 344 | ; | 516 | 1247 1271 1123 1208 759 1273 901 | 1244 1245 1273 1274 1152 1246 1154 | 7 459 1144 5 1143 3 1146 4 | 347 1271 3 560 | 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | 1193 1033 875 1150 876 1151 1152 608 361 299 502 |

| | | | | | | | | | | 5 | Subject | | | | | | , | | | |
|----------------------|---|-----|----|---|-------------------|------------|----------------------|-----------|------|--------------------|--|------------|-------|--|---|----------------------------|--|---|---------------|--|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB– 1 | IIB- | IIC | IIIA- | IIIA- 2 | IIIA- | IIIB– 1 | IIIB- 2a | IIIB- 2bi | IIIB– 2bii | | IIIB- 2biv | IV |
| Picea glauca (con't) | | | | | | | | | | | | | | | | | | 1087 478 880 | ; | |
| Picea mariana | 946 67 382 68 719 564 873 402 46 978 1316 | | 68 | 421 68 731 10 1176 564 311 873 784 1080 720 1044 402 1052 49 607 47 48 1359 | 12 873 | 747 253 | 978 1359 1130 |)) | | 731 1286 | 1080 978 | | | 1181 971 974 | 759 | | 1144 | | | 1176 607 1316 |
| Picea rubens | 859 382 1343 | | | 859 | | 1111 | | | | | | 446 | | 859 783 1181 | | | 1147 1148 1144 1143 | 1085 | ' | |
| Pinus spp. | 498 497 182 849 610 617 | | | 524 497 301 975 617 815 | 273 | 253 849 | | | 1090 | 1073 | 429 | 150 849 | | 498 497 922 631 1090 1091 975 1181 701 849 1092 138 617 815 94 | | 1245 708 | 5 792 3 1145 | | ! ! | 429 108 |
| Pinus banksiana | 946 1118 2 229 230 509 1332 67 68 1124 608 46 377 396 422 | 835 | | 946 1118 860 230 1124 1301 297 699 731 1176 10 873 1079 1080 47 | 1072 12 873 | 747 | 1300 1302 1216 | | 1100 | 699 1215 731 | 599 2 230 422 1071 1079 1080 | 690 446 | | 775 843 280 1181 701 812 1283 516 810 | 767 1123 915 916 385 217 | 1244 448 1043 916 | 916 674 459 74 1144 676 1143 | 346 952 233 1122 387 293 | | 599 1193 1009 1124 1033 1215 1216 1176 608 361 1316 217 |

| | | | | | | | | | | | Subjec | t | · | | | | | | | |
|-------------------------|--|----|---------------------------|----------------------------------|--------------------|--|---|-------------------|------|----------|---|---|---|--|---|--|--|--|---|---------------------------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- | IIIA- | - IIIA- | - IIIB 1 | IIIB- 2a | IIIB- 2bi | IIIB- 2bii | | IIIB- 2biv | IV |
| Pinus banksiana (con't) | 719 767 1071 623 297 305 873 1079 1200 375 865 1065 | | | 48 516 1359 377 1065 | | | | | | | | | | | | | | 915 916 1206 1205 900 615 288 674 675 84 | 5 5 5 5 8 8 8 | |
| Pinus echinata | 789 244 653 578 6 656 254 1259 1048 1227 597 127 124 | 22 | | 653 | 580 1306 174 | 270 1165 939 733 1129 | 666 656 1302 | | | 1000 724 | | | | 578 1006 810 1354 188 189 1181 802 816 803 804 817 818 805 812 820 1016 806 679 73 807 110 809 | 5 580 1194 254 3 451 1352 1352 1354 5 533 5 534 5 597 5 521 | 270 4 4 1 1 7 3 4 3 4 7 | | 2 1194 2 254 2 517 3 1353 3 1356 5 1007 5 1354 | | 275 796 739 738 110 |
| Pinus pungens | | | | | | - | | | | | | | | 1181 | | | | | | |
| Pinus resinosa | 1330 498 4 295 382 1124 417 1071 836 1207 | | 1293 837 525 381 | 29 4 295 1124 665 | 873 | 416 71 853 836 744 1293 657 747 | 1001 665 1301 673 360 712 661 | 6 663 740 3 | 2 | 3 121 | 107 1136 762 1086 1126 714 882 978 | 6 5 2 120 0 113 0 65 4 80 2 44 | 2 17 60 71 77 71 66 71 69 2 01 39 66 4 | 5 322 2 324 5 323 4 1001 5 672 | 2 1008 1141 3 587 1051 2 175 3 1129 834 1123 | 3 1010 1 708 7 1244 1 1066 5 448 9 1043 4 916 3 674 | 915 3 1148 4 1143 6 1144 8 916 3 674 6 74 4 676 | 399 3 1087 3 1246 4 940 5 572 4 489 4 346 5 952 | 1193 1009 1338 1124 1301 796 1033 2 1215 | |

| | | | | | | | | | | S | Subject | | | | · · | | | , . | | |
|------------------------|--|-----|-----|--|-----------|----------|--|-----------------|-----------|-----------------|------------|------------|-------|--|-------------|---|-------------------------|--|--|---------------------------------|
| Species | IA,B | IC | ID | IE- 1 | IE- 1a | IE- 2 | IIA | IIB- | IIB- 2 | IIC | IIIA- 1 | IIIA- 2 | IIIA- | IIIB– 1 | IIIB- 2a | IIIB- 2bi | IIIB– 2bii | IIIB– 2biii | | IV |
| Pinus resinosa (con't) | 185 1290 403 978 378 376 380 762 873 1291 1083 1200 1292 375 1293 307 1294 608 45 1120 714 381 1297 377 | | 377 | 1207 51 185 376 873 1293 657 389 1080 450 116 49 716 381 362 377 1297 378 | | | 313 657 978 593 1132 1133 | 7 3 3 3 2 2 3 3 | | | | 378 380 | 1133 | 1296 807 1297 39 34 831 1277 1295 1149 1294 450 395 812 1292 887 640 1298 593 973 38 36 35 810 | | 353 626 397 1246 398 376 1085 | 3 6 7 7 6 8 8 9 6 6 5 5 | 271 233 347 594 1338 387 293 1198 916 615 616 674 32 30 932 933 207 204 973 1068 478 879 917 685 1170 1169 272 587 1269 266 1204 560 920 846 158 1007 915 582 1024 31 1081 680 1277 537 830 837 | 836 352 1266 608 572 45 1127 882 11 381 38 | |
| Pinus rigida | 1285 440 724 789 1329 441 | 440 | | 860 789 727 21 441 | | | 1220 1217 1329 | | 1100 | 723 78 80 | 789 | 789 446 | | 322 323 789 325 1181 1220 1049 812 810 | 383 | 207 | 1012 1013 | 323 | | 796 383 739 572 738 |

| | | | | | | | | | | | Subject | | | | | | | | | |
|------------------|---|-----|--|---------------------------|--|--------------------------------------|--|-----------------------------------|------|-------------------|------------|--|----------------------|-------------|---|--|--|---|---|--|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- | IIB- | IIC | IIIA- 1 | IIIA- | IIIA- | IIIB- 1 | IIIB- 2a | IIIB– 2bi | IIIB- 2bii | IIIB– 2biii | | iv |
| Pinus serotina | | | | 727 78 79 80 | 580 | 1129 | | | | | | | | 1181 810 | | 1108 | 1012 1013 1015 | | | |
| Pinus strobus | 406 946 349 859 1119 498 641 497 760 851 382 68 417 1071 1109 918 583 185 403 930 1343 762 117 999 375 608 543 978 896 724 | | 1119 641 68 375 543 740 | 349 859 460 1119 | 90 1072 765 1131 12 624 1202 | 416 758 71 1129 919 9 | 665 775 776 1090 403 870 1213 978 | 1073 737 736 1202 740 | 1093 | | 1071 | 710 690 150 444 151 445 152 659 148 149 | 390 1287 | 783 | 68 1141 1172 66 1131 1123 560 1078 6403 871 896 1287 923 6614 131 1126 | 1172 1010 1210 1244 1043 1249 1248 1249 1248 1249 1248 1249 1248 1249 1248 1249 1248 1249 1248 1249 1248 1249 1249 1249 1249 1249 1249 1249 1249 | 2 428 1347 1147 1012 3 1148 1013 1015 | 346 322 1339 1251 231 90 | 6 160 2 30 6 80 933 327 681 685 56 478 2 160 3 32 5 15 6 32 9 32 5 15 6 33 4 204 3 364 6 896 6 644 6 896 6 644 | 1193 1009 1338 1033 796 1210 1215 918 352 875 608 1257 877 |
| Pinus sylvestris | 649 700 498 509 619 728 317 1334 902 401 999 117 179 608 | 180 | 700 1289 | 317 1334 731 386 | 999 53 558 799 | 53 308 379 | 1090 395 978 799 177 | 700 1289 3 | | 731 627 724 | 1249 |) | 5 1051 712 390 | 498 | 3 266 3 871 3 3 4 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 | 1043 625 | 1347 5 1147 3 1148 5 1144 7 1143 5 5 5 8 | 594 1122 3 293 4 922 | 1 | 416 1334 902 179 608 572 287 |

| | - | | | | | | | | | , | Subject | | | | | | | | | |
|--------------------------|---|--|--|--|---|--|---|--|------|------------------|------------|---------------------|------------|--|---|----------------------------|--------------------------|---|---------------|--|
| Species | IA,B | IC | ID | IE- 1 | IE la | IE- 2 | IIA | IIB | IIB- | IIC | IIIA | IIIA- | IIIA- | IIIB- | IIIB- 2a | IIIB 2bi | IIIB- 2bii | IIIB- 2biii | IIIB- 2biv | IV |
| Pinus sylvestris (con't) | 375 370 1023 308 543 978 377 724 | | | | | | - | eter. | | | | | | | | | | 162 117 572 1246 1085 785 | 7 } } | |
| Pinus taeda | 1360 244 653 578 245 | 430 193 722 1134 102 1344 5 194 | 960 578 82 968 1227 119 669 121 122 124 | 960 336 42 665 938 21 82 1259 254 1048 451 78 | 906 580 482 430 172 194 174 199 624 1230 1306 1229 1231 | 146 1227 431 433 722 1165 416 1129 1095 1352 770 | 665 1341 673 656 861 668 82 671 172 870 24 892 | 477 431 432 433 505 504 965 508 | | 80 163 724 | 894 798 | 1053 446 1054 | 3 567 5 | 1017 242 239 110 821 807 1327 751 813 812 322 323 578 1341 672 668 1260 325 476 1227 1354 975 188 1181 189 977 810 257 809 822 1159 75 833 94 240 241 808 1306 802 816 962 | 595 938 906 580 1194 254 517 451 168 290 1227 1354 255 533 213 534 521 749 116 751 292 506 | 1250 530 581 1262 |) 488) 487 l 1227 | 3 595 7 1194 7 264 3 254 1 517 2 325 3 754 5 1356 4 158 | | 416 796 1094 1348 1095 463 118 1350 1229 119 1201 1157 739 432 1158 506 1262 1263 110 102 1230 505 157 126 504 283 500 764 56 943 702 738 1261 844 507 508 203 1017 |

| | | | | | | | | | | | Subject | | | | | | | | | |
|-----------------------|---|--------------|-----|--------------------------------|---------------------------|---------------------------|------------|-----------|-----------|-----|------------|------------|-------------|---------------------|---------------------------|--------------|---|--------------------------------|---------------|---------------------------|
| Species | IA,B | IC | ID | IE- 1 | IE- 1a | IE- 2 | IIA | IIB- 1 | IIB- 2 | IIC | IIIA- 1 | IIIA- 2 | IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | IIIB– 2bii | | IIIB- 2biv | IV |
| Pinus taeda (con't) | | | | | | | | | | | | | | | | | | 1088 55 899 18 688 | | |
| Pinus virginiana | 629 | | | 629 21 476 567 816 | 174 | | | 477 | , | | 894 | | 567 415 | | 383 415 | } | 792 1147 1012 1148 1013 1015 | 1007 612 932 514 | | 775 |
| Platanus spp. | | | | | | | | | | | | | | 1181 | 839 |) | | | | |
| Platanus occidentalis | | | 418 | 128 568 | | 795 568 418 128 | 569 | 1 | | | 870 | | | 94 809 | | | | 1125 681 685 | | 795 418 |
| Populus spp. | 869 1196 1028 501 | 1196 1028 | | 1300 93 | | | | | | | 14 | 150 |) 282 14 | | 839 1123 871 501 | | 3 792 | 1269 310 | | |
| Populus alba | 405 | 405 | | 755 405 841 | | | - | | | | | | | 1181 | 871 | | | | | |
| Populus balsamifera | 946 176 869 465 1070 298 | | 176 | 946 | 12 | 176 298 948 1040 | | | | | | | | 1181 | | | | | | |
| Populus deltoides | 1305 1332 1236 1028 358 64 | 1299 | 462 | | 455 1333 482 565 | 795 | 568 799 | | | | 15 | | | 748 1221 1223 | 1235 871 | | 3 797 | 32 863 | 1235 | 171 795 1341 419 |

| | | | | | | | | | | 5 | Subjec | t | | | | | | | | |
|--|---|----|------|---|-----------|---|------------------|------------------------|------|-----|-------------------|------------------------|-------------|---|------------------|--------------|---------------|---|---------------|-----|
| Species | IA,B | IC | ID | IE- 1 | IE- 1a | IE- 2 | IIA | IIB– 1 | IIB- | IIC | IIIA- 1 | IIIA- 2 | - IIIA- | - IIIB– 1 | IIIB 2a | IIIB- 2bi | IIIB– 2bii | | IIIB- 2biv | IV |
| Populus deltoides (con't) | 501 | | | 1341 | | 427 1040 | | | | | | | | | | | | | | |
| Populus deltoides var. occidentalis | | | | | | 128 | | | | | | | | | | | | *************************************** | | |
| Populus grandidentata | 335 1070 356 1349 77 793 | | | 356 77 1049 | | 357 335 76 355 356 303 304 1349 563 77 1049 793 | 226 72 661 | | | | 77 | 7 69 30 44 79 | 4 6 | 1223 807 810 | 7 | | | | | |
| Populus nigra var. italica | 1061 | | 1062 | | | 1063 1062 | | | | | | | | 1181 | 87 | 1 | | | | |
| Populus tremuloides | 176 | | 176 | 296 109 356 77 1049 424 425 1164 | 12 | 1039 1038 1035 1042 523 1163 793 424 1034 357 425 61 62 1045 645 1069 176 296 1027 1040 1036 103 522 1041 1164 794 76 191 348 356 355 303 449 | 226 | 5 1163 1041 1042 | | 724 | 762 77 1164 | 7 69 | 0 4 6 | 1181 680 1049 1223 807 810 |)) 3 7 | 4 | | 104: 680 135: 100: |) 7 | 348 |

| | | | | | | | | | | | Subject | t | | | | | | | | |
|-----------------------------|---|----|------------|--|----------------------------------|--|--------------------|---------------------|-----------|-----|---------|---|----------------------------|---|---|------------------|---------------|--|---|-----------------------------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- | IIB- 2 | IIC | IIIA- | IIIA- 2 | IIIA- | IIIB- | IIIB- 2a | IIIB- 2bi | IIIB- 2bii | | IIIB- 2biv | IV |
| Populus tremuloides (con't) | | | | | | 304 563 190 77 1049 948 | | | | | | | | | | | | | | |
| Prunus spp. | 547 | | | 898 | | | | | | | | | 60 | l | | 448 | 3 792 | 2 1269 |) | |
| Prunus pensylvanica | 382 | | | | | | | | | | | | | | | | | | | |
| Prunus virginiana | 229 1332 | | | | 1332 | | | | | | | | | | | | | | | |
| Prunus serotina | 382 762 181 | | | 872 72 | | | 226 72 713 | | | | 762 | 2 710 440 | | 809 |) | | | | | |
| Quercus spp. | 547 652 1178 653 245 182 375 184 999 711 1285 1329 1212 | | 375 462 | 652 182 999 711 566 | 481 482 174 | 838 711 795 | 852 1329 145 | 1 | 2 | | 245 | 6 69 69 32 15 119 96 44 | 1 1323 2 3 7 7 | 3 324 | 966 1141 1210 573 1209 871 | 5 l) 3 | 3 792 | 2 652 222 690 691 1269 1210 208 1345 329 527 1197 689 207 28 169 | 2 0 1 3 5 5 7 7 7 7 8 | 966 329 1268 795 1209 |
| Quercus alba | 394 652 1178 244 245 336 1056 660 414 220 156 1135 854 1285 183 706 1329 787 | | 780 | 652 1178 244 336 414 220 156 1135 706 707 | 906 156 1060 104 707 | 733 | 226 | 124: 124: 78: | 2 | 787 | 245 | 5 98 113: 69: 119: 61: 446 | 5 3 5 | 779 780 220 1181 239 240 241 810 | 906 573 871 | 5 3 | 792 | 2 272 98 169 208 693 1173 613 1175 454 1067 1240 | 3 3 3 3 3 3 4 7 | 1241 1242 |
| Quercus bicolor | 220 | | | 220 | 455 104 | | | | | | | | | 220 |) | | | 169 |) | |

| | | | | | | | | | | | Subject | t | | | | | | | | |
|--------------------------------------|---|----|----|--------------------------------|---|------------|------------|------|-----------|-----|----------|--------------------------------|-------------|----------------------------|-------------|----------------|-----------------|--|---------------|------|
| Species | IA,B | IC | ID | IE- 1 | IE– la | IE- 2 | IIA | IIB- | IIB- 2 | IIC | IIIA- | IIIA- 2 | IIIA- | IIIB- | IIIB- 2a | - IIIB- 2bi | - IIIB- 2bii | | IIIB- 2biv | IV |
| Quercus coccinea | 1178 156 1285 183 1329 | | | 1178 156 | 156 | | 1329 | , | | | | 159 586 446 | 5 | | | | | 272 169 159 493 |) | |
| Quercus ellipsoidalis | 305 | | | | | | 611 | | | | | 98 693 329 659 446 | 3)) | | | | | 98 169 693 329 | | 329 |
| Quercus falcata | | | | | 481 174 | | | | | | | 159 |) | 1181 145 | | | | 169 159 | | |
| Quercus falcata var. pagodaefolia | | | | | 565 566 174 | | | | | | | | | | | | | 169 1104 | | |
| Quercus lyrata | 656 | | | | 481 906 482 174 | 656 | 656 | | | | | | | | | | | | | |
| Quercus macrocarpa | 544 1253 1332 509 274 13 211 305 403 854 | | | 544 128 509 274 15 | 1332 455 12 104 | 128 | 661 | | | | 15 13 | |) | 748 1253 326 1181 | • | 3 | | 272 690 691 98 169 693 881 | | |
| Quercus marilandica | 244 336 156 711 | | | 244 336 156 711 | 481 156 174 | 733 711 | | | 1325 | | | 1195 613 446 753 | 3 | | | | | 272 169 613 | | |
| Quercus muehlenbergii | 781 | | | | 174 | | | | | | | | | | | | | 169 | 1 | |
| Quercus nigra | | | | | 481 482 174 | | | | | | | 446 | 5 | · | | | | 169 208 1175 | | |
| Quercus palustis | 202 220 | | | 220 568 | 164 455 1333 482 565 568 566 569 315 104 | 568 | 568 569 | | | | | | | 220 810 | | | 792 1337 | 2 272 7 169 | | 202 |
| Quercus prinus | 1318 1319 | | | 1135 | 481 906 | 1319 | 851 906 | | | | | 1135 446 | 5 | 1181 | 900 | 5 | 1147 1148 | | | 1319 |

| | | | | | | | | | | 5 | Subject | | | | | | | | | |
|------------------------|-------------|----|-----|-------------|-------------|--------------|------|--------------|-----------|---------|------------|-------------|-------------|--------------|-------------|--------------|---------------|------------|--------|-------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB– 1 | IIB- 2 | IIC | IIIA- 1 | IIIA- 2 | IIIA- | IIIB– 1 | IIIB- 2a | IIIB– 2bi | IIIB– 2bii | | | IV |
| Quercus prinus (con't) | 437 1135 | | | | 909 | | | | | | , | | | | | | | | | - |
| Quercus rubra | 781 | | 462 | 133 | 15 | | 664 | | | | | 1182 | | 779 | | | | | 2 1235 | |
| | 652 1178 | | 705 | 652 1178 | 906 | 128 1248 | 756 | | | | 13 | | | 220 | | | | 272 | | 1182 |
| | 544 | | 343 | 544 | 130 | 705 | | 1241 1242 | | | 762 | 693 1346 | | 1181 1049 | | | | 169 693 | | 1336 526 |
| | 244 | | | | 1177 | 219 | 906 | | | | | 1336 | | 810 | | | | 1346 | | 608 |
| | 382 | | | | 1060 | 359 | 799 | | | | | 159 | | | | | | 159 | | 1214 |
| | 336 | | | 336 | | 886 | | | | | | 446 |) | | | | | 113 | | 219 |
| | 417 | | | 417 | 104 | | | | | | | | | | | | | 454 | | |
| | 1056 15 | | | 455 156 | 799 | | | | | | | | | | | | | 1067 | | |
| | 13 | | | 220 | | | | | | | | | | | | | | 1240 |) | |
| | 979 | | | 1135 | | | | | | | | | | | | | | | | |
| | 156 | | | 117 | | | | | | | | | | | | | | | | |
| | 220 | | | 908 | | | | | | | | | | | | | | | | |
| | 211 | | | 705 | | | | | | | | | | | | | | | | |
| | 1135 762 | | | 709 707 | | | | | | | | | | | | | | | | |
| | 117 | | | 706 | | | | | | | | | | | | | | | | |
| | 854 | | | 700 | | | | | | | | | | | | | | | | |
| | 908 | | | | | | | | | | | | | | | | | | | |
| | 608 | | | | | | | | | | | | | | | | | | | |
| | 543 706 | | | | | | | | | | | | | | | | | | | |
| Quercus stellata | 1178 | | | 1178 | 481 | 733 | 1328 | 345 | | 345 | | 446 | <u></u> | | | | - | 169 |) | |
| | 244 | | | 244 | 156 | | | | | | | 753 | | | | | | | | |
| | 336 156 | | | | 1060 174 | | | | | | | | | | | | | | | |
| Quercus velutina | 244 | | | | 1060 | 733 | | 1241 | | | | 691 | | | 1209 | | | 272 | | |
| | 336 | | | 336 | 104 | | | 913 | | | | 1195 | | 1049 | | | | 691 | | 1241 |
| | 854 | | | | | | | | | | | 159 | | 810 | 913 | | | 169 | | |
| | | | | | | | | | | | | 613 446 | | | | | | 159 613 | | |
| | | | | | | | | | | | | 770 | , | | | | | 493 | | |
| | | | | | | | | | | | | | | | | | | 1240 | | |
| Rhododendron spp. | 726 | | | | | | | | | | | | | | | | 792 1347 | | , | |
| Rhododendron maximum | 1284 | | | | | | | | | | | | | | | • | | 259 |) | |
| Rhus spp. | | | | | | | | | | | | | | | | 448 | 3 | | | |
| Rhus copallina | 335 | | | 335 | | 335 | | | | | | 446 753 | | | | | | | | |
| Rhus glabra | | | | | | | | | | · · · - | | 753 | | | | | | | | |
| Robinia pseudoacacia | 509 | | 766 | 133 | 481 | 133 | 72 | | 1323 | | 133 | 446 | 282 | . 227 | 857 | 606 | 5 1147 | 7 272 | , | 416 |
| r | 195 | | | 1155 | 174 | 416 | 703 | | 1100 | | 195 | | 925 | 588 | 521 | | 1148 | | | 171 |
| | 465 | | | 195 | | 227 | | | | | 13 | | 227 | | | | | | | 1215 |
| | 13 | | | 72 | | 704 | | | | | 703 | | 588 | | | | | | | 1232 |
| | 660 543 | | | 766 1317 | | 1155 1138 | | | | | | | 589 1323 | | | | | | | 1317 |
| | 343 | | | 841 | | 1130 | | | | | | | 1323 | 831 | | | | | | |

| | | | | | | | | | | S | ubjec | t | | | | | | | | |
|--------------------|--|----|------------------|--|--------------------------------|---|-----------|-----------|-----------|--------------------|-------|--------------------------|--------|---------------------------|------------|--------------|---------------|--|-----------------------|------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- 1 | IIB- 2 | IIC | IIIA- | IIIA- 2 | IIIA- | IIIB- | IIIB 2a | IIIB- 2bi | IIIB– 2bii | | IIIB- 2biv | IV |
| Salix spp. | 1139 501 723 1178 1236 | | | 411 250 731 93 1139 1115 163 1236 | 250 558 52 481 455 | 218 1236 | 163 | | 52 | 1115 163 724 | | | | 1181 | 839 | 448 | 3 792 | 207 | 7 | |
| Salix alba | 509 897 | | 462 | | 558 427 | 427 | | 462 | | | | | | ****** | 871 | | 797 | 996 | 5 | |
| Salix nigra | 724 | | | 95 568 1115 | | | | | | 1115 724 | | | | | | | | | - | 795 |
| Sassafrass albidum | 547 335 415 | | | | 481 482 174 104 | | | | 415 | | | 440 | 5 41: | 5 | | | | | | |
| Taxodium distichum | 825 312 1178 198 673 309 | | | 825 312 198 309 | 1178 198 | 694 198 1226 | | | | 673 | | | | 94 | 825 906 | | | 1007 996 | | 416 314 |
| Thuja spp. | | | | · · · · · · | | 253 | | | | | | 150 |) | | 923 | 708 | | | | |
| Thuja occidentalis | 859 68 70 382 69 71 276 884 437 469 375 878 | | 68 70 375 | 860 | | 746 747 416 939 69 71 884 | | | | | | 710 658 659 440 | 8 9 | 859 884 | | , | 792 1144 | 2 884 1269 1206 1024 192 32 30 1089 |) 5 1 2 2 | 416 201 |
| Tilia spp. | | | | | 481 1335 | | | | | 1335 | | 150 |) | 1181 | 1335 | | | | | |
| Tilia americana | 781 544 1332 382 211 | | 462 780 59 | 1332 | 12 58 1177 174 | 745 59 | 226 72 | | | | 15 | 5 710 440 | | 779 780 1181 905 | | 448 | | 1269 |) | 59 608 |

| | | | | | | | | | | S | ubjec | t | | | | | | | | |
|-------------------------|--|----|-----------|-------------------------------|--------------------------------|--------------------------|-------------------|-------------|-----------|------|------------|--|-------|-------------|--|--------------|---------------|--|-------------|-------------------|
| Species | IA,B | IC | ID | IE- 1 | IE– 1a | IE- 2 | IIA | IIB- | IIB- 2 | IIC | IIIA- 1 | IIIA- 2 | IIIA- | IIIB- 1 | IIIB- 2a | IIIB- 2bi | IIIB- 2bii | | | IV |
| Tilia americana (con't) | 58 608 | | | 57 | | | | | | | | | | | | | | | | |
| Tsuga spp. | | | | | | | | | | | | 150 | | 1181 | 839 | | | | | |
| Tsuga canadensis | 407 1178 68 382 1343 930 375 724 | | 68 375 | 407 1178 68 72 | | 747 1156 71 | 72 1131 | | | 724 | | 903 893 658 659 285 446 | | 407 1181 | | | 3 792 1347 | |) | 1082 |
| Ulmus spp. | 223 | | | | | | | | | | | 150 | 282 | 324 1181 | | 448 | 3 792 | | | |
| Ulmus alata | 781 238 1332 509 201 225 224 382 202 1200 1050 | | | | 481 778 482 174 | | | | | | | | | | | | | | | |
| Ulmus americana | | | 462 | 201 90 224 665 15 | 455 778 482 306 12 | 747 416 755 568 | 1001 665 72 | l 5 2 | 2 | 1335 | 238 | | 415 | | 1235 1209 1141 1210 993 1335 415 | | 797 | 1024 1205 528 32 883 277 995 | 3 2 3 | 416 224 202 |
| Ulmus rubra | | , | | 72 | | | 72 | 2 | | | | | | | | | | | | |
| Ulmus thomasii | 608 | | | | 12 | | | | | | | | | | | | | | | 608 |





